



On the Endomorphism Ring of a Noetherian Chain Module

Hanan A. Alolaiyan^{1*} and Ahmad M. Alghamdi²

¹Department of Mathematics, Faculty of Sciences, King Saud University, Riyadh, Saudi Arabia.

²Department of Mathematical Sciences, Faculty of Applied Sciences, Umm Alqura University, P.O.Box 14035, Makkah 21955, Saudi Arabia.

Authors' contributions

This work was carried out in collaboration between both authors. Author AMA suggested the idea of the paper, and wrote the first draft of the manuscript, managed the literature searches as well as revised the paper several times. Author HAA modified the first draft, wrote the proofs, revised the paper as well as contacted the publishing agency. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/BJMCS/2016/23383

Editor(s):

(1) Raducanu Razvan, Department of Applied Mathematics, Al. I. Cuza University, Romania.

Reviewers:

(1) Burcu Nisanci Turkmen, Amasya University, Turkey.

(2) Leonardo Simal Moreira, University Centre of Volta Redonda, Volta Redonda, Brazil.

(3) Ali Mutlu, Celal Bayar University, Turkey.

Complete Peer review History: <http://sciencedomain.org/review-history/13052>

Received: 28th November 2015

Accepted: 13th January 2016

Published: 23rd January 2016

Original Research Article

Abstract

The aim of this paper is to further develop the theory of the well-known Schur's lemma on endomorphism rings for application to wider structures. We show that for a commutative Noetherian chain ring R and Noetherian chain modules M and N , the left R -module $\text{Hom}_R(M, N)$ will be a chain module. Then, we conclude that the endomorphism ring $\text{End}_R(M)$ of the Noetherian chain module M over such R is a Noetherian chain ring. As a special case, we consider that the dual R -module $M^* = \text{Hom}_R(M, R)$ of the Noetherian chain R -module M is a Noetherian chain module.

*Corresponding author: E-mail: holayan@ksu.edu.sa;

Keywords: Chain rings; chain module; Noetherian ring; Noetherian module; endomorphism ring; Schur's lemma.

2010 Mathematics Subject Classification: 06B99, 13F05, 13E05, 13C99.

1 Introduction

The chain ring is a generalization of the concepts of the division ring and commutative valuation rings. Several approaches, both algebraic and geometric, have been employed in studying this concept (see [1]). This research will be supported by Alolaiyan's thesis " *The structure of chain rings*" [2], which is focused on chain rings, generalized chain rings and their relationship with Galois groups.

In this paper, we shall recall a parallel concept for a module over a chain ring. A chain module indicates a module with a unique series of submodules. In other words, even if two submodules of a module M are arbitrary, they should be comparable. The main objective is to investigate the structure of the endomorphism ring of a Noetherian chain module. This investigation is followed by a study of the behavior of a chain ring under the assumptions that the module under construction is a Noetherian chain module and that the coefficients come from a commutative Noetherian chain ring.

This paper is organized as follows. In Section 2, we discuss the set of ideals in a commutative Noetherian chain ring and the set of submodules in the chain modules. Then, in Section 3, we present the main results for the endomorphism ring of a Noetherian chain module and the dual of a chain module. In Section 4, we give a brief conclusion about this work.

We assume that all rings are associative commutative with identity and that all modules are unitary modules. By an R -module (R -submodule), we refer to the left R -module (R -submodule), where R is a ring with identity and is a commutative Noetherian chain ring. For more information about chain rings and modules, the reader can consult the survey article [1] and the two books [3, 4]. For new results in the notion of endomorphism rings, the reader can see [5, 6, 7, 8]. See also [9, 10].

2 Preliminaries and Definitions

In this section, we recast some definitions and standard facts and describe certain properties that characterize the set of ideals and submodules in specific algebraic structures, which are then used to reach our conclusion.

Definition 2.1. A ring R is called a *left chain ring* if for all $a, b \in R$, we have either $Ra \subseteq Rb$ or $Rb \subseteq Ra$. Similarly, R is called a *right chain ring* if for all $a, b \in R$, we have either $aR \subseteq bR$ or $bR \subseteq aR$. The ring R is called a *chain ring* if its ideals are comparable.

Definition 2.2. A left (right) module M over a ring R is called a **left (right) chain module** if any two left (right) submodules of M are comparable.

Definition 2.3. Let R be a commutative ring, and let M, N be R -modules. Then, the module of all module homomorphisms from M to N is called the *homomorphism R -module* from M to N and is denoted by $Hom_R(M, N)$.

Definition 2.4. Let R be a commutative ring, and let M be an R -module. Then, the ring of all module homomorphisms from M to M is called the *endomorphism ring* of M and is denoted by $End_R(M)$.

Definition 2.5. Let R be a commutative ring, and let M be an R -module. Then, the module of all module homomorphisms from M to R is called the **dual module** of M and is denoted by $M^* = Hom_R(M, R)$.

Proposition 2.1. Let R be a commutative Noetherian chain ring. Then, its ideals form a unique series of the form $R \supset \langle \theta \rangle \supset \langle \theta^2 \rangle \supset \dots$, for some $\theta \in R$.

Proof. We will show that any Noetherian chain ring is a principle ideal ring. As R is Noetherian, all of its ideals are finitely generated. Now, let L be any ideal of R , and suppose that a, b generates L . Because R is a chain ring, either $\langle a \rangle \subset \langle b \rangle$ or $\langle b \rangle \subset \langle a \rangle$, and hence, L is principle ideal. We can easily use induction to show that any finitely generated ideal is principle. As R is a chain ring, it is also a local ring, and hence, $J(R)$ is the maximal ideal of R . Based on the above, $J(R)$ is generated by a single element, which we shall call θ . Thus, it is clear that

$$R \supset \langle \theta \rangle \supset \langle \theta^2 \rangle \supset \dots$$

To show that the above result is a series, suppose that there is an ideal K in R such that

$$J^i(R) = \langle \theta^i \rangle \supset K \supset \langle \theta^{i+1} \rangle = J^{i+1}(R),$$

for some i . Because R is a chain ring, K is principle. Let $K = \langle a \rangle$; then, $a \in J^i(R)$, $a = r\theta^i$, r is a unit; otherwise, if r in $J(R)$, then $a \in J(R)^{i+1}$, which is a contradiction. Thus, $r^{-1}a = \theta^i$ and $\theta \in K$, which implies that the above series is a unique series. \square

In the following proposition, the Jacobson radical of an R -module M denotes the intersection of all maximal R -submodules of M .

Proposition 2.2. Let R be a ring and M be a chain R -module. Then, the Jacobson radical $Rad(M)$ of M is a maximal R -submodule.

Proposition 2.3. Let R be a ring and M be a Noetherian chain R -module. Then, the Jacobson radical $Rad(M)$ of M is generated by a single element.

Proof. Suppose that $Rad(M)$ cannot be generated by one element and is thus generated by at least two elements, say, u, v . Therefore, Ru is not contained in Rv , and Rv is not contained in Ru , which produces a contradiction, as M is a chain R -module. Thus, $Rad(M)$ is generated by a single element. \square

Proposition 2.4. Let R be a commutative Noetherian chain ring, and let M be a Noetherian chain R -module. Then, the set of all R -submodules of M forms a unique series as follows:

$$M = \langle w \rangle \supset \langle \theta w \rangle \supset \langle \theta^2 w \rangle \supset \dots,$$

where $J(R)w = Rad(M)$, and $J(R) = \langle \theta \rangle$.

Proof. Let N be an R -submodule of M . It is clear that N is generated by a single element. Let $N = Ra$ for some a in M . As $N \subset RadM$, $a = r\theta w$, for some $r \in R$, $r = s\theta^i$, s is a unit; therefore, $N \subseteq J(R)^{i+1}w$. Moreover, $\theta^{i+1}w = s^{-1}r\theta w = s^{-1}a \in N$, and hence, $N = J(R)^{i+1}w$. Thus, all R -submodules of M form a chain as follows: $M = \langle w \rangle \supset \langle \theta w \rangle \supset \langle \theta^2 w \rangle \supset \dots$, where $J(R)w = Rad(M)$ and $J(R) = \langle \theta \rangle$. \square

3 Main Results

It is well known that for a ring R and a simple R -module M , $End_R(M)$ is a division ring. We shall replace the simplicity property with the chain property for an R -module M , assume that R itself is a chain ring, and study the ring $End_R(M)$. In this section, we would like to investigate certain properties of the R -module $Hom_R(M, N)$, where M, N are Noetherian chain R -modules. Then, we will discuss the endomorphism ring of M over a Noetherian chain ring R and show that the endomorphism ring $End_R(M)$ is a chain ring. Moreover, we shall study the special case in which $Hom_R(M, N)$ is the dual R -module $M^* = Hom_R(M, R)$.

In the following lemma, we will attempt to find a characterization for the left submodules of an R -module $Hom_R(M, N)$, where R is a commutative Noetherian chain ring and M, N are Noetherian chain R -modules.

Lemma 3.1. Let R be a commutative Noetherian chain ring, and let M, N be Noetherian chain R -modules. Then, each left submodule L of an R -module $K = Hom_R(M, N)$ is of the form $L = Rf$, for some $f \in K$.

Proof. Let L be a left submodule of an R -module $K = Hom_R(M, N)$; thus, any $g \in L$ takes any left submodule of M to a left submodule in N . The left submodules of M, N form a unique series as follows:

$$M = \langle w \rangle \supset \langle \theta w \rangle \supset \langle \theta^2 w \rangle \supset \dots,$$

where $J(R)w = Rad(M)$, and $J(R) = \langle \theta \rangle$,

$$N = \langle v \rangle \supset \langle \theta v \rangle \supset \langle \theta^2 v \rangle \supset \dots,$$

where $J(R)v = Rad(N)$, and $J(R) = \langle \theta \rangle$. As g is an R -module homomorphism, $g(w) = r\theta^i v$, r is a unit in R , $i \in 0, 1, 2, \dots$. Let f be the element in L for which $f(w) = r\theta^i v$, r is a unit in R and i is the smallest power for all elements in L . If h in L has the above property, such that $f(w) = r\theta^i v$, r is a unit in R and $h(w) = s\theta^i v$, s is a unit in R , then $rs^{-1}h = f$. Thus, we can take $r = 1$, $f(w) = \theta^i v$, where i is the smallest integer for all elements in L ; thus, $Rf \subseteq L$. Let g be any element in L . Then, $g(w) = t\theta^j v$, t is a unit in R , $j \geq i$, such that $t\theta^{j-i}f = g$, $g \in Rf$. Hence, $L = Rf$. \square

We will call the homomorphism f in the above proof the **minimal element** in L and i the **degree of f** . Now, we will study the ideals in the ring $End_R(M)$, where R is a Noetherian chain ring and M is a Noetherian chain R -module.

Lemma 3.2. Let R be a commutative Noetherian chain ring, and let M be a Noetherian chain R -module. Then, $End_R(M)$ is a principle ideal ring.

Proof. Let I be an ideal of the ring $End_R(M)$, so that any $f \in I$ takes any submodule of M to a submodule of M . The submodules of M form the following unique series:

$$M = \langle w \rangle \supset \langle \theta w \rangle \supset \langle \theta^2 w \rangle \supset \dots,$$

where $J(R)w = Rad(M)$ and $J(R) = \langle \theta \rangle$.

Therefore, because f is an R -module endomorphism, $f(w) = r\theta^i w$, r is a unit in R , $i \in 0, 1, 2, \dots$. Let g be the minimal element in I such that $g(w) = \theta^i w$. Then, for any $h \in End_R(M)$, we have $h(w) = t\theta^j w$, and thus,

$$hg(w) = h(\theta^i w) = \theta^i h(w) = \theta^i t\theta^j w = t'\theta^{i+j} w,$$

for some $t' \in R$ and hence $hg \in I$. Similarly, we can show that $gh \in I$ and $\langle g \rangle \subseteq I$. Now, let f be any element in I . Then, $f(w) = t\theta^j w$, t is a unit in R , $j \geq i$, which implies that $t^{-1}\theta^{j-i}g = f$, and thus $f \in \langle g \rangle$. Hence, $I = \langle g \rangle$, and $End_R(M)$ is a principle ideal ring. \square

Theorem 3.3. If R is a commutative Noetherian chain ring, and M, N are Noetherian chain R -modules, then the module $Hom_R(M, N)$ is a Noetherian chain R -module. Moreover, if N is of finite length, then the length of the composition series of $Hom_R(M, N)$ is equal to the length of the composition series of N .

Proof. Suppose that the composition series of M and N are as follows:

$$M = \langle w \rangle \supset \langle \theta w \rangle \supset \langle \theta^2 w \rangle \supset \dots,$$

$$N = \langle v \rangle \supset \langle \theta v \rangle \supset \langle \theta^2 v \rangle \supset \dots,$$

where $J(R) = \langle \theta \rangle$. Let I be a submodule of $Hom_R(M, N)$, and let $f \in I$. Then, the image of any submodule of M under f is a submodule of N . Assume that f is the minimal element in I , so that $f(w) = \theta^i v$, $i \in 0, 1, 2, \dots$ is the smallest power for all elements in I . Let K be any submodule in $Hom_R(M, N)$. Then, by lemma 2.1, $K = Rg$, g is the minimal element in K . Now, $g(w) = \theta^j v$, $j \in 0, 1, 2, \dots$. If $i \leq j$, $g(w) = \theta^j v = \theta^{j-i}\theta^i v = \theta^{j-i}f(w)$, and hence $Rg \subseteq Rf$. Conversely, if $j \leq i$, then $Rf \subseteq Rg$. Thus, $Hom_R(M, N)$ is a Noetherian chain R -module. Now, we will show that the unique series of $Hom_R(M, N)$ is as follows:

$$Hom_R(M, N) = Rh \supset R\theta h \supset R\theta^2 h \supset R\theta^3 h \supset \dots,$$

where $h(w) = v$. Let K be any submodule of $Hom_R(M, N)$. Then, $K = Rg$, $g(w) = \theta^j v$, $j \in 0, 1, 2, \dots$, is the minimal element in K , so $g = \theta^j h$, and hence, the above series is a unique series.

It is clear that if N is of finite length, then the length of the composition series of $Hom_R(M, N)$ is equal to the length of the composition series of N . \square

Here, we present a development of Schur's lemma, asserting that for a ring R and a simple R -module M , $End_R(M)$ is a division ring.

Corollary 3.4. Let R be a commutative Noetherian chain ring, and let M be a Noetherian chain R -module. Then, the endomorphism ring $End_R(M)$ is a Noetherian chain ring.

Proof. Let I, L be two ideals of $End_R(M)$. Then $I = \langle g \rangle, L = \langle h \rangle$, where g, h are the minimal elements in I and L and i, j are the degrees of g, h respectively. Assume that $i \leq j$. Then, $g(w) = \theta^i w, h(w) = \theta^j w = \theta^{i+(j-i)} w = \theta^{j-i}g(w)$, and hence, $h = \theta^{j-i}g \in \langle g \rangle, L \subseteq I$. Because I, L are arbitrary, we conclude that $End_R(M)$ is a chain ring. \square

Corollary 3.5. Let R be a commutative Noetherian chain ring, and let M be a Noetherian chain R -module. Then, $Hom_R(M, N)$ is an Artinian R -module if R is an Artinian ring.

Proof. From the proof for Theorem 3.3, we can find that the length of the composition series of R as a ring is greater than or equal to the length of composition series of the R -module $Hom_R(M, N)$. Hence, if R is an Artinian ring, then $Hom_R(M, N)$ is an Artinian R -module. \square

In general, the converse of the above corollary is not true. For example, if we assume that N has a finite length as an R -module, say n , and R is a chain ring that is not Artinian, then the length of the composition series of $Hom_R(M, N)$ is equal to n , and hence $Hom_R(M, N)$ is an Artinian R -module.

Corollary 3.6. Let R be a commutative Noetherian chain ring and M a Noetherian chain R -module. Then, $End_R(M)$ is an Artinian ring if R is an Artinian ring.

To obtain the following corollary, we need to assert that if R is a Noetherian chain ring, then R is a Noetherian chain R -module. Assuming that R is a chain ring, we aim to show that R is itself a chain R -module. Suppose that $a, b \in R$. As $J(R) = \langle \theta \rangle$ is the maximal left ideal of R , $Ra, Rb \subset J(R)$, $a = r\theta^i$ for some $i \in 0, 1, 2, \dots$, r is a unit with the smallest power of θ in Ra . Similarly, for Rb , we have $b = t\theta^j$, $j \in 0, 1, 2, \dots$, t is a unit with the smallest power of θ in Rb . Now (without loss of generality), assume that $i \leq j$. Then, $b = t\theta^j = t\theta^{j-i}r^{-1}r\theta^i = t\theta^{j-i}r^{-1}a \in Ra$, which implies that $Rb \subseteq Ra$. Hence, R is a Noetherian chain R -module.

Now, for the dual R -module M^* of M , we will set $N = R$ and obtain the following result.

Corollary 3.7. Let R be a commutative Noetherian chain ring, and let M be a Noetherian chain R -module. Then, $M^* = Hom_R(M, R)$ is a Noetherian chain module. If R is of finite length as an R -module, then the length of the composition series of M^* is equal to the length of the composition series of the left ideals in R .

4 Conclusions

We summarize our work in this paper as follows:

- a In Section (2), we recast some well known definitions of basic facts in this study. Namely, we define chain ring and chain module and we apply the Hom functor of these objects.
- b In Section (3), we have obtained results by fixing a commutative Noetherian chain ring and apply the Hom functor of a chain module with coefficients from this ring. In particular, we present the main results for the endomorphism ring of a Noetherian chain module and the dual of a chain module. We believe that our construction as well as our results will open new gates in the area of research in this direction.

Acknowledgement

The authors would like to thank Professor A. Laradji for correspondence regarding this work and for providing us with papers [8] and [10].

Competing Interests

The authors declare that no competing interests exist.

References

- [1] Bessenrodt C, Brungs HH. Right chain rings, part 1. SM-DU-181; 1990.
- [2] Al-olaiyan HA. The structure of chain rings. Department of Mathematics, College of Science, King Saud University, Riyadh, Saudi Arabia, 2001, Unpublished Ph. D thesis.
- [3] Jacobson N. Basic algebra 2, 2nd ed. Dover Publications; 2009.
- [4] Lam TY. Lectures on modules and rings. Graduate texts in Mathematics, 189. Springer-Vellag, New York; 1999.

- [5] Wang F, Kim H. Modules satisfying certain chain conditions and their endomorphisms. Bulletin of the Korean Mathematical Society. 2015;52(2):549-556.
- [6] Lee G, Roman CS, Zhang X. Modules whose endomorphism rings are division rings. Communications in Algebra. 2014;42(12):5205-5223.
- [7] Lee G, Rizvi ST, Roman CS. Transfer of certain properties from Modules to their endomorphism rings. Contemporary Mathematics. 2014;609:197-222.
- [8] Smith PF. Mapping between module lattices. International Electronic Journal of Algebra. 2014;15:173-195.
- [9] Smith PF. Complete homomorphisms between module lattices. International Electronic Journal of Algebra. 2014;16:16-31.
- [10] Tunganbaev AA. Flat modules and rings finitely generated as modules over their center. Mathematical Notes. 1996;60(2):254-277.

©2016 Alolaiyan and Alghamdi; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<http://sciencedomain.org/review-history/13052>