



Management of Sunflower Necrosis and Sunflower Leaf Curl Diseases by Using Insecticides

Poornima ^{a*}, Vijaykumar N. Ghante ^a, Vikas V. Kulakarni ^a
and Umesh M.R. ^a

^a AICRP (Sunflower), MARS, UAS, Raichur-584 104, Karnataka, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i72552>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117893>

Original Research Article

Received: 20/03/2024

Accepted: 29/05/2024

Published: 02/06/2024

ABSTRACT

A field experiment on management of Sunflower Necrosis and Sunflower Leaf Curl disease was conducted for three years during rabi season at Main Agricultural Research Station, Raichur. The experiment was planned with seven treatments (six insecticides and one untreated) in three replications for three successive years i.e., rabi 2016-17, 2017-18 and 2018-19. The insecticides used were imidacloprid 17.8 SL, fipronil 5 SC, triazophos 40 EC, diafenthiuron 50 WP, flonicamid 50 WG and spiromesifen 24 SC. For six treatments (T1 to T6) seed treatment was done with imidacloprid 600 FS and for T7 was maintained without seed treatment. Pooled analysis revealed that the treatment T4 :(Seed treatment with imidacloprid 600 FS@ 5ml/kg seed+ foliar spray with Flonicamide 50WG @ 0.25g/l at 30, 45 and 60 DAS recorded the lesser Sunflower necrosis and Sunflower leaf curl incidence of 4.21% and 4.54% respectively with higher yield of 1630 kg/ha, which is followed by the treatment T5 : (Seed treatment with imidacloprid 600FS @ 5ml/kg seed +

*Corresponding author: E-mail: poornimaagri@gmail.com;

foliar spray with Diafenthiuron 50 wp @ 1gm/lit at 30, 45 and 60 DAS) recorded Sunflower necrosis and Sunflower leaf curl incidence of 6.34% and 7.06% respectively with higher yield of 1560 kg/ha and both were at par. Both the treatments recorded B:C ratio of 1.76 and 1.59 respectively. Higher incidence of Sunflower necrosis and Sunflower leaf curl 10.70% and 18.46% respectively and lower yield of 968.4 kg/ha was recorded in untreated control.

Keywords: Sunflower necrosis; sunflower leaf curl; sunflower; management; insecticide.

1. INTRODUCTION

“Sunflower (*Helianthus annuus* L.), is one of the most important oilseed crops grown throughout the year across India. Sunflower is a newly introduced oilseed crop in India, but it was grown as ornamental plant since ancient times. Andhra Pradesh, Karnataka, Maharashtra and Tamil Nadu are the main sunflower growing states of India. In recent years, its cultivation was taken up in non-traditional states like Punjab, Haryana, Uttar Pradesh, Gujarat, Orissa, Madhya Pradesh and Rajasthan. In recent years sunflower area is rapidly declining and among many reasons diseases are one of the major contributors for cropped area reduction. Among viral diseases Sunflower necrosis and Sunflower leaf curl diseases are the important diseases. The Sunflower Necrosis disease was first observed during 1997 at Bagepally (Kolar) near Bangalore” [1], which occurred in an epidemic form consecutively for four years (1997-2001) with the incidence ranging from 10-80 per cent in both open pollinated and hybrid varieties and Sunflower leaf curl is a serious disease reported in India apart from necrosis [2]. “This disease was reported first time on Sunbred-275 during rabi, 2009 at Main Agricultural Research Station, Raichur. It is caused by sunflower leaf curl virus which belongs to Begomo virus group and is infecting many crops. Symptoms of sunflower necrosis disease (SND) comprises of chlorotic and necrotic ringspots, leaf distortion, leaf and stem necrosis and eventual death” [3]. “The causal virus of the disease has been identified as Tobacco Streak Virus (TSV) and shown to be transmitted by thrips in the presence of infected pollen grains” [3]. “The prominent symptoms of the Sunflower leaf curl disease observed were: small leaf size, malformed leaves, leaf and veinal thickening, enations and upward leaf curling. Emerging leaves exhibited yellow discoloration and severe reduction in leaf size. The infected plants were very much stunted when infected at early stages of crop and no ear head was erected. The disease significantly affects the plant height, head diameter, seed weight and oil percentage, etc” [4]. It is a whitefly (*Bemisia*

tabaci) transmitted virus and the vector attacks numerous fiber (particularly cotton), food, vegetable and ornamental plants apart from sunflower because of its polyphagous nature. A vector plays an important role in transmitting the virus. Keeping the above points in view, the present investigation was carried out to manage the vectors responsible for the transmission of virus. As the chemical management is highly effective and an important way to manage the vectors, this experiment was planned with six insecticides for management of Sunflower Necrosis and Sunflower Leaf Curl disease.

2. MATERIALS AND METHODS

A field experiment was conducted at Main Agriculture Research Station, Raichur, University of Agricultural Sciences, Raichur (16°12' N 77° 19'3 E, 407 m elevation) of Karnataka for a period of three seasons from rabi, 2017-18 to 2019-20 under All India Coordinated Research Project on Sunflower scheme. The climate of the region is semi-arid and subtropical. The soil of the area was medium to deep black, low in organic carbon (0.36 %), high in available P₂O₅ (45 kg /ha) and available K₂O (536 kg /ha).

Randomized Block Design with seven different treatments in three replications. Plot area was maintained as 4.2 x 3 m² (60 plants per plot), spacing of 60x30 cm and fertilizer schedule of 75- 90-30 NPK kg/ha was followed. Seeds were dressed with liquid formulation of insecticide. Three irrigations were given at different stages i.e., star bud stage, flowering and seed setting in the crop period. All agronomic practices were followed as per package of practices. The soil of the experimental field was black, with pH 8.3 and EC 0.26 dS/m. Seeds of susceptible check (KBSH-44) were obtained from Indian Institute of Oilseed Research, Hyderabad. Three irrigations were given at different stages i.e., star bud stage, flowering and seed setting in the crop period. All agronomic practices were followed as per package of practices. The foliar application of insecticides was done as per schedule at 30, 45 and 60 days after sowing.

Data on Sunflower necrosis and leaf curl incidence was recorded monthly interval starting from sowing date and final data was taken into consideration. Percent of Disease incidence was calculated by using the following formula:

$$\text{PDI (\%)} = \frac{\text{Number of plants infected}}{\text{Total number of plants}} \times 100$$

3. RESULTS AND DISCUSSION

The present investigation was carried out using insecticides to manage viral diseases of sunflower. The results indicated that during 2016-17, germination percentage recorded was more than 90% in all the treatments and it was found that there was no significant difference among the treatments. The results are elucidated in Table 1. Among the imposed treatments, less Sunflower necrosis and Sunflower leaf curl incidence of 4.21% and 4.54% respectively was observed in the treatment T5 (Seed treatment with Imidacloprid 600FS @ 5ml/kg seed + foliar spray with Flonicamide 50WG @ 0.25g/l at 30, 45 and 60 Date after sowing) followed by the treatment T4 (Seed treatment with Imidacloprid 600FS @ 5ml/kg seed + foliar spray with Diafenthiuron 50 WP @ 1gm/lit at 30, 45 and 60 DAS) recorded disease incidence of Sunflower necrosis 5.17% and Sunflower leaf curl disease incidence of 3.85% respectively. Higher Sunflower necrosis and Sunflower leaf curl incidence of 28.75% and 16.26% respectively was recorded in untreated plot. Higher yield of 1630 kg/ha was recorded in the treatment T5, followed by T4 which recorded seed yield of 1560. kg/ha and the treatment T2 recorded 510 kg/ha; however, all these treatments were at par with each other and significantly differed with untreated plot. Less yield of 863.3 kg/ha was observed in untreated plot among all the treatments and it was non-significant with treatments T5 and T4.

During 2018-19 also, similar trend was observed i.e., the treatment T5 recorded the lowest Sunflower necrosis and Sunflower leaf curl incidence of 3.89% and 2.78% respectively followed by the treatment T4 recorded Sunflower necrosis (4.82%) and Sunflower leaf curl (4.10%) both the treatments were at par each other (Table 1) More Sunflower necrosis and Sunflower leaf curl disease incidence of 13.63% and 18.25% respectively was observed in T7 (untreated plot). Regarding yield, all the treatments showed higher seed yield over that of untreated plot. However, significant differences

were found between untreated plot and T5 treatment.

During the year 2018-19, data (Table 1) indicated that the treatment T5 recorded lesser disease incidence of Sunflower necrosis 3.58% and Sunflower leaf curl 6.99%. Like previous two years results, untreated plot recorded more Sunflower necrosis and Sunflower leaf curl disease incidence of 14.76% and 20.87% respectively.

Pooled analysis results indicated that, Among the imposed seven treatments, the treatment T5 recorded less Sunflower necrosis and Sunflower leaf curl incidence of 4.21% and 4.54% and the same treatment has recorded higher yield of 1630 kg/ha, followed by the treatment T4 which recorded Sunflower necrosis (6.34%) and Sunflower leaf curl (7.06%) with seed yield of 1485.6 kg/ha. Higher Sunflower necrosis and Sunflower leaf curl incidence of 10.70% and 18.46% respectively with lower yield of 968 kg/ha was observed in untreated plot. These results indicated that insecticides in most of the treatments reduced Sunflower necrosis and Sunflower leaf curl incidence, by suppressing the thrips and whitefly (vector) population and among them Flonicamide was found to be most effective. The results are in accordance with Venkataramanamma et. al. 2022 where treatment T4 (Seed treatment with Imidacloprid 600FS @ 5ml/kg seed + foliar spray with Diafenthiuron 50 wp @ 1gm/lit) was observed with least leaf curl incidence of 23.71% with higher seed yield of 1591 kg/ha followed by T4 Seed treatment with Imidacloprid 600FS @ 5ml/kg seed + foliar spray with Flonicamide 50WG @0.25g/l recorded leaf curl incidence of 24.29% with yield 1563kg/ha.

Deepa et al. [5] conducted an experiment on management of leaf curl incidence of sunflower with insecticides, leaf extracts and bioagents and the results concluded that seed treatment with imidacloprid @ 5 g/kg along with two sprays of imidacloprid @ 0.5 ml/l or triazophos @ 1.5 ml/l were effective in controlling the leaf curl incidence and vector population with increased yield. In the present experiment also, disease reduction was found with insecticides but results of foliar spray with imidacloprid and fipronil were at par with untreated plot. Rajasekhar et al. [6], who tested 8 insecticides against whiteflies of cotton under normal and high density planting system, found that diafenthiuron 50% WP @ 1.25 g/litre was effective in reducing whitefly

Table 1. Management of viral diseases of sunflower using insecticides

Treatments	Sunflower Necrosis disease (PDI)				Sunflower leaf curl (PDI)			
	2016-17	2017-18	2018-19	Pooled	2016-17	2017-18	2018-19	Pooled
Seed treatment with Imidacloprid 600FS @ 5ml/kg seed and foliar spray with Imidacloprid 17.8SL @ 0.5ml/l at 30, 45 and 60 DAS	14.35 (22.04)	8.21 (16.65)	9.78 (18.17)	9.17 (17.60)	9.89 (18.33)	6.53 (14.80)	16.22 (23.75)	10.88 (18.96)
Seed treatment with Imidacloprid 600FS @ 5ml/kg seed + foliar spray with Fipronil 5SC @ 1ml/lit at 30, 45 and 60 DAS	8.79 (16.95)	6.53 (14.81)	7.09 (15.39)	7.47 (15.80)	6.63 (14.92)	5.58 (13.66)	16.81 (24.20)	9.67 (17.60)
Seed treatment with Imidacloprid 600FS @ 5ml/kg seed + foliar spray with Trizophos 40EC @ 1ml/lit at 30, 45 and 60 DAS	10.49 (18.73)	5.88 (14.02)	8.53 (16.82)	8.30 (16.68)	8.28 (16.73)	5.24 (13.33)	15.99 (23.57)	9.84 (17.84)
Seed treatment with Imidacloprid 600FS @ 5ml/kg seed + foliar spray with Diafenthiuron 50 wp @ 1gm/lit at 30, 45 and 60 DAS	7.94 (15.99)	4.82 (12.68)	6.25 (14.44)	6.34 (14.48)	5.98 (14.15)	4.10 (11.69)	11.09 (19.46)	7.06 (15.10)
Seed treatment with Imidacloprid 600FS @ 5ml/kg seed + foliar spray with Flonicamide 50WG @ 0.25g/l at 30, 45 and 60 DAS	5.17 (12.88)	3.89 (11.36)	3.58 (10.61)	4.21 (11.73)	3.85 (11.31)	2.78 (9.59)	6.99 (15.33)	4.54 (12.08)
Seed treatment with Imidacloprid 600FS @ 5ml/kg seed + foliar spray with Spiromesifen 24SC @ 1ml/l at 30, 45 and 60 DAS	12.99 (20.83)	4.59 (12.36)	4.39 (12.07)	7.32 (15.65)	7.85 (16.27)	3.90 (11.39)	13.95 (21.93)	8.57 (16.53)
Control	28.75 (32.37)	13.63 (21.66)	14.76 (22.48)	10.70 (19.06)	16.26 (23.78)	18.25 (25.29)	20.87 (27.18)	18.46 (25.42)
S. Em.±1	2.39	0.39	1.44	1.04	1.30	0.88	1.55	1.01
C. D. at 5%	7.36	1.21	4.43	3.21	4.00	2.72	4.78	3.11

Table 2. Seed yield (Kg/ha) and B: C Ratio under various treatments

Treatments	Yield (Kg/ha)				B:C Ratio
	2016-17	2017-18	2018-19	Pooled	
Seed treatment with Imidacloprid 600FS @ 5ml/kg seed and foliar spray with Imidacloprid 17.8SL @ 0.5ml/l at 30, 45 and 60 DAS	1248.33	1363	1167	1259.4	1.22
Seed treatment with Imidacloprid 600FS @ 5ml/kg seed + foliar spray with Fipronil 5SC @ 1ml/lit at 30, 45 and 60 DAS	1510.83	1455	1311	1425.6	1.62
Seed treatment with Imidacloprid 600FS @ 5ml/kg seed + foliar spray with Trizophos 40EC @ 1ml/lit at 30, 45 and 60 DAS	1364.67	1542	1268	1391.5	1.62
Seed treatment with Imidacloprid 600FS @ 5ml/kg seed + foliar spray with Diafenthiuron 50 wp @ 1gm/lit at 30, 45 and 60 DAS	1560.00	1648	1249	1485.6	1.59
Seed treatment with Imidacloprid 600FS @ 5ml/kg seed + foliar spray with Flonicamide 50WG @ 0.25g/l at 30, 45 and 60 DAS	1630.00	1750	1668	1682.6	1.76
Seed treatment with Imidacloprid 600FS @ 5ml/kg seed + foliar spray with Spiromesifen 24SC @ 1ml/l at 30, 45 and 60 DAS	1455.00	1685	1454	1531.3	1.58
Control	863.33	1025	1017	968.4	0.84
S. Em.±1	102.8	100.04	96.13	81.49	
C. D. at 5%	316.75	297.22	296.20	251.1	

population followed by flonicamid 50%WG@ 0.3 g/litre and acephate 75 SP@ 1.5 g/ litre, but promising results were not obtained with imidacloprid 17.8% SL and fipronil 5% SC. Shirshikar, [7] recorded the use of imidacloprid for seed treatment along with imidacloprid foliar spray and this treatment were found to be useful in the management of sunflower necrosis disease. In the present study seed treatment with imidacloprid was common for all the treatments and compared with different insecticides. Pandey et al. [8] reported that lowest thrips population and highest bulb yield by applying fipronil. Similarly fipronil and imidacloprid reduced the thrips damage severity and increased the onion bulb yield [9,10]. Seed treatment was done with imidacloprid 600FS @5g/kg of seed in all the treatments to suppress the sucking pest population in initial days of sowing. The effect of seed treatment was in conformity with Singh et al. [11], who suggested that seed treatment with imidacloprid (Gaucho 70WS) @5g/kg of seed reduced whitefly (vector population) incidence compared to untreated plot and reduced leaf curl incidence in cotton. Similar results were reported by Chandel et al. [12].

Similarly, the highest B:C ratio 1.76 was recorded in T5 followed by T2 and T3 recorded 1.62 whereas untreated control recorded lowest B:C ratio of 0.84. Evaluation of various molecules of insecticides against the thrips and whitefly (vector) transmitted Sunflower necrosis and Sunflower leaf curl disease respectively revealed that even though the seed treatment was common in all the treatments, the spraying of insecticide diversified the results. Foliar spray with Flonicamide 50WG @ 0.25g/l at 30, 45 and 60 DAS exhibited less Sunflower necrosis and Sunflower leaf curl disease incidence (4.21% and 4.54% respectively) whereas, foliar spray with Diafenthiuron 50 wp @ 1gm/lit at 30, 45 and 60 DAS resulted in good seed yield. Rational use of these insecticides plays a significant role in management of *Thrips palmi* and *Bemisia tabaci* and consequently Sunflower necrosis and Sunflower leaf curl incidence. Based on the above data, it was concluded that the two chemicals (flonicamid and diafenthiuron) will be effective at recommended concentration for management of vector and hence Sunflower necrosis and Sunflower leaf curl incidence [13-16].

4. CONCLUSION

Among different treatments, the treatment T4: (Seed treatment with imidacloprid 600 FS@

5ml/kg seed+ foliar spray with Flonicamide 50WG @ 0.25g/l at 30, 45 and 60 DAS was found effective for the management of Sunflower necrosis and Sunflower Leaf curl disease.

ACKNOWLEDGEMENT

The authors are thankful to the AICRP Sunflower, MARS,UAS, Raichur, for constant encouragement and providing necessary facilities to carry out the research work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Singh SJ, Nagaraju, Krishna Reddy M, Muniyappa V, Virupakshappa K. Sunflower necrosis a new virus disease from India. In: Nat. Symp. Eco. Imp. Diseases of Crop Plant, Dec 18-20th, IPS (S-Zone), Univ. Agric. Sci., Bangalore, India; 1997.
2. Govindappa MR, Shankar gouda I, Shankarappa KS, Wickramaarachchi WART, Anjaneya Reddy B. Molecular detection and partial characterization Begomovirus associated with leaf curl disease of sunflower (*Helianthus annuus*) in Southern India. Plant Pathology Journal. 2011;10(1):29-35.
3. Ravi KS, Buttgeritt A, Kitkaru AS, Deshmukh S, Lesemann DE, Winter S. Sunflower necrosis disease from India is caused by an ilarvirus related to Tobacco streak virus. New Disease Report. 2001; 50(6):800.
4. Deepa, Sunkad G. Survey and characterizations of symptoms of the sunflower leaf curl virus disease transmitted by whitefly. Ecology Environment Conservation. 2015;22(4): 1999-2006.
5. Deepa, Gururaj S, Govindappa MR, Naik MK, Somasekhar. Management of new sunflower leaf curl virus disease caused by Begomovirus. Trends in Bioscience. 2014; 7(16):2127-2131.
6. Rajasekhar N, Durgaprasad NVVS, Sairamkumar DV, Adinarayana M. Incidence and management of cotton whitefly *Bemisia tabaci* under High density planting system (HDPS). International Journal Current Microbiology Applied Science. 2018;7(8):2074-79.

7. Shirshikar SP. Integrated management of sunflower necrosis disease. *Helia*. 2008; 31(49):27-34.
8. Pandey S, Singh BK, Gupta RP. Effect of neembased botanicals, chemicals and bio-insecticides for the management of thrips in onion. *Indian Journal Agriculture Research*. 2013;47(6):545 - 548.
9. Ullah F, Mulk M, Farid A, Saeed MQ, Sattar S. Population dynamics and chemical control of onion thrips (*Thrips tabaci* Lindemann). *Pakistan Journal Zoology*. 2010;42(4):401 - 406.
10. Gachu SM, Muthomi JW, Narla RD, Nderitu JH, Olubayo FM, Waga JM. Management of thrips (*Thrips tabaci*) in bulb onion by using of vegetable inter crops. *International Journal of Agriculture*. 2012;2(5):393 - 402.
11. Singh D, Singh K, Garg HR. Management of cotton leaf curl virus vector (CLCuV) disease managing its vector, (*Bemisia tabaci* Genn.). *World Cotton Research*; 2003
12. Chandel RS, Banyal DK, Singh BP, Kamlesh M, Lakra BS. Integrated management of whitefly, *Bemisia tabaci* (Gennadius) and potato apical leaf curl virus in India. *Potato Research*. 2010; 53(2):129-39.
13. Venkataramanamma K, Neelima S, Prabhakar K, Lakshmi Kalyani D. Management of Leaf Curl Disease of Sunflower under field conditions. *Agriculture Research Journal*. 2022;59(3): 447-452.
14. Dole PD, Gaikwad SD, Digvijay C. Optimizing tomato health: Holistic approach to early blight disease management. *Journal of Advances in Biology and Biotechnology*. 2024;27(6): 492–497. Available:<https://doi.org/10.9734/jabb/2024/v27i6909>
15. Jayashree A, Suresh KP, Raaga R. Advancing Coffee Leaf Rust Disease Management: A Deep Learning Approach for Accurate Detection and Classification Using Convolutional Neural Networks. *Journal of Experimental Agriculture International*. 2024;46(2):108–118. Available:<https://doi.org/10.9734/jeai/2024/v46i22313>
16. Salari K, Heydarnejad J, Massumi H, Hasanvand V, Varsani A. Incidence of cotton leaf curl Gezira virus and the associated alphasatellites and betasatellites in crops and ornamental plants in southern Iran. *Tropical Plant Pathology*. 2023;48(1):62-72.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/117893>