



# Remediation of Saline Soils Using Halo-Tolerant Plant Growth Promoting Rhizobacteria

Charan Singh <sup>a++</sup>, Gayatri Kumari <sup>b\*</sup>, Lalita <sup>a++</sup>,  
Vishal Gandhi <sup>a++</sup>, Ashish Jain <sup>a++</sup>, Sukham Madaan <sup>a++</sup>,  
Sumit Saini <sup>a++</sup>, Amit Kumar <sup>a++</sup>, Mahaveer <sup>a++</sup>  
and Rakesh Kumar <sup>a++</sup>

<sup>a</sup> Rice Research Station, CCS HAU, Kaul, Kaithal, Haryana, India.

<sup>b</sup> Department of Botany and Plant Physiology, CCS HAU, Hisar, Haryana, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijecc/2024/v14i64208>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118111>

Review Article

Received: 28/03/2024

Accepted: 03/06/2024

Published: 04/06/2024

## ABSTRACT

Soil salinization poses a significant threat to global agriculture, affecting approximately 6.73m Ha land area in India. Salinity stress impacts plant growth and soil health negatively, leading to reduced crop yields and soil degradation. This review examines the sources and effects of soil salinity, highlighting the intricate interplay between salinity and soil nutrients and its remediation. Traditional methods for soil remediation often have detrimental long-term effects, prompting the exploration of alternative strategies such as the use of halo-tolerant plant growth-promoting

<sup>++</sup> Assistant Scientist;

\*Corresponding author: E-mail: ydvgayatri714@gmail.com;

**Cite as:** Singh, Charan, Gayatri Kumari, Lalita, Vishal Gandhi, Ashish Jain, Sukham Madaan, Sumit Saini, Amit Kumar, Mahaveer, and Rakesh Kumar. 2024. "Remediation of Saline Soils Using Halo-Tolerant Plant Growth Promoting Rhizobacteria". *International Journal of Environment and Climate Change* 14 (6):24-35. <https://doi.org/10.9734/ijecc/2024/v14i64208>.

rhizobacteria. HT-PGPR offer a promising solution for sustainable agriculture by enhancing soil fertility and plant resilience to salinity stress through various mechanisms. Furthermore, this review identifies research gaps in understanding the metabolic pathways and strain selection of HT-PGPR, as well as their interactions with soil microbiota. Future research directions include field-scale experiments to validate the effectiveness and economic viability of HT-PGPR inoculation for large-scale application in saline soils. Overall, leveraging the potential of HT-PGPR represents a critical step towards mitigating the global challenge of soil salinity and ensuring food security in the face of climate change.

**Keywords:** Soil salinity; plant growth; HT-PGPR; field application; soil nutrients.

## 1. INTRODUCTION

Globally, 424 million hectare (Mha) surface (0-30 cm) and 833 Mha sub soil (30-100 cm) which is about 73% of the land area is salinity affected [1]. Agricultural practices rely significantly on the state of land and water use, both of which are progressively influenced by climate change [2]. Saline soil is identified by having an  $EC_e$  of about 4 dS/m or more in the root zone with 15% ESP and a pH below 8.5 [3]. Although multiple salt ions contribute to salinity, sodium ( $Na^+$ ), calcium ( $Ca^{2+}$ ), magnesium ( $Mg^{2+}$ ), chloride ( $Cl^-$ ), sulphate ( $SO_4^{2-}$ ) and carbonate ( $CO_3^{2-}$ ) are the most prevalent and display the greatest levels of phytotoxicity [4]. Soil alkalinity, a variant of soil salinity, arises when the soil is primarily saturated with sodium carbonate, resulting in a higher soil pH. The effects of soil alkalinity can be more severe than those of normal salinized soils due to its impact on nutrient availability and soil microbial activity [5]. Salinization causes nutritional disorders in plants by modifying nutrient availability and disrupting processes like competitive absorption, transport and distribution within the plant. The interplay between salinity and primary mineral nutrition is intricate and can negatively impact essential soil nutrients such as NPK [6]. Cultivating crops under saline soils is a major problem that farmers face today as salinity in the soil results in yield losses of the crop. To counter the negative effect of salinity, farmers use different physical and chemical techniques including excess use of fertilizers and the use of chemicals. These methods can have negative effects on plant and soil health. An alternative to this is the use of halo tolerant plant growth

promoting rhizobacteria (HT-PGPR) which help the soil to regain its fertility and do not have any ill effects on soil and plants. Inoculation of HT-PGPR can alleviate plant salt stress by triggering various physiological and molecular mechanisms. This includes modifying root systems, activating antioxidant mechanisms, producing siderophores and exopolysaccharides (EPS), regulating phytohormones, synthesizing osmolytes, absorbing minerals and controlling phytopathogens [7]. It has been observed that several halotolerant soil bacterial species, including *Arthrobacter*, *Azospirillum*, *Alcaligenes Bacillus*, *Burkholderia*, *Enterobacter*, *Flavobacterium*, *Pseudomonas* and *Rhizobium*, can help crops under salinity stress [8]. Overall, the use of HT-PGPR bioinoculants represents a promising strategy for sustainable agriculture, offering multiple benefits including improved soil health, enhanced water retention and increased productivity in saline environments while minimizing environmental impact.

## 2. SOURCES OF SOIL SALINIZATION

According to international standards, saline and alkaline soils can be classified into three categories based on specific salinity and alkalinity parameters. These classifications utilize soil reaction pH, electrical conductivity (EC) and either the sodium adsorption ratio (SAR) or the exchangeable sodium percentage (ESP) (Table 1). Salinization is governed by various factors, including environmental conditions, water supply and management systems (irrigation and drainage), and agricultural practices (such as plant cover type, density and rooting

**Table 1. Classification of salt-affected soils**

Classification of Salt-Affected Soil	$EC_e$ (dS $m^{-1}$ )	pH	SAR	Soil Physical Condition
Saline	> 4.0	< 8.5	< 13	Normal
Saline-sodic	> 4.0	< 8.5	> 13	Normal
Sodic	< 4.0	> 8.5	> 13	Poor

characteristics) [9]. These factors influence the soil-water balance, thereby affecting the movement and accumulation of salts within the soil. Salinization typically occurs under conditions where soluble salts are present, the water table is high, evaporation rates are high and annual rainfall is low. Soil salinization can result from various natural and human-induced factors such as weathering of rocks and improper agricultural practices etc. Okur and Örcen, [10].

### 3. NATURAL PROCESSES

The weathering of the parent rocks and soil, atmospheric deposition and groundwater are the sources of ions, including carbonate and sulphate. Minerals undergo weathering as a result of water action and dissolved CO<sub>2</sub> [11]. Salinity has an impact on coastal ecosystems, including salt marshes. There are usually three types of marsh zones in tidal salt marshes: high, middle and low. Near the high marsh zone, the salinity of the soil reaches a maximum. The shortened tidal inundation period, which permits evapotranspiration to concentrate pore water salinity and salt to build, is one of the primary causes of this salinity increase [12]. The capillary rise in groundwater also results in soil salinization as in areas with high groundwater tables, capillary action can draw saline water up to the soil surface, where it evaporates and leaves salts behind [13].

### 4. HUMAN-INDUCED PROCESSES

First and foremost, salinization is a naturally occurring process brought about by the movement and dispersal of water-soluble salts from a salt source location to an area that was initially salt-free [14]. Salinization is also a human-induced process, known as secondary salinization, that either causes salt-free soils to become contaminated due to poor water and land management practices or increases the concentration of salt in soils that are already affected by salt.

Salt concentrations in the root zone can result from the use of saline groundwater for irrigation, particularly if internal drainage is impeded and leaching is inadequate [15]. Small areas are affected by this salinization process, which happens in arid conditions. A build-up of salt concentration in the soil can occur from the prolonged use of irrigation water, even in cases when the amount of dissolved salt is low, in regions with significant evapotranspiration [16].

The use of excess fertilizers also results in soil salinization as some fertilizers contain salt ions that can accumulate in the soil due to excess and prolonged use [17]. High crop demands and continuous cropping can deplete soil nutrients and contribute to salinity. Discharge of industrial wastewater can also introduce salts into the soil [18].

## 5. EFFECT OF SALINITY ON SOIL PROFILE

Crops exhibit various responses to salinity stress, which not only affects their productivity but also induces changes in soil physicochemical characteristics and ecological balance. Salinity exerts detrimental effects on agricultural yield by reducing plant water uptake capacity, resulting in soil erosion and diminished economic returns. It represents a substantial challenge in various regions worldwide [19].

### 5.1 Plant Available Water

Soil salinity presents a challenge when an accumulation of soluble salt ions in the root zone reaches levels that harm plant growth. Elevated levels of salt ions in the root zone hinder a plant's capacity to absorb water from the surrounding soil, reducing available water for plants [20]. In saline environments, water retention in the soil is enhanced, but its accessibility for plant uptake is reduced due to osmotic stress. Consequently, this is likely to result in reduced growth and manifestations similar to drought stress, such as wilting or leaf abscission [21]. Following irrigation, soil salinity is minimal and plant-available water reaches its peak. The water is transpired by plants or evaporates into the atmosphere, the remaining soil water becomes more saline. This is attributed to the retention of salts while water diminishes in volume due to transpiration and evaporation. This phenomenon is particularly significant during high temperatures and the peak of the growing season which increases the evapotranspiration rate [22].

### 5.2 Flocculation under Salinity

Increased salinity in soil can induce flocculation, a process where fine soil particles clump together to form aggregates. Higher salt levels in the soil solution encourage the aggregation of clay particles which increases the flocculation rate in the soil. On the other hand, the spaces between soil aggregates are comparatively larger

than in non-flocculated soil, enhancing soil permeability and reducing the likelihood of waterlogging upon wetting [23,24].

### 5.3 Infiltration Rates

The ratio of salinity to sodicity is indicated by the SAR value which is crucial in determining the impact of salts and sodium on the soil while infiltration rates determine the effect of saline and sodicity on the soil physical health [25]. The overall salinity level of the soil solution impacts certain physicochemical properties of the soil. Conversely, the presence of Na<sup>+</sup> in the soil solution and its absorption into the soil exchange complex can induce distinct effects on soil physicochemical traits [26]. Salt ions viz. Na<sup>+</sup> are dissolved in the soil water solution and are thus transported downward along with soil water, primarily through drainage within the soil profile. However, sodium being attached to soil particles, remains immobile and does not migrate downward with soil water [27]. Elevated sodium concentration induces soil dispersion, while higher salinity levels promote soil flocculation. Rainfall can enhance the possibility of sodium-induced dispersion and decrease the possibility of soil aggregations caused by salts.

The physicochemical attributes of salt-affected soils exhibit seasonal fluctuations. These variations manifest in the range of salinity from 0.76 to 1.24 dSm<sup>-1</sup>, with salinity levels typically escalating in the surface layers during summer due to elevated evaporation rates and decreasing during the monsoon period owing to leaching processes [28]. Salt-affected soils exhibit increased salinity and dominance of Na<sup>+</sup> and Cl<sup>-</sup> ions, leading to higher SAR values in the

summer compared to other seasons. Consequently, optimal conditions for seed sowing or cultivation in such soils are likely during periods of minimal salinity, notably in the monsoon season [29].

## 6. EFFECT OF SALINIZATION ON PLANTS

Salinity has a major impact on nearly every facet of agricultural development, including decreased yield potential and productivity and the onset of soil contamination and degradation. Around the world, salinized soils are known for causing salinity stress, which inhibits the growth of salt-intolerant plants [30]. The negative impacts of salinity typically occur in two stages. Initially, the altered osmotic pressure due to excess salt ions in the root zone disrupts regular water uptake, known as the physiological drought conditions. This disruption impairs cell division and differentiation, resulting in stunted growth and development. Furthermore, excessive salt ion accumulation in the cell walls offers osmotic stress and ionic toxicity to plants and sometimes can also result in cell death [31] (Fig. 1). Salinity results in membrane injury of the cell and lowers the total carotenoid and chlorophyll content thus affecting photosynthesis in the plants [32]. Certain plants modify their photosynthetic pathway in response to salinity stress. For instance, *Mesembryanthemum crystallinum* and *Atriplex lentiformis* respond to water scarcity in saline conditions by switching from the C3 to CAM and C3 to C4 pathways respectively [33]. Increased soil salinity offers ionic toxicity to plants which results in the formation of ROS in the plant cells [34]. These ROS alter membrane permeability and protein structures results




Fig. 1. Impact of salinity stress on plants

in resulting in disruption of plant growth and metabolism [35]. Salinity results in deteriorative plant growth as under such conditions, plant cells become more permeable to  $\text{Na}^+$  as compared to  $\text{K}^+$  which changes membrane potential and also causes  $\text{K}^+$  deficiency in the plants. The absorption of  $\text{K}^+$ ,  $\text{NO}_3^-$ ,  $\text{PO}_4^{2-}$ ,  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  reduce under salinity and as a result, the ratios of  $\text{K}^+/\text{Na}^+$ ,  $\text{Ca}^{2+}/\text{Na}^+$ ,  $\text{Mg}^{2+}/\text{Na}^+$  and  $\text{PO}_4^{2-}/\text{Cl}^-$  also decreases which causes various macro and micronutrient deficiencies in the plants. High sodium and chloride concentrations can also cause ion imbalance and toxicity to specific ions [4].

Excess salinity results in various deteriorative effects on plants which makes plants more prone to various diseases and biotic stresses [36]. However, some plants become tolerant to salinity by improving growth and survival through various metabolism and adaptations which include maintaining ionic balance, enhanced antioxidative enzymatic activity and production of osmolytes to maintain osmotic balance in the plants and sometimes plants try to avoid the adverse effect of salinity by shortening their life cycle.

## 7. ALTERNATIVE AND TRADITIONAL METHODS FOR REMEDIATION OF SALINE SOILS

Efforts have been made from time to time to address salinization issues and the urgent need for fertile agricultural lands, including soil reclamations and crop modifications. Standard methods for soil reclamation include physical techniques (salt scrapping, flushing and soil washing), chemical amendments (gypsum) and various agricultural management practices (irrigation management, crop rotation and intercropping). But in practice, budgetary constraints and the quality of the water supply constantly place restrictions on such projects [37]. Furthermore, overuse of pesticides is typically involved by farmers since plants under salinity stress are more susceptible to disease and pests [38]. These short-term fixes will ultimately cause even more soluble salt accumulations from the chemical additives, which will exacerbate the salinity issues [39]. Different molecular approaches have also been discussed for the reclamation of saline soil which include the use of modified crops using genetic engineering, induced mutant breeding, priming

agent treatments, conventional selective breeding and other methods to give crops new potential traits like increased salt tolerance. Saline soil reclamation require a sustainable and reliable approach with no long-term ill effect on soil health and improved plant growth under saline conditions [9]. The application of HT-PGPR inoculation is an effective alternate strategy for mitigating salinity hazards and enhancing crop performance. Naturally, the plant tolerance to salt and biotic stresses is increased by these advantageous microorganisms significant growth-promoting and salinity-defensive characteristics. HT-PGPR alleviate salinity hazards through various pathways that activate complex physiological, biochemical and molecular responses in plants to support their growth [40].

## 8. HALOTOLERANT BACTERIA AS SOIL SALINITY REMEDIATORS

Using halotolerant bacteria for soil remediation offers a sustainable approach to address salinity issues, especially in agricultural areas where saline soils can severely impact crop productivity (Fig. 2). Research findings have validated that inoculation with HT-PGPR positively influences nutrient mineral uptake, soil physicochemical properties, organic carbon (OC) content, pH, EC and the accumulation of ionic salts in the soil AbuQamar *et. al.*, [7], Arora *et. al.*, [41] Numerous studies have shown that applying HT-PGPR can improve the salt index of saline soil. Additionally, HT-PGPR addresses nutrient deficiencies by increasing levels of nitrogen (N), phosphorus (P), potassium (K), iron (Fe) and zinc (Zn) in saline soils, thereby restoring vegetation and promoting agricultural sustainability [42].

Halotolerant bacteria can actively participate in the remediation of salt-affected soil involving enzymatic pathways which are capable of degrading specific salt compounds, converting them into less harmful substances [43]. Some bacteria can also bind or sequester salt ions, reducing their availability in the soil and mitigating their negative effects on plant growth known as salt immobilization [44]. Halotolerant bacteria may contribute to the improvement of soil structure and fertility in saline soils by producing EPS which act as soil binders, improving soil aggregation and water retention [45]. HT-PGPR in the soil helps in enhancing

organic matter decomposition which contributes to soil nutrient cycling [40]. Beyond nutritional aspects, soil quality is greatly influenced by aggregation, which enhances the infiltration rate, root aeration and penetration and the development of micropores [46]. HT-PGPR enhance soil physical quality through the synthesis of EPS, particularly under stress conditions [47]. This process promotes the formation of microaggregates of size <250 µm and macroaggregates of size >250 µm, which effectively sequester nutrients and water molecules [48]. The establishment of a cohesive EPS-soil matrix offers a protective microenvironment, particularly against salinity stress, thereby safeguarding both plant health and the HT-PGPR population within the rhizosphere. Functioning as a defensive barrier, the EPS within the biofilm complex serves to enhance soil water retention capacity and ameliorate water activity in plants during salinity and drought stress conditions [49]. Research has shown that *Pseudomonas putida* strain GAP-P45 EPS improved soil water retention capacity and aggregation under saline conditions [50]. Biofilm formation within soil particles and on root surfaces is marked by high concentrations of root exudates, signalling

molecules, water content and OC content [51]. This complex matrix plays a crucial role in the establishment and selection of different microbial populations. The predominant constituent of EPS regulates organic matter dynamics by serving as a carbon source and facilitating soil particle coagulation, thus promoting the formation of stable organic carbon compounds known as humic substances [52]. An increase in salinity diminishes the bacterial populations. The exogenous application of HT-PGPR enhances soil structure and promotes the assimilation of organic matter, thereby fostering microbial interactions. An increase in microbial population and dehydrogenase activity was observed in saline soil after introducing *Bacillus cereus* strain Pb25, which belongs to the category of HT-PGPR [53]. This research provides compelling evidence of the role and potential application of HT-PGPR in enhancing soil quality under abiotic stress conditions such as salinity. Table 2 provides an insight into previous research revolving around how inoculation of HT-PGPR in different crops helps mitigate salinity stress through different mechanisms resulting in improved growth and metabolism of the crop plants.




Fig. 2. Remediation of salt-affected soil with inoculation of HT-PGPR

**Table 2. HT-PGPR mediated tolerance in different crops under salinity stress**

HT-PGPR	Crop Evaluated	Mechanism that Promotes Plant Growth	Effect on the Plant	References
<i>Gracilibacillus saliphilus</i> , <i>Staphylococcus petrasii</i> , <i>Bacillus licheniformis</i> , <i>Brevibacterium halotolerans</i>	<i>Zea mays</i>	IAA production increased ACC deaminase activity and EPS formation	Increased growth by higher root-to-shoot ratio and increase in fresh and dry biomass	Aslam et. al., [54]
<i>Bacillus safensis</i> , <i>Bacillus haynesii</i>	<i>Amaranthus viridis</i>	GA, IAA, exopolysaccharides, defence-related enzymatic activities	Growth promotion	Patel et. al., [55]
<i>Hartmannibacter diazotrophicus</i>	<i>Hordeum vulgare</i> L.	Reduced Na <sup>+</sup> uptake by the roots, increased phytohormone production	Growth promotion	Saurez et. al., [56]
<i>Pseudomonas pseudoalcaligenes</i> , <i>Bacillus subtilis</i>	<i>Glycine max</i> L.	ACC-deaminase activity, production of siderophores and IAA	Increase in shoot and root dry weight, higher K <sup>+</sup> /Na <sup>+</sup> in shoot and root and increased proline concentration	Yasmin et. al., [57]
<i>Pantoea ananatis</i>	<i>Oryza sativa</i>	ACC deaminase, P-solubilization, IAA, Siderophore	Higher K <sup>+</sup> /Na <sup>+</sup> ratio in the roots, increased chlorophyll content	Lu et. al., [58]
<i>Sinorhizobium mellilote</i> and <i>Rhizobium leguminosarum</i> b.v <i>phaseoli</i>	<i>Brassica napus</i> L.	IAA, ACC deaminase activity and phosphate solubilizing	Improved growth, higher nutrient uptake and lower Na <sup>+</sup> /K <sup>+</sup> ratio in the roots	Saghafi et. al., [59]
<i>Bacillus</i> sp. (EN1), <i>Zhihengliuella halotolerans</i> (EN3), <i>Bacillus</i> sp. (EN5), <i>Bacillus gibsonii</i> (EN6 and EN10), <i>Oceanobacillus oncorhynchi</i> (EN8), <i>Zhihengliuella</i> sp. (EN12), and <i>Halomonas</i> sp. (IA)	<i>Triticum aestivum</i> L.	Ammonia, IAA and ACC deaminase production, phosphate solubilizing and N <sub>2</sub> fixation	Increased the root and shoot length and total biomass of the plants	Orhan, [60]
<i>Bacillus</i> sp. and <i>Pseudomonas</i> sp.	<i>Musa acuminata</i>	higher levels of K <sup>+</sup> , Ca <sup>2+</sup> , and Mg <sup>2+</sup> and lower levels of Cl <sup>-</sup> and Na <sup>+</sup> , defence-related enzyme activities	Improved growth parameters and development	Kaleh et. al., [61]
<i>Bacillus licheniformis</i> NJ04 strain	<i>Solanum lycopersicum</i> L.	Lower Na <sup>+</sup> /K <sup>+</sup> ratio in the roots	Increased root and shoot length of the plant	James et. al., [62]

## 9. RESEARCH GAPS

Rapid salinization of agricultural land adversely affected our crop yields and posed a serious threat to food security. Global climate change results in the conversion of fertile agricultural land to saline soils every year. Different physical, and chemical amendments and agricultural techniques have been used by farmers from time to time for the reclamation of saline soils but these techniques negatively affect the soil health and also yield potential of the crops. HT-PGPR has proven to be an effective and sustainable solution from time to time for the reclamation of saline soils for crop production [41]. Several PGPR metabolites and genes that respond to salinity stress have been found as a result of approach and method advancements. Still, more research is needed to understand the HT-PGPR's metabolome in both physiologically stressed and unstressed plant interactions [63]. Furthermore, optimal strain selection is important for the proper reclamation of saline soils and investigation is required for the identification of the most effective HT-PGPR for various saline soil types and conditions [64]. A better understanding of the mechanism of tolerance for mitigating stress mediated by halotolerant bacteria and interacting with plants and soil in saline environments to promote growth [65]. Investigating the interactions between introduced halotolerant bacteria and the indigenous soil microbiome is crucial to understand their impact on soil ecology and microbial diversity [66]. Research should focus on assessing the long-term effects of halotolerant bacterial inoculation on soil salinity levels, soil health, and crop productivity to determine the sustainability of this approach [67]. More field-scale experiments are necessary to validate the effectiveness of halotolerant bacterial inoculation under field conditions and to evaluate its practical feasibility and economic viability [68]. Developing scalable and cost-effective strategies for the large-scale application of halotolerant plant growth-promoting bacteria in saline soils is essential for practical implementation and widespread adoption. These research gaps will contribute to the development of more effective and sustainable strategies for remediating saline soils using halotolerant plant growth-promoting bacteria [69].

## 10. CONCLUSION

Changing climate increases the threats of abiotic stresses such as soil salinity which negatively

affects the agriculture system and food security. Salinity negatively affects the plant and soil health posing ionic and osmotic stress to the plants and thereby reducing yields of the crops. Various unsustainable traditional methods have been used by farmers to increase yield under saline soils. Their unsustainable approaches had long-term effects on soil microorganisms and their health. Inoculation of HTPGPR can be a sustainable and reliable approach for the reclamation of saline soils as it improves soil and plant health. Furthermore, areas such as the interaction of HT-PGPR with other soil microbes and the mechanism of salt tolerance to salinity and be explored and more research on this innovative microbial technology is required to control the pandemic of soil salinity and increase crop yield which will help better understanding of HT-PGPR mediated salinity tolerance and its effect on soil health.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. FAO. Global Map of Salt Affected Soils Version 1.0; 2021. Available:<https://www.fao.org/soils-portal/data-hub/soil-maps-and-databases/global-map-of-salt-affected-soils/en/>. Accessed December 2021.
2. Hertel TW, Lobell DB. Agricultural adaptation to climate change in rich and poor countries: Current modeling practice and potential for empirical contributions. *Energy Economics*. 2014;46:562-75.
3. Jamil A, Riaz S, Ashraf M, Foolad MR. Gene expression profiling of plants under salt stress. *Critical Reviews in Plant Sciences*. 2011;30(5):435-58.
4. Mahmood MZ, Odeibat HA, Ahmad R, Gatasheh MK, Shahzad M, Abbasi AM. Low apoplastic Na<sup>+</sup> and intracellular ionic homeostasis confer salinity tolerance upon Ca<sub>2</sub>SiO<sub>4</sub> chemigation in *Zea mays* L. under salt stress. *Frontiers in Plant Science*. 2024;14:1268750.
5. Rengasamy P, de Lacerda CF, Gheyri HR. Salinity, sodicity and alkalinity. In: *Subsoil constraints for crop production* Cham: Springer International Publishing. 2022;83-107.
6. Ehtaiwesh AF. The effect of salinity on nutrient availability and uptake in crop



- plants. *Scientific Journal of Applied Sciences of Sabratha University*. 2022;55-73.
7. AbuQamar SF, El-Saadony MT, Saad AM, Desoky ES, Elrys AS, Abd El-Mageed TA, Semida W, Abdelkhalik A, Mosa WF, Al Kafaas SS, Naser S. Halotolerant Plant Growth-promoting Rhizobacteria Improve Soil Fertility and Plant Salinity Tolerance for Sustainable Agriculture—A Review. *Plant Stress*. 2024;100482.
  8. Bhadreacha P, Singh S, Dwibedi V. A plant's major strength in rhizosphere: The plant growth promoting rhizobacteria. *Archives of Microbiology*. 2023;205(5):165.
  9. Singh A. Soil salinity: A global threat to sustainable development. *Soil Use and Management*. 2022;38(1):39-67.
  10. Okur B, Örçen N. Soil salinization and climate change. In *Climate change and soil interactions*. Elsevier. 2020;331-350
  11. Vinnarasi F, Srinivasamoorthy K, Saravanan K, Gopinath S, Prakash R, Ponnumani G, Babu C. Chemical weathering and atmospheric carbon dioxide (CO<sub>2</sub>) consumption in Shanmuganadhi, South India: Evidences from groundwater geochemistry. *Environmental Geochemistry and Health*. 2021;43:771-90.
  12. Visser JM, Midway S, Baltz DM, Sasser CE. Ecosystem structure of tidal saline marshes. In *Coastal wetlands*. Elsevier. 2019;519-538
  13. Yu X, Xin P, Hong L. Effect of evaporation on soil salinization caused by ocean surge inundation. *Journal of Hydrology*. 2021;597:126200.
  14. Gupta IC, Gupta SK. *Crop production in salt affected soils*. Scientific Publishers; 2019 Apr 1.
  15. Singh A. Soil salinization management for sustainable development: A review. *Journal of Environmental Management*. 2021;277: 111383.
  16. Hopmans JW, Qureshi AS, Kisekka I, Munns R, Grattan SR, Rengasamy P, Ben-Gal A, Assouline S, Javaux M, Minhas PS, Raats PA. Critical knowledge gaps and research priorities in global soil salinity. *Advances in Agronomy*. 2021;169:1-91.
  17. Mohanavelu A, Naganna SR, Al-Ansari N. Irrigation induced salinity and sodicity hazards on soil and groundwater: An overview of its causes, impacts and mitigation strategies. *Agriculture*. 2021;11 (10):983.
  18. Ondrasek G, Rengel Z. Environmental salinization processes: Detection, implications & solutions. *Science of the Total Environment*. 2021;754:142432.
  19. Hailu B, Mehari H. Impacts of soil salinity/sodicity on soil-water relations and plant growth in dry land areas: A review. *J. Nat. Sci. Res*. 2021;12:1-0.
  20. Stavi I, Thevs N, Priori S. Soil salinity and sodicity in drylands: A review of causes, effects, monitoring, and restoration measures. *Frontiers in Environmental Science*. 2021;9:712831.
  21. Kazemi ArPanahi Ū, Mahlooji M, Marashi SK, Maojdam M, Sakinezhad T. Morphophysiological responses of barley genotypes to concentration of Zinc sulphate under drought and salinity conditions. *Crop Science Research in Arid Regions*; 2023.
  22. Abdalla M, Ahmed MA, Cai G, Zarebanadkauri M, Carminati A. Coupled effects of soil drying and salinity on soil-plant hydraulics. *Plant Physiology*. 2022 Oct 1;190(2):1228-41.
  23. Le Bissonnais Y. Soil characteristics and aggregate stability. In *Soil erosion, conservation, and rehabilitation*. CRC Press. 2023;41-60.
  24. Chi W, Nan Q, Liu Y, Dong D, Qin Y, Li S, Wu W. Stress resistance enhancing with biochar application and promotion on crop growth. *Biochar*. 2024;6(1):43.
  25. Singh P, Sharma S, Nisar S, Choudhary OP. Structural stability and organic matter stabilization in soils: Differential impacts of soil salinity and sodicity. *Journal of Soil Science and Plant Nutrition*. 2023;23(2): 1751-73.
  26. Wang H, Zheng C, Ning S, Cao C, Li K, Dang H, Wu Y, Zhang J. Impacts of long-term saline water irrigation on soil properties and crop yields under maize-wheat crop rotation. *Agricultural Water Management*. 2023;286:108383.
  27. Sharma D. *Reclamation and Management of Salt-affected Soils for Crop Production with Special Emphasis on Coastal region: Trend and Prospects*. Souvenir. 2021;11.
  28. Mali BS, S. thengal S, Pate PN. Physico-chemical characteristics of salt affected soil from Barhanpur, MS, India; 2021.
  29. Hiranmai RY, Neeraj A. Assessment of salt-affected soil in Indo-Gangetic Plain of Uttar Pradesh and prediction using regression model. *Land Degradation & Development*. 2024;35(3):1087-97.

30. Raza A, Tabassum J, Fakhar AZ, Sharif R, Chen H, Zhang C, Ju L, Fotopoulos V, Siddique KH, Singh RK, Zhuang W. Smart reprogramming of plants against salinity stress using modern biotechnological tools. *Critical Reviews in Biotechnology*. 2023; 43(7):1035-62.
31. Lu Y, Fricke W. Salt Stress—Regulation of root water uptake in a whole-plant and diurnal context. *International Journal of Molecular Sciences*. 2023;24(9):8070.
32. Hnilickova H, Kraus K, Vachova P, Hnilicka F. Salinity stress affects photosynthesis, malondialdehyde formation, and proline content in *Portulaca oleracea* L. *Plants*. 2021;10(5):845.
33. Wungrampha S, Rawat N, Singla-Pareek SL, Pareek A. Survival strategies in halophytes: adaptation and regulation. In *Handbook of Halophytes: From Molecules to Ecosystems towards Biosaline Agriculture* Cham: Springer International Publishing. 2020;1-22.
34. Hasanuzzaman M, Fujita M. Plant responses and tolerance to salt stress: Physiological and molecular interventions. *International Journal of Molecular Sciences*. 2022;23(9):4810.
35. Sachdev S, Ansari SA, Ansari MI, Fujita M, Hasanuzzaman M. Abiotic stress and reactive oxygen species: Generation, signaling, and defense mechanisms. *Antioxidants*. 2021;10(2):277.
36. Chourasia KN, Lal MK, Tiwari RK, Dev D, Kardile HB, Patil VU, Kumar A, Vanishree G, Kumar D, Bhardwaj V, Meena JK. Salinity stress in potato: Understanding physiological, biochemical and molecular responses. *Life*. 2021;11(6):545.
37. Farooqi ZU, Qadir AA, Ilić P, Zeeshan N, Tunguz V, Pržulj N. Restoration and preservation of degraded soils for crop production. *ОДРЖИВИ РАЗВОЈ И УПРАВЉАЊЕ ПРИРОДНИМ РЕСУРСИМА РЕПУБЛИКЕ СРПСКЕ*. 2023;9(9).
38. Shahid M, Khan MS, Singh UB. Pesticide-tolerant microbial consortia: Potential candidates for remediation/clean-up of pesticide-contaminated agricultural soil. *Environmental Research*. 2023; 116724.
39. Zhang P, Jiang Z, Wu X, Lu Q, Lin Y, Zhang Y, Zhang X, Liu Y, Wang S, Zang S. Effects of biochar and organic additives on CO<sub>2</sub> emissions and the microbial community at two water saturations in saline–alkaline soil. *Agronomy*. 2023;13(7): 1745.
40. Peng M, Jiang Z, Xiang Z, Zhou A, Wang C, Wang Z, Zhou F. Genomic features of a plant growth-promoting endophytic *Enterobacter cancerogenus* JY65 dominant in microbiota of halophyte *Suaeda salsa*. *Plant and Soil*. 2024;496 (1):269-87.
41. Arora NK, Fatima T, Mishra J, Mishra I, Verma S, Verma R, Verma M, Bhattacharya A, Verma P, Mishra P, Bharti C. Halo-tolerant plant growth promoting rhizobacteria for improving productivity and remediation of saline soils. *Journal of Advanced Research*. 2020;26:69-82.
42. Channab BE, Idrissi AE, Ammar A, Dardari O, Marrane SE, El Gharrak A, Akil A, Essemli Y, Zahouily M. Recent advances in nano-fertilizers: synthesis, crop yield impact, and economic analysis. *Nanoscale*; 2024.
43. Kumawat C, Kumar A, Parshad J, Sharma SS, Patra A, Dogra P, Yadav GK, Dadhich SK, Verma R, Kumawat GL. Microbial diversity and adaptation under salt-affected soils: A review. *Sustainability*. 2022;14(15): 9280.
44. Arora NK, Fatima T, Mishra J, Mishra I, Verma S, Verma R, Verma M, Bhattacharya A, Verma P, Mishra P, Bharti C. Halo-tolerant plant growth promoting rhizobacteria for improving productivity and remediation of saline soils. *Journal of Advanced Research*. 2020;26:69-82.
45. Agbodjato NA, Babalola OO. Promoting sustainable agriculture by exploiting plant growth-promoting rhizobacteria (PGPR) to improve maize and cowpea crops. *PeerJ*. 2024;12:e16836.
46. Sharma PK, Kumar S. Soil Structure and Plant Growth. In *Soil Physical Environment and Plant Growth: Evaluation and Management* Cham: Springer International Publishing. 2023;125-154.
47. Kumar V, Raghuvanshi N, Pandey AK, Kumar A, Thoday-Kennedy E, Kant S. Role of halotolerant plant growth-promoting rhizobacteria in mitigating salinity stress: Recent advances and possibilities. *Agriculture*. 2023;13(1):168.
48. Abdullatif Y, Soliman E, Hammad SA, El-Ghamry A, Mansour MM. Impact of Carbon Nanoparticles on Aggregation and Carbon Sequestration under Soil Degradation—a Review. *Journal of Soil Sciences and Agricultural Engineering*. 2024;15(4):85-92.

49. Santra HK, Banerjee D. Microbial extracellular polymeric substance: function and role against environmental stress. In *Bacterial Secondary Metabolites*. Elsevier. 2024;83-106.
50. Sandhya V, Ali SZ. The production of exopolysaccharide by *Pseudomonas putida* GAP-P45 under various abiotic stress conditions and its role in soil aggregation. *Microbiology*. 2015;84:512-9.
51. Guhra T, Stolze K, Totsche KU. Pathways of biogenically excreted organic matter into soil aggregates. *Soil Biology and Biochemistry*. 2022;164:108483.
52. Ali N, Abbas SA, Sharif L, Shafiq M, Kamran Z, Haseeb M, Shahid MA. Microbial extracellular polymeric substance and impacts on soil aggregation. In *Bacterial Secondary Metabolites*. Elsevier. 2024;221-237.
53. Islam S, Akanda AM, Prova A, Islam MT, Hossain MM. Isolation and identification of plant growth promoting rhizobacteria from cucumber rhizosphere and their effect on plant growth promotion and disease suppression. *Frontiers in Microbiology*. 2016;6:165532.
54. Aslam F, Ali B. Halotolerant bacterial diversity associated with *Suaeda fruticosa* (L.) forssk. improved growth of maize under salinity stress. *Agronomy*. 2018;8(8): 131.
55. Patel M, Vurukonda SS, Patel A. Multi-trait halotolerant plant growth-promoting bacteria mitigate induced salt stress and enhance growth of *Amaranthus viridis*. *Journal of Soil Science and Plant Nutrition*. 2023;23(2):1860-83.
56. Suarez C, Cardinale M, Ratering S, Steffens D, Jung S, Montoya AM, Geissler-Plaum R, Schnell S. Plant growth-promoting effects of *Hartmannibacter diazotrophicus* on summer barley (*Hordeum vulgare* L.) under salt stress. *Applied Soil Ecology*. 2015;95:23-30.
57. Yasmin H, Naeem S, Bakhtawar M, Jabeen Z, Nosheen A, Naz R, Keyani R, Mumtaz S, Hassan MN. Halotolerant rhizobacteria *Pseudomonas pseudoalcaligenes* and *Bacillus subtilis* mediate systemic tolerance in hydroponically grown soybean (*Glycine max* L.) against salinity stress. *PLoS One*. 2020;15(4):e0231348.
58. Lu L, Chang M, Han X, Wang Q, Wang J, Yang H, Guan Q, Dai S. Beneficial effects of endophytic *Pantoea ananatis* with ability to promote rice growth under saline stress. *Journal of Applied Microbiology*. 2021;131 (4):1919-31.
59. Saghafi D, Ghorbanpour M, Lajayer BA. Efficiency of Rhizobium strains as plant growth promoting rhizobacteria on morpho-physiological properties of *Brassica napus* L. under salinity stress. *Journal of Soil Science and Plant Nutrition*. 2018;18 (1):253-68.
60. Orhan F. Alleviation of salt stress by halotolerant and halophilic plant growth-promoting bacteria in wheat (*Triticum aestivum*). *Brazilian Journal of Microbiology*. 2016;47:621-7.
61. Kaleh AM, Singh P, Mazumdar P, Wong GR, Chua KO, Harikrishna JA. A halotolerant plant growth promoting consortium of *Bacillus* sp. RB3 and *Pseudomonas* sp. EB3 primes banana, *Musa acuminata* cv. Berangan, against salinity and Foc-TR4 stresses. *Current Plant Biology*. 2023;35:100294.
62. James N, Umesh M, Sarojini S, Shanmugam S, Nasif O, Alharbi SA, Chi NT, Brindhadevi K. Unravelling the potential plant growth activity of halotolerant *Bacillus licheniformis* NJ04 isolated from soil and its possible use as a green bioinoculant on *Solanum lycopersicum* L. *Environmental Research*. 2023;216:114620.
63. Al-Turki A, Murali M, Omar AF, Rehan M, Sayyed RZ. Recent advances in PGPR-mediated resilience toward interactive effects of drought and salt stress in plants. *Frontiers in Microbiology*. 2023;14: 1214845.
64. Teo HM, Bhubalan K, CI MS, Ng LC. Setting a plausible route for saline soil-based crop cultivations by application of beneficial halophyte-associated bacteria: A review. *Microorganisms*. 2022; 10(3):657.
65. Hernández-Canseco J, Bautista-Cruz A, Sánchez-Mendoza S, Aquino-Bolaños T, Sánchez-Medina PS. Plant growth-promoting halobacteria and their ability to protect crops from abiotic stress: An eco-friendly alternative for saline soils. *Agronomy*. 2022;12(4):804.
66. Ma Y, Zheng C, Bo Y, Song C, Zhu F. Improving crop salt tolerance through soil legacy effects. *Frontiers in Plant Science*. 2024;15:1396754.

67. Phour M, Sindhu SS. Soil salinity and climate change: Microbiome-based strategies for mitigation of salt stress to sustainable agriculture. In Climate change and microbiome dynamics: carbon cycle feedbacks Cham: Springer International Publishing. 2023; 191-243.
68. Malik A, Garg VK. Bioremediation for Sustainable Environmental Cleanup. CRC Press; 2024.
69. Arora S. Microbial approaches for bio-amelioration and management of salt affected soils. Soil Science: Fundamentals to Recent Advances. 2021; 433-47.

---

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/118111>