



# Myocardial Infarction Due to Coronary Artery Embolism in a Patient with a Mechanical Mitral Valve: A Case Report

Sara Abouradi <sup>a\*</sup>, Asmaa Elamraoui <sup>a</sup>, Zineb Eljaouhari <sup>a</sup>,  
Salim Arrous <sup>a</sup>, Ghali Bennouna <sup>a</sup> and Rachida Habbal <sup>a</sup>

<sup>a</sup> *Department of Cardiology, Ibn Rochd Hospital University, Casablanca, Morocco.*

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96333>

**Case Study**

**Received: 09/12/2022**

**Accepted: 16/02/2023**

**Published: 24/02/2023**

## **ABSTRACT**

**Background:** Mechanical valves are inherently thrombogenic and require meticulous anticoagulation. Thrombosis of cardiac valve prostheses is a dangerous complication, that practitioners should clearly explain to their patients the potential risks of stopping treatment.

**Case Presentation:** A 48-year-old female patient without any cardiovascular risk factor with a history of mechanical prosthesis for rheumatic mitral stenosis, who stopped anticoagulant treatment for dental extraction, admitted for Ischemic cerebral vascular accident complicated during her hospitalization by supraventricular tachycardia with a coronary embolism leading to an ST segment elevation due to a non obstructive thrombus causing embolic acute myocardial infarction.

**Conclusions:** Education regarding an adequate anticoagulation in these patients is important. Clinicians should keep in mind the importance of providing to their patients all the informations concerning the potential risks of stopping their treatment.

*Keywords: Mechanical valves; thrombosis of prosthesis; anticoagulation; case report.*

\*Corresponding author: Email: saraabouradi17@gmail.com;

## ABBREVIATIONS

VKA : Antivitamine K  
GCS : Glasgow Coma Scale  
ECG : Electrocardiogram  
LV : Left Ventricle  
LVEF : Left Ventricular Ejection Fraction

## 1. INTRODUCTION

Thrombosis of cardiac valve prostheses is a serious and potentially lethal complication. Its diagnosis must be made early by analysing clinical and echocardiographic results [1,2,3], particularly the transoesophageal one.

The surgical intervention, often performed in an emergency situation, is related to a high mortality percentage [4].

Prevention of valve thrombosis is essential. It is based on a strict control of the anticoagulant treatment. Both clinician and biologist doctor are responsible for the management of all patients with a mechanical valve. "Although mechanical valves have long term durability, yet the incidence of malfunction is not negligible (0.1-6%) and mechanical prosthetic valve thrombosis is one of those serious complications, where patients often deteriorate rapidly, compromising management outcomes" [5,6].

## 2. CASE PRESENTATION

A 48-year-old female patient with no cardiovascular risk factors, with a history of mitral valve replacement for rheumatic mitral stenosis, who stopped by herself the anticoagulant treatment for dental extraction for five days.

The patient was admitted to the cardiology department for Ischemic cerebral vascular accident.

Clinical examination on admission revealed a confused patient GCS 14/15, hemiplegic, hemodynamically and respiratory stable : blood pressure at 135/75mmHg, heart rate at 80bpm, SpO2 at 96% without oxygenation, cardiac auscultation was normal, and the prosthetic click audible, a transthoracic echocardiography was performed showing no prosthetic thrombus. Regarding the blood tests results, we hadn't notice any abnormalities, the cerebral CT scan has confirmed the presence of a left frontal and a right occipital ischemic vascular accident.

The patient received aspirin 75mg per day , fluoxet 20mg per day, and a placement of a feeding nasogastric probe. We have contacted the neurologist doctor, we have decided to stop the anticoagulation. After 48h of hospitalization, we have performed a cerebral CT scan showing a non surgical hemorrhagic remodeling becoming a factor of poor prognosis.

The day after, the patient had presented a sudden supraventricular tachycardia with hemodynamic instability (Fig. 1) shocked three times, and immediately after that, the electrocardiogram showed a ST-segment elevation in inferior derivation (Fig. 2).

The echocardiographic control had showed a homogeneous hyperechoic formation attached to the mitral prosthesis related to a thrombus (Fig. 3).

The patient subsequently had presented a cardiorespiratory arrest that was not recovered after 30 min of cardiopulmonary resuscitation.





Fig. 1. Electrocardiogram showing a supraventricular tachycardia



**Fig. 2. Electrocardiogram showing the st segment elevation in inferior derivation**



**Fig. 3. Apical section four cavities objectifying the thrombus on the mitral prosthesis**

### 3. DISCUSSION

Coronary embolism MI is a rare disease. Most emboli (75%) involve the left reticular formation [1,4]. "This is due to its larger caliber and the implantation angle of the left coronary artery is smaller than the acute angle of the right coronary artery" [7]. "Mechanical heart valves are thrombosed, and anticoagulation is essential to prevent thromboembolism and acute thrombotic obstruction [8]". Coronary embolism is a common cause of acute transmural myocardial infarction,

Occurs in 10% to 13% of autopsy series [9] and represents an important entity in etiology and clinical management [10].

"The consequences of coronary embolism leading to myocardial infarction depend on the size of the embolus and the lumen diameter of the vessel" [9,11].

"Coronary embolism should always be considered, especially in patients with sudden chest pain in patients with prosthetic valves,

chronic atrial fibrillation, dilated cardiomyopathy, infective endocarditis, intracardiac shunts, cardiac myxomas, mural thrombi, and hypercoagulable states case" [11].

Major embolic events occurred in 1% of patients with valve prostheses treated with warfarin [10]. Among serial coronary embolisms in patients with valvular heart disease at autopsy, 8% occurred in patients with prostheses [12]. The risk of embolism is greater with mitral valve prostheses, rigid prostheses, or multiple valve prostheses [13,14].

Coronary embolism as a cause of myocardial infarction is an uncommon but important entity both in terms of aetiology and treatment. Previous cases of coronary emboli in association with prosthetic mechanical valves have been reported previously [15] but the mechanism of pharmacology and lack of patient awareness of medication importance is quite unique in this case.

Inadequate anticoagulation is the most frequent factor involved [15,8], which was found in our patient. Therapeutic education of patients with prosthetic valves is one of the major preventive strategies for these serious, sometimes fatal, embolic events.

"Embolism myocardial infarction is underdiagnosed and it is important to diagnose the source of embolism and treat the cause" [16,15].

Patient education is vital in our battle to prevent this entity in high-risk patients as in our case.

#### 4. CONCLUSIONS

Myocardial infarction by coronary embolism from mitral prosthesis thrombosis is probably an underestimated situation and often occurs in the presence of ineffective anticoagulation. Therapeutic education is necessary to prevent these complications.

#### CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

#### ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Wenger KN, Bauer S. Coronary embolism. review of the literature and presentation of 15 cases. *Am J Med.* 1958;25:549-57.
2. Alpert JS. The thrombosed prosthetic valve. *J Am Coll Cardiol* 2003;41:659-60.
3. Tong TA, Roudout R, Ozkan M, et al. On behalf of the prosthetic valve thrombolysis – role of transoesophageal echocardiography (pro-tee) registry investigators/ transoesophageal echocardiography improves risk assessment of thrombolysis of prosthetic valve thrombosis: Results of the international pro-tee registry. *J Am Coll Cardiol.* 2004;43:77-84.
4. Prizel KR, Hutchins GM, Bulkley BH. Coronary artery embolism and myocardial infarction, a clinicopathologic study of 55 patients. *Ann Intern Med.* 1978;88:155-61.
5. Hassouna A, El-Ghanam M, Moftah H, Samir K, Refaat K. Index of deterioration of patients with mechanical prosthetic heart valve thrombosis. *The Cardiothoracic Surgeon.* 2020;28(1):1-1.
6. Bezanjani FN, Gohari S, Bassiri HA, Ahangar H, Reshadmanesh T. Risk factors associated with heart valve thrombosis in patients with prosthetic heart valve dysfunction. *Archives of Iranian Medicine.* 2020;23(9):600-4.
7. Gully C, Baron O, Nguyen Khal O. et al, Coronary embolism: a not so rare cause of myocardial infarction? Review of the literature about five suspected cases of patients in atrial fibrillation. *Ann Cardiol Angeiol.* 2008;57:290-94.
8. Cannegiester SC, Rosendaal FR, Brier E. Thromboembolic and bleeding complications in patients with mechanical heart valve prosthesis. *Circulation.* 1994;89:635-41.
9. Aslam MS, Sanghi V, Hersh S, Lakier JB. Embolie coronarienne en selle et infarctus du myocarde chez un patient porteur d'une valve mitrale prothétique. *Cathétérisme et Interventions Cardiovasculaires.* 2002;57(3):367–370.
10. Charles RG, Epstein EJ, Holt S, Coulshed N. Coronary embolism in valvular heart disease. *Q J Med.* 1982;51:147-61

11. Waller BF, Dixon DS, Kim RW, Roberts WC. "Embolus to the left main coronary artery." *Journal Américain De Cardiologie*. 1982;50(3):658–660.
12. Cannegiester SC, Rosendaal Fr, Winten AR, Van der Meer FJM, Vandenbrouke JP, Brier E. Optimal oral anticoagulant therapy in patients with mechanical heart valves. *NEJM*. 1995;333:11-7.
13. Srilakshmi MA, Shetty GG, santosh MJ, et al. Acute st elevation myocardial infarction in a patient with caged ball mitral valve prosthesis: A case report. *International Journal of Cardiology*. 2008;128:e101–e103.
14. M. Dogan, S. Acikel, Aksoy, et al. Embolie coronarienne en selle provoquant un infarctus du myocarde chez un patient MMN porteur d'une prothèse de valve mitrale mécanique : traitement par thérapie thrombolytique. *Journal International De Cardiologie*, 2009;135(2):e47-e48.
15. Kiernan TJ, Flynn AM, Kearney P. Coronary embolism causing myocardial infarction in a patient with mechanical aortic valve prosthesis. *International Journal of Cardiology*. 2006;112:e14–e16.
16. Lavoie P, Leduc L, Mercier LA. Infarctus du myocarde embolique chez une femme enceinte porteuse d'une valve cardiaque mécanique sous héparine de bas poids moléculaire. *Journal Canadien De Cardiologie*. 2004;20(9):917–919.

© 2023 Abouradi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/96333>