



# **Phytochemicals against COVID -19, Pharmacological Perspectives: A Systematic Review**

**Ahmed S. Ali <sup>a,b</sup>, Ekram A. Saied <sup>c\*</sup>, Sharifa M. Theeban <sup>d#</sup>,  
Mazen M. Humayran <sup>e<sup>o</sup></sup>, Assmaa Ahmed Shaker <sup>f</sup>,  
Mohammed S. Almuhayawi <sup>g,h</sup> and Abdelbagi Alfadil <sup>g</sup>**

<sup>a</sup> Department of Pharmacology, Faculty of Medicine, King Abdulaziz University, KSA.

<sup>b</sup> Department of Pharmaceutics, Faculty of Pharmacy, Assiut University, Egypt.

<sup>c</sup> Department of Clinical Pharmacy and Pharmacology, Ibn Sina National College for Medical Studies, Jeddah, KSA.

<sup>d</sup> Jeddah International College, Jeddah, KSA.

<sup>e</sup> Faculty of Medicine, Medical Intern at King Abdulaziz University Hospital/King Abdulaziz University, Jeddah, KSA.

<sup>f</sup> Ibn Sina National College for Medical Studies, Jeddah, KSA.

<sup>g</sup> Department of Medical Microbiology and Parasitology, Faculty of Medicine, King Abdulaziz University, Jeddah, KSA.

<sup>h</sup> Y. A. Jameel Scientific Chair of Prophetic Medical Applications, King Abdulaziz University, KSA.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/JPRI/2021/v33i53B33678

### Editor(s):

- (1) Dr. Fahmida Khan, National Institute of Technology Raipur, India.
- (2) Dr. P. Veera Muthumari, V.V. Vanniaperumal College for Women, India.
- (3) Dr. Sung-Kun Kim, Northeastern State University, USA.

### Reviewers:

- (1) Mark M. Alipio, Philippine Christian University, Philippines.
- (2) Rajesh Kumar, Rajasthan University of Veterinary and Animal Sciences (RAJUVAS), India.
- (3) A. Vijaya Anand, Bharathiar University, India.
- (4) Veeresh Yadav, N.E.T. Pharmacy College, India.
- (5) G. Bhuvaneshwari, Saveetha Institute of Medical and Technical Sciences, India.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/76654>

**Review Article**

**Received 29 October 2021**

**Accepted 29 November 2021**

**Published 06 December 2021**

# Lecturer;

<sup>o</sup> MBBS Student;

\*Corresponding author: E-mail: ekramnemr@ibnsina.edu.sa;

## ABSTRACT

**Background:** Many publications discussed the potential role of medicinal plants in the management of COVID-19. However, clinical studies of the efficacy and safety of specified phytochemical(s) are limited.

**Objectives:** To explore the pharmacological profile of specified compounds against COVID-19

**Method:** Systematic literature search of academic databases to explore specified phytochemicals for the management of COVID-19 using appropriate search terms. Rayyan software was used to organize 786 citations of which, 236 articles were included in this review.

**Results:** Initially 70 compounds were identified to have a potential role in the management of COVID-19. In this review, 18 compounds were selected for further search

**Conclusion:** In vitro anti-SARS-CoV-2 activity has been demonstrated for a variety of natural compounds. However, preclinical research for most phytochemicals is scarce, and only a few compounds have been evaluated in clinical trials against COVID-19. A comprehensive pharmacological profile of these phytochemicals is urgently needed.

*Keywords: Phytochemicals; COVID-19; SARS-CoV-2; flavonoids; alkaloids; essential oils.*

## 1. INTRODUCTION


Coronavirus disease 2019 (COVID-19), which was first reported in Wuhan, China, in December 2019, is an ongoing global pandemic, highly infective respiratory viral illness. It is caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). It leads to extreme human and economic losses. As of 29 June 2021, more than 181 million cases have been identified, with more than 3.92 million confirmed deaths [1]. More details are available in these reviews [2-4]. The development of diagnostic tests, vaccinations, and medications for COVID-19 management has received a lot of attention [5].

Numerous meta-analyses have shown the potential beneficial role of medicinal plants in reducing the severity of COVID-19 when combined with standard pharmacological therapy [6-30]. Most clinical trials involved Chinese herbs [10, 12-15, 17, 18, 20, 22-29], few studies discussed the role of herbal medicine in specific countries, such as Nigeria [16] or India, [19]. Some clinical trials focused on a single plant or food as bee products [7, 11], plant-derived food-grade substances [21], Lianhua Qingwen (a Chinese herbal medicine) [28, 30]. However, it should be noted that integrated research on isolated phytochemicals (pure substances) is sparse, which may explain why most decision-makers are reluctant to study the efficacy and safety of phytochemicals in COVID-19 clinical studies. Therefore, the goal of this systemic review is to explore the potential role of phytochemicals in the management of COVID-19.

## 2. METHODS

Two databases were used, namely PubMed (NCBI), Lit-COVID 19, last accessed in November 2021. The research was restricted to 2019-2021, full text, and English language. All citations were imported to endnote 8, duplicates were excluded. Rayyan program (for Systemic review) was used to organize and select the relevant articles [31]. PRISMA flow chart is shown in Fig. A [32].

The first objective was to identify phytochemical(s) for the management of COVID-19. The 2<sup>nd</sup> objective explores the efficacy and safety, pharmacology of specified phytochemicals. The study was conducted in two phases, search terms in phase one were: # 1 "SARS-CoV-2" OR "COVID-19", OR "coronavirus". # 2: "Phytochemical\*" OR "medicinal plants", OR "herbal medicine" followed by advanced (combined) search: #1 AND#2. 70 compounds were identified to have potential activity against the coronavirus (Tables 1-3). 18 compounds were selected by two investigators: given their low IC<sub>50</sub> against SARS-CoV-2, immunomodulating effect and good safety profile. Key terms in phase 2: each of the phytochemicals e.g. # 3Thymoquinone, 4 # "clinical trial" OR "antiviral" OR "anti-inflammatory" OR "lung injury" OR "Pharmacokinetic" OR "Formulation" OR "safety". Followed by advanced search #3 AND #4.



**Fig. A. PRISMA flow chart of the search for safety and efficacy of specified phytochemicals against COVID 19**

\*1- Only journals were included, 2- Only review articles, systemic reviews, clinical trials were included

#### Abbreviations and glossary of terms

ACE2	: Angiotensin converting enzyme-2.
ALI	: Acute lung injury.
COVID-19	: Coronavirus disease 2019.
CPE	: Cell-protective effects.
FDA	: Food and Drug Administration.
IC <sub>50</sub> /EC <sub>50</sub> values	: The 50% inhibitory and effective concentration respectively.
LPS	: Lipopolysaccharide.
M Wt.	: Molecular weight.
Mpro	: Main protease.
PK	: Pharmacokinetics; describes absorption, distribution, metabolism, and elimination of drugs.
PLpro	: Papain like proteases, and 3CLpro: 3-chymotrypsin like protease: enzymes essential for viral replication.

SARS	: Severe acute respiratory syndrome.
SARS-CoV or SARS-CoV-1	: severe acute respiratory syndrome coronavirus 1, is a strain of virus that causes SARS.
SARS-CoV-2	: severe acute respiratory syndrome coronavirus 2, that causes COVID-19.


### 3. POTENTIAL ROLE OF PHYTO-CHEMICALS IN THE MANAGEMENT OF COVID-19


#### 3.1 Overview

A comprehensive web-based resource for natural products (about 200 compounds) was launched in 2020 [33]. Several review articles included an extensive search for medicinal plant extracts, compounds that have been identified and proven to suppress the life cycle of the coronavirus. These reviews summarized the IC<sub>50</sub>, selectivity, mechanisms of action and structure of specified compounds as polyphenols [34-36], flavonoids [37,38] of which myricetin and scutellarin [39] were highlighted. Alkaloids [40,41] particularly lycorine [42,43] was of considerable interest, other alkaloids include tetrandrine, tanghinoline, cepharanthine [44] and reserpine [45]. Other classes of phytochemicals include: sterols [46],

terpenoids [47,48], stilbenoid, resveratrol [49], triterpene glycoside (saponin) glycyrrhizin [50], and saponin (Aescin) [45]. Tannins (tannic acid), 3-isothaflavin-3-gallate, theaflavin-3,3'-digallate [51], and essential oils [52-54].

Many publications provided an overview of the utility of phytochemicals against COVID-19 [55-57]. Some provided a detailed description of a specified compound(s) e.g., curcumin [58], valinomycin [59], tanshinones [60], resveratrol [61] and glycyrrhizin [62]; or plant e.g. *Nigella sativa* [63]. A non-comprehensive list (about 70 compounds) was provided in Tables 1-3. From this list, 18 compounds were selected for further search to explore their pharmacological profile (their structure is provided in Fig. 1). The selection was based on low IC<sub>50</sub>, immunomodulating effect and known safety profile by two authors.





**Fig. 1. Structure of selected phytochemicals of a potential role in the management of COVID-19**

**Table 1. Compounds suggested interacting with the specified target(s) of SARS-CoV-2: Spike protein, 3CLpro or PLpro [33]**

Compound	Class/ source	mechanism
Betulinic acid	Pentacyclic triterpenoid	3CLpro
Dihomo- $\gamma$ -linolenic acid	$\omega$ -6 fatty acid	3CLpro
Dihydrotanshinone	<i>Salvia miltiorrhiza</i>	Spike protein
Tanshinonella	<i>Salvia miltiorrhiza</i>	PLpro and 3CLpro
Cryptotanshinone	Quinone	PLpro and 3CLpro
Lignan	Low M Wt. polyphenols	3CLpro
Moupinamide	A phenolic amide	PLpro
N-cis-feruloyl tyramine	A phenolic amide	PLpro and 3CLpro
Coumaroyl tyramine	A phenolic amide	PLpro and 3CLpro
Quercetin	Flavonol	PLpro and 3CLpro
Kaempferol	Flavonol	PLpro and 3CLpro
Sugiol	A diterpene	3CLpro

### 3.2 Pharmacological Profile of Selected Phytochemicals

Out of 18 selected phytochemicals (Fig. 1), two clinical trials were found, the 1<sup>st</sup> trial revealed that resveratrol, reduces angiotensin converting enzyme-2 (ACE2) expression in human adipose tissue [75]. The 2<sup>nd</sup> trial documented that, nano-curcumin treatment is a promising technique for controlling inflammatory cytokines in COVID-19

patients [76]. Aescin (was registered in China and Italy for clinical trials to demonstrate its potential role as an adjuvant in the management of COVID-19 [77].

The following is a summary of what has been collected regarding the pharmacology profile of these compounds, with a focus on their potential role against COVID-19.

**Table 2. Example of compounds with IC<sub>50</sub>/EC<sub>50</sub> values tested against an infectious coronavirus [64]**

Compound	Chemical group	IC <sub>50</sub> /EC <sub>50</sub> values	Coronavirus type targeted
Glycyrrhizin	Saponin	EC <sub>50</sub> = 364.5 μM	SARS-CoV
Tetra-O-galloyl-β-D-glucose	Polyphenol	EC <sub>50</sub> = 4.5 μM	SARS-CoV
Luteolin	Flavonoid	EC <sub>50</sub> = 10.6 μM	SARS-CoV
Sinigrin	Polyphenol	IC <sub>50</sub> = 217 μM	SARS-CoV
β-Sitosterol	Phytosterol	IC <sub>50</sub> = 1210 μM	SARS-CoV
Hesperetin	Flavonoid	IC <sub>50</sub> = 8.3 μM	SARS-CoV
Amentoflavone	Flavonoid	IC <sub>50</sub> = 8.3 μM	SARS-CoV
Luteolin	Flavonoid	IC <sub>50</sub> = 20.2 μM	SARS-CoV
Quercetin	Flavonoid	IC <sub>50</sub> = 23.8 μM	SARS-CoV
Isobavachalcone	Flavonoid	IC <sub>50</sub> = 7.3 μM	SARS-CoV
Psoralidin	Flavonoid	IC <sub>50</sub> = 4.2 μM	SARS-CoV
Tomentin A	Flavonoid	IC <sub>50</sub> = 6.2 μM	SARS-CoV
Tomentin B	Flavonoid	IC <sub>50</sub> = 6.1 μM	SARS-CoV
Tomentin E	Flavonoid	IC <sub>50</sub> = 5.0 μM	SARS-CoV
3'-O-Methyldiplacol	Flavonoid	IC <sub>50</sub> = 9.5 μM	SARS-CoV
Isoliquiritigenin	Flavonoid	IC <sub>50</sub> = 61.9 μM	SARS-CoV
Kaempferol	Flavonoid	IC <sub>50</sub> = 116.3 μM	SARS-CoV
Kazinol F	Flavonoid	IC <sub>50</sub> = 43.3 μM	SARS-CoV
Broussouchalcone B	Flavonoid	IC <sub>50</sub> = 57.8 μM	SARS-CoV
Papyriflavonol A	Flavonoid	IC <sub>50</sub> = 103.6 μM	SARS-CoV
Terrestriamine	Cinnamic amide	IC <sub>50</sub> = 15.8 μM	SARS-CoV
Blancoxanthone	Xanthone	EC <sub>50</sub> = 3 μg/ml	HCoV 229E
Pyranojacareubin	Xanthone	EC <sub>50</sub> = 15 μg/ml	HCoV 229E
Lycorine	Alkaloid	EC <sub>50</sub> = 15.7 IU/ml	SARS-CoV
Tingenone	Triterpene	IC <sub>50</sub> = 9.9 μM	SARS-CoV
Iguesterin	Triterpene	IC <sub>50</sub> = 2.6 μM	SARS-CoV
Pristimererin	Triterpene	IC <sub>50</sub> = 5.5 μM	SARS-CoV
Dihydrotanshinone I	Diterpene	IC <sub>50</sub> = 4.9 μM	SARS-CoV
Cryptotanshinone	Diterpene	IC <sub>50</sub> = 0.8 μM	SARS-CoV
Tanshinone IIA	Diterpene	IC <sub>50</sub> = 1.6 μM	SARS-CoV
Xanthoangelol	Chalcone	IC <sub>50</sub> = 11.4 μM	SARS-CoV
Hirsutenone	Diarylheptanoid	IC <sub>50</sub> = 3.0 μM	SARS-CoV
Rubranoside	Diarylheptanoid	IC <sub>50</sub> = 7.2 μM	SARS-CoV
Curcumin	Diarylheptanoid	IC <sub>50</sub> = 5.7 μM	SARS-CoV
FIt3 receptor-interacting lectin (FRIL).	Lectin	EC <sub>50</sub> = 6.25 μg/ml	SARS-CoV-2
Abrus precatorius (APA) Lectin	Lectin	EC <sub>50</sub> = 0.45 μg/ml	SARS-CoV
Urtica dioica agglutinin (UDA) Lectin	Lectin	EC <sub>50</sub> = 1.3 μg/ml	SARS-CoV

Other compounds are not listed above. Resveratrol, EmodinAescin, Ginsenoside-Rb1, Leptodactylone, Celastrol, Tetrandrine, Cephalaranthine, S-aikosaponinB<sub>2</sub>, Quercetin-3-b-galactoside, Chalcones, Theaflavin, Myricetin, Scutellarein[40, 65-70].

Essential oils with potential antiviral and immunomodulating activity, 8-cineole (eucalyptol), allyl sulfide, farnesene, farnesol, nerolidol; eugenol, menthol, carvacrol, cinnamaldehyde [71-74]

### 3.2.1 Thymoquinone

It is a quinone compound found in the seeds of *Nigella sativa* (*Ranunculaceae*) It has a broad pharmacological activity, and a good safety profile [78]. Its potential role in the management of COVID -19 was suggested based on its antiviral activity and ability to attenuate lung injury and inflammation [79-82]. Thymoquinone has poor oral bioavailability [83] likely due to its

low solubility in water. Pharmaceutical formulations and nano delivery systems were suggested to improve their bioavailability and distribution [84-86].

### 3.2.2 Lycorine

Lycorine is a phenanthridine alkaloid isolated from the bulbs of *Lycoris radiata* (*Amaryllidaceae*). It has broad pharmacological

activity [87], it has potent activity against SARS-CoV [42]. Its ability to suppress SARS-CoV-2 replication was confirmed in Viro-6 cells, the EC<sub>50</sub> of lycorine, and chloroquine were, 0.18 and 1.36  $\mu$ M, respectively, CC value >40  $\mu$ M [69]. An animal study showed its very high volume of distribution and short half-life [88]. More, pre-clinical investigations are needed to determine other PK parameters such as protein binding and metabolism. Lycorine, in particular, has a wide range of pharmacological effects on a variety of disorders while having relatively low toxicity and minimal side effects [89].

### 3.2.3 Glycyrrhizin

It is triterpene glycoside (saponin), which is also known as glycyrrhizic acid isolated from the roots of *Glycyrrhiza glabra* (*Fabaceae*). It has several pharmacological activities [90]. The compound or its active metabolite expresses antiviral effects against SARS-CoV in cell culture [91]. *In vitro* studies demonstrated that glycyrrhizin potently neutralizes SARS-CoV-2 by inhibiting the viral main protease (Mpro) [92]. It exhibited anti-inflammatory potential by several mechanisms including suppression of several proinflammatory mediators [93]. Its low bioavailability, low volume of distribution suggest that it is not likely to achieve an effective antiviral concentration of glycyrrhizin in lung tissues by oral or IV administration of safe doses 150-300 mg/day. An advanced delivery system such as inhalation can be considered to enhance its access to lung tissues.

### 3.2.4. Aescin(escin)

It's a blend of triterpene saponins extracted from the seeds of the horse chestnut tree; *Aesculus Hippocastanum* (*Sapindaceae*). It has a high molecular weight [94]. It possesses antiviral and immunomodulatory activities [95]. It has been widely used in the treatment of traumatic induced oedema, and chronic venous insufficiency [96]. It was demonstrated to inhibit acute inflammation similar to corticosteroids [45]. Its EC<sub>50</sub> against SARS-CoV was 6  $\mu$ M [45]. Given these pharmacological characteristics; it was suggested to consider escin as an add-on therapy in acute lung injury (ALI) associated with severe COVID-19 infection [96]. Injection formulation was registered in China and Italy for clinical trials to treat patients with COVID-19 pneumonia [77]. Aescin is not absorbed orally (bioavailability < 0.3%); likely due to extensive first-pass metabolism [97]. Aescin PK features

were studied in rats following IV injection at dosages ranging from 0.5 to 2.0 mg/kg. The half-life of Aescin was about 7-12 hr., Vd : 3-6 L/Kg [97]. PK data in humans not available.

### 3.2.5 Resveratrol

Resveratrol, a polyphenol compound (3,5,4'-trihydroxystilbene) found in grapes; *Vitis vinifera* (*Vitaceae*); Mulberry or *Morus nigra* (*Moraceae*) and Peanuts or *Arachis hypogaea* (*Fabaceae*). It possesses antioxidant, antitumor, antiviral and free radical scavenging properties [98]. Among the seven drugs tested against HCoV-229E *in vitro*, resveratrol demonstrated a favourable antiviral effect (EC<sub>50</sub>) = 4.6  $\mu$ M which was superior to two drugs namely lopinavir/ritonavir (EC<sub>50</sub> = 8.8  $\mu$ M), and Chloroquine (EC<sub>50</sub> = 5  $\mu$ M). Resveratrol has also the best selectivity index (SI) of 45.65 [70]. It showed the ability to attenuate lung injury in animal models [99]. In this context, studies documented its ability to reduce biomarkers of inflammation, tumor necrosis factor (TNF- $\alpha$ ), and C-reactive protein [100].

Resveratrol has poor oral bioavailability due to extensive metabolism in the gastrointestinal tract (GIT) and liver. It has a very short half-life, but its active sulfate metabolites showed a half-life of about 8 hr. It was speculated that levels of resveratrol in serum after oral administration are not likely to achieve an effective antiviral level [99]. However, the optimized pharmaceutical formulation may solve this limitation e.g., nanotechnology-based formulations and liposomes [72,101]. Clinical studies showed favorable safety outcomes of resveratrol. A good safety profile was assured with doses up to 500 mg/d for 60 days [102]. Most reported adverse effects are mild GIT symptoms such as diarrhea [103]. However, an advanced delivery system is needed to improve its PK profile.

### 3.2.6 Emodin

Emodin (6-methyl-1,3,8-trihydroxyanthraquinone) is found in many plants such as Chinese rhubarb; *Rheum palmatum* (*Polygonaceae*) [104]. It has many effective preventive and therapeutic effects [105]. Emodin was demonstrated to block the binding of SARS-CoV spike protein binding to its ACE2 receptors, hence it is likely to block viral entry into host cells [66]. It is one of the top 16 network-predicted repurposed drugs against SARS-CoV and is suggested among drug combinations (toremifene

plus emodin) [106]. It is included among traditional Chinese medicine against viral infections [107]. Its ability to attenuate lung injury is well documented in an animal model [108]. In animal studies, it has poor oral bioavailability due to extensive metabolism by glucuronidation. Emodin can cause reproductive, liver, and kidney toxicity, especially after chronic use [109].

### 3.2.7 Ginsenoside-Rb1 (G-Rb1)

G-Rb1 appears to be most abundant in *Panax quinquefolius* (Araliaceae) (American Ginseng). Ginsenosides possess a variety of potential health effects [110]. The ability of ginseng to attenuate experimentally induced lung injury are well documented and several mechanisms were suggested to explain its pharmacological effect [111]. Molecular docking analysis indicated that ginsenosides have the potential to inhibit the SARS-CoV-2 Mpro receptor [112]. Genesiondis have very low bioavailability after oral administration [113]. Several studies documented the feasibility of enhancing its bioavailability by adopting several pharmaceutical technology approaches [114, 115]. When ginseng extract was standardized at a concentration of 4 mg ginsenosides/100 mg capsule and given at a dose of up to 114 g ginsenoside/kg to humans for up to 12 weeks, no significant adverse effects were demonstrated [116].

### 3.2.8 Eugenol

It is a phenolic molecule (methoxy phenols, M Wt 164.2) found in several plants such as cinnamon *Cinnamomum* (Lauraceae), clove *Syzygium aromaticum* (Myrtaceae), Tulsi *Ocimum tenuiflorum* (Lamiaceae). The potential value of clove and eugenol against COVID-19 was suggested based on their antiviral, anti-inflammatory, and antithrombotic effects [117]. Docking analysis indicated the possible interaction of  $\alpha$ -pinene and eugenol with SARS-CoV spike protein [118]. The ability of eugenol to attenuate experimentally acute induced lung injury was documented [68]. It is likely mediated by the inhibition of lipoxygenase and cyclooxygenase pathways [119].

In mice, eugenol was readily absorbed after oral administration. The mean plasma half-life was about 14 hrs. suggesting a potential accumulation of the drug following repeated administrations [120]. A good safety profile was demonstrated in animal toxicity studies [121]. It is generally

recognized as safe by the food and drug administration (FDA) [122]. However, local hypersensitivity reaction was reported in dental practice [123]. Based on its pharmacology profile, it is strongly recommended for pre-clinical studies and clinical studies, optimized formulations to explore its utility in the management of respiratory viral infections.

### 3.2.9 Gingerol

Gingerol, (6-gingerol) is a phenolic compound found in fresh ginger, *Zingiber officinale* (Zingiberaceae). Pre-clinical *in vitro* and animal models documented a broad spectrum of therapeutic values [124]. The favorable impact of ginger and its active ingredients on the immune system are extensively studied, *in vitro* studies included; modulation of inflammatory cytokines levels [125], inhibition of lipopolysaccharide (LPS)-induced inflammatory responses in macrophages [126], the potential of the anti-inflammatory effect of some drugs such as paracetamol [127]. Other studies documented the therapeutic benefits in the management of allergic asthma and allergic rhinitis [128, 129]. Pharmacokinetic (PK) studies showed that, in general, is well absorbed orally and has good tissue distribution [130,131]. A good safety profile was reported in clinical trials [132], several pharmaceutical nanotechnology approaches were suggested to enhance its systemic availability or distribution [133, 134].

### 3.2.10 Eucalyptol (1,8-cineole)

It is a cyclic ether and monoterpenoid, it controls airway mucus hypersecretion and asthma via anti-inflammatory cytokine inhibition. It is abundant in many *Eucalyptus species* (Myrtaceae) [135]. It has low toxicity (LD<sub>50</sub> for rats 2.5 g/kg) [116]. It showed the ability to attenuate ALI in an animal model by different mechanisms [136,137]. A comprehensive review of the ability of essential oil to suppress several respiratory viruses was studied by Wani et al. [138]. Eucalyptus essential oils have been used to treat a variety of respiratory diseases, including sinusitis, pharyngitis, and bronchitis. One of its active ingredients (1,8-cineole), showed smooth muscle relaxant effects [139]. Furthermore, investigations have shown that inhaling cineole has analgesic and anti-inflammatory properties, suggesting that it could be utilized to treat chronic obstructive pulmonary disease (COPD) and asthma [140]. Sharma et al. [141] used molecular docking to show that

eucalyptol (1,8-cineole) may bind to Mpro and suppress viral multiplication. Strong ionic, hydrogen bonds and hydrophobic interactions have been described in the eucalyptol/Mpro complexes.

Several investigations found that eucalyptus oil and its active component, eucalyptol, have strong immunomodulatory activities. They were able to reduce the release of pro-inflammatory cytokines from macrophages and monocytes while maintaining their phagocytic properties [142].

In conclusion, it has an anti-inflammatory, bronchodilator, antibacterial effect, mucolytic, and antiviral activity. It is well absorbed orally, can be inhaled, and has good lipophilicity which suggests adequate access to lung cells. It has a good safety profile; therefore, the compound is suggested for more investigations in the management of respiratory viral infections.

### 3.2.11 Cinnamaldehyde

Cinnamaldehyde is an active ingredient in the bark of Cinnamon trees *Cinnamomum verum* (*Lauraceae*). The essential oil of cinnamon bark contains > 90% cinnamaldehyde (predominantly the Trans E- isomer) synthetic compound is also available. According to docking experiments, cinnamaldehyde may prevent SARS-CoV-2 from attaching to its target receptors [143]. Cinnamaldehyde's preventive benefits in the animal model of ALI have been demonstrated [144]. Different mechanisms were demonstrated; including neutrophils, macrophages, and total cell number in bronchoalveolar lavage fluid is inhibited. It reduced the levels of inflammatory cytokines such TNF-, interleukin (IL-6), IL-13, and IL-1 [145].

Cinnamaldehyde is approved by the FDA for use within allergenic percutaneous patch tests [146]. It has had a low potential for toxicity but weak sensitization reactions in animals and human skin allergy were reported [147]. It is metabolized to cinnamyl alcohol and methyl cinnamate and cinnamic acid, these compounds are potentially toxic [146]. In acute studies, these materials have a low to moderate order of oral toxicity (LD<sub>50</sub> values of 1.5–39 g/kg body weight) [148].

About 52% of cinnamaldehyde is absorbed through the skin and shown to be rapidly absorbed from the gut. It is metabolized and excreted as polar metabolites, the final major

urinary metabolite is hippuric acid [148]. Novel intravenous sub-micrometre emulsion showed good tissue distribution and enhanced antitumor efficacy without significant toxicity in an animal model of cancer [149].

### 3.2.12 Kaempferol (KMF)

Kaempferol (3,5,7-trihydroxy-2-(4-hydroxyphenyl)-4H-chromen-4-one) is a polyphenol abundant in fruits and vegetables, herbal medications, and beverages derived from plants. It's used to treat a variety of diseases [150]. Several studies suggested KMF has superior efficacy compared to many other natural compounds against SARS-CoV-2 [151,152]. KMF shares structural similarities with other flavonoids such as myricetin and dihydromyricetin, which showed strong inhibitory activity against the 3-chymotrypsin like protease(3CLpro) of SARS-CoV-2 [153]. KMF was shown to inhibit MERS 3CLpro (IC<sub>50</sub> = 35.3 μM) and SARS 3CLpro (IC<sub>50</sub> = 116.3 μM) [154]. Molecular docking documented strong interaction of KMF with SARs-CoV-2 Mpro 3CLpro [155]. This finding was confirmed by another docking study, which also documented an inhibition rate of about 90 % at a concentration of 62.5 μM cell protective effect (CPE) inhibition assay. The authors concluded that KMF was found to protect cells against virus-induced cell death, suggesting that it could be a promising SARS-CoV-2 antiviral treatment [151].

The ability of KMF to attenuate the experimentally induced lung injury are well documented, various mechanisms were suggested [156,157]. KMF is expected to show low bioavailability, therefore several studies suggested the utility of pharmaceutical approaches such as phospholipid complexes; solid dispersion, self-emulsifying formulation to enhance its bioavailability [158], Nano formulations were also investigated to improve PK characteristics of KMF [159]. On the other hand, KMF was found to enhance the bioavailability of some drugs after oral administration [160].

### 3.2.13 Rutin

Rutin, (a glycoside) also known as rutoside or quercetin-3-O-rutinoside, is a citrus flavonoid found in a range of plants, including Citrus (*Rutaceae*) and Buckwheat or Fagopyrum esculentum (*Polygonaceae*) species [161]. Several computations and *in vitro* testing suggest its activity against SARS-CoV-2 [162,163]. Based

on the crystal structures of 3CLpro and RdRp, docking studies revealed that sulfate or glucuronide metabolite of rutin can inhibit these enzymes; which are essential for replication of SARS-CoV-2 [164]. Moreover, it demonstrated a good ability to attenuate experimentally induced ALI [165], and potent thrombolytic activity [166].

Before being absorbed into the circulation, rutinoides like rutin and nicotiflorin are deglycosylated and subsequently conjugated mostly with glucuronate and sulfate, which are the predominant forms in plasma [167]. Intravenous or intranasal administration was suggested to improve the delivery of rutin [164]. Despite several preclinical mechanistic studies on its anticancer activities, the lack of well-designed randomized clinical trials on rutin's safety and therapeutic activity highlights the need for greater clinical research [168].

### 3.2.14 Leptodactylone

It is a coumarin isolated from *Leptodactylon pungens* (Polemoniaceae) and *Linanthus demissus* (Polemoniaceae) species, [169]. *In vitro* study was applied in Vero-E6 cells infected by SARS-CoV and measured the CPE by fluorescence microscope revealed that it produced 60 % protection of the infected cells (EC %) at 100 micro mols [170]. PK and safety data of Leptodactylone are not available.

### 3.2.15 Celastrol

It is a pentacyclic triterpenoid (belong to triterpene quinone methides) extracted from *Tripterygium wilfordii* ( Celastraceae) [171]. It has broad pharmacological activities [172]. Celastrol inhibits SARS-CoV-2 Mpro 3CLpro and acts as a superoxide radical scavenger, according to docking and other relevant studies [173]. Moreover, it showed promising *in vivo* results in animal models for inflammatory pulmonary diseases [174]. In rats, the pure compound showed poor bioavailability which improved after using an optimized tablet dosage form [175]. Several studies showed the utility of pharmaceutical techniques to enhance its PK characteristics [176,177]. However, liver and kidney induced toxicity was demonstrated in animal studies [178]. More studies are needed to clarify the safety concern.

### 3.2.16 Tetrandrine (TET)

It is a bis-benzylisoquinoline alkaloid calcium channel blocker. It is isolated from the plant

*Stephania tetrandra* (Menispermaceae). It has immunomodulatory effects and was found to be beneficial against inflammation, and lung cancer in clinical trials, with a favorable safety profile. It was demonstrated as effective against Ebola virus, Mycobacterium TB, Plasmodium falciparum, and Candida albicans. Tetrandrine's pharmacological properties have been demonstrated by its effects on many signaling pathways, including, calcium channel, suppression of formation of reactive oxygen species, increased autophagy, reversal of multidrug resistance, and caspase pathway [179]. It was suggested to act against the coronavirus by antagonism of the two-pore channel 2 (TPC2) [180]. It also decreased human coronavirus strain OC43 (HCoV-OC43) infection of MRC-5 human lung cells *in vitro*, with an inhibitory concentration (IC<sub>50</sub>) of 0.33 μM/L [40]. It showed promising effects in the management of silicosis, and in one study it was given by inhalation to ameliorate lung fibrosis [181].

A published hypothesis, discussed in detail that TET can be considered a promising treatment for COVID-19. The research reviewed the multiple properties of the drug, including its ability to inhibit replication of related coronavirus at very low concentrations. Then, the researcher also discussed its PK and the possibility of achieving an effective concentration in lung tissue. They also reviewed the expected side effects; drug interactions and precautions to be considered in certain situations [180].

### 3.2.17 Pristimerin (Pris)

Pristimerin, a quinonoid triterpene, isolated from the Chinese herbal plant *Tripterygium wilfordii* (Celastraceae), and it has been claimed that pristimerin has anti-fungal, anti-oxidant, anti-bacterial, and antiviral characteristics [182,183]. It is active against SARS-CoV with an IC<sub>50</sub> = 5.5 0.7 M According to several studies, pristimerin and its biological counterpart URB602 are monoglyceride lipase inhibitors, [184]. The formation of proteasomes was suppressed, NF-κB activity and cyclin D1 expression were downregulated, and Bax, caspase, and cleavage of poly (ADP-ribose) polymerase PARP-1 were activated as part of pristimerin's anticancer activities [185]. Studies in animals showed that Pris has a half-life of about 2 hours and poor bioavailability of about 28 % after oral administration. It is primarily metabolized by phase I enzymes [186].

**Table 3. Bioavailability of specified phytochemicals**

<b>Phytochemicals</b>	<b>Oral route</b>	<b>Intravenously (IV)/ Intraperitoneal Route (IP)</b>	<b>Enhancer and notes</b>
<b>Glycyrrhizin [ 209]</b>	Extremely low		Sodium caprate > sodium laurate > sodium caprylate> sodium oleate Nasal-80-fold greater compared with oral administration. Rectal administration, greater compared with oral
<b>Resveratrol [ 210, 211]</b>	Preferred route		Dose-dependent. Low water solubility which explains its slow absorption,
<b>Lycorine [88]</b>	Not available	No significant difference between (IP, iv)- The concentration in plasma was undetectable, which indicate its rapid clearance	
<b>Emodin [109, 212]</b>	Poor intestinal absorption, fast elimination, and low bioavailability.		Gender-dependent
<b>Aescin [213]</b>	Film-coated tablet with sustained- release allowed high bioavailability		
<b>G-RB1 [ 113]</b>	The low oral bioavailability of RB1 and rapid reduction of Rg1 in blood indicated that pharmaceutical technology is necessary		
<b>CELASTROL [175]</b>	Tablets with enhanced bioavailability was reported		Gender-dependent

### 3.2.18 Curcumin

It is described as a drug with multitarget for multiple chronic diseases. It has broad pharmacological activity. Its potential role in the management of COVID-19 was suggested [187, 188]. Curcumin is a bright yellow chemical produced by plants of the *Curcuma longa* (Zingiberaceae) that could potentially block ACE2 to prevent COVID-19 entrance into host cells [189]. It also has a potential inhibitory effect on COVID-19 Mpro [190]. An *in vitro* study showed that treatment with cationic carbon dots based on curcumin can suppress coronavirus replication by stimulating the production of interferon-stimulating genes (ISGs) and cytokines (IL8 and IL6) of Vero cells by triggering the innate immunity of the host [191]. Curcumin has been shown to lower influenza A virus infection and its associated pneumonia. It suppresses virus-induced activation of several inflammatory pathways (TLR2/4, MAPK, and NFB). Given these findings, it was suggested that curcumin has a potential role in the treatment of COVID-19 induced lung complications [192]. Curcumin has anti-inflammatory and anti-fibrotic effects by several mechanisms [193]. Furthermore, in experimental models of lung fibrosis, curcumin has been demonstrated to decrease collagen formation [194]. A study showed that prophylactic application of curcumin decreased the inflammation and reduces the influx of fluid in the lungs of a rat model of induced hypoxia [195].

Curcumin has poor bioavailability and as such, it is not likely to show good tissue distribution. [196]. Some clinical trials suggested its good safety profile after oral administration [197]. However, it is considered unsafe for pregnant women. It can cause GIT side effects, and there are some reports of liver injury [198, 199].

### 3.2.19 Betulinic acid (BTA)

BTA is lupine-type pentacyclic triterpenoid saponin (3 $\beta$ -hydroxy-lup-20). It is found in the bark of several natural plants, primarily *Betula pendula* (Betulaceae). Its anti-inflammatory, anti-angiogenic, and immunomodulatory properties, as well as its anti-human immunodeficiency virus effects, have been demonstrated. The majority of studies focused on its anticancer effects and postulated the underlying mechanisms [200, 201]. BTA was discovered to be a strong inhibitor of (SARS-CoV) *in vitro* [202]. Molecular modelling analysis indicated that BTA interacts

with coronavirus (SARS-CoV) 3CL protease [203]. Several studies suggested BTA ability to attribute experimentally induced lung injury by several mechanisms [204, 205]. BTA has very low water solubility that explains its low bioavailability. Some derivatives were suggested to improve its bioavailability while retaining its antitumor activity [206]. Other studies utilized pharmaceutical techniques such as liposomes, nanosuspension, nanoencapsulation analogue [5, 207] to improve its bioavailability. In animal experiments, a good safety profile was documented [208].

## 4. CONCLUSION

- 1) Many phytochemicals have been suggested to have potent anti-SARS-CoV-2 activity based on *in vitro* testing. However, preclinical research is sparse, and only a few compounds were tested in clinical trials against COVID-19.
- 2) Extensive research is needed to provide a full pharmacological profile of phytochemicals that showed promising efficacy against SARS-CoV-2.
- 3) The pharmacokinetics (PK) features of phytochemicals could be improved with pharmaceutical technologies. Table 3 summarizes the bioavailability of several substances as well as techniques to improve them.
- 4) Decision-makers, researchers, and pharmaceutical companies will benefit from this study.

## CONSENT AND ETHICAL APPROVAL

This systematic review is approved by Ibn Sina National College for Medical Studies- Research Center (ISNC-RC) number 011CPP23112021.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Roser M, et al. Coronavirus pandemic (COVID-19), in Our world in data. Our world in data; 2021.
2. Singh SP, et al. Microstructure, pathophysiology, and potential therapeutics of COVID-19: A comprehensive review. J Med Virol. 2021;93(1):275-299.

3. Yang X, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: A single-centered, retrospective, observational study. *The Lancet Respiratory Medicine*. 2020;8(5):475-481.
4. Chakraborty I, Maity P. COVID-19 outbreak: Migration, effects on society, global environment and prevention. *Science of the Total Environment*. 2020; 728:138882.
5. Wang R, et al. Decoding SARS-CoV-2 transmission and evolution and ramifications for COVID-19 diagnosis, vaccine, and medicine. *Journal of chemical information and modeling*. 2020;60(12):5853-5865.
6. Ang L, et al. Herbal medicine for the treatment of coronavirus disease 2019 (COVID-19): A systematic review and meta-analysis of randomized controlled trials. *J Clin Med*. 2020;9(5):1583.
7. Arentz S, et al. Honeybee products for the treatment and recovery from viral respiratory infections including SARS-COV-2: A rapid systematic review. *Integr Med Res*. 2021;10(Suppl):100779.
8. Aschale Y, et al. A systematic review on traditional medicinal plants used for the treatment of viral and fungal infections in Ethiopia. *J Exp Pharmacol*. 2021;13:807-815.
9. Aucoin M, et al. A systematic review on the effects of Echinacea supplementation on cytokine levels: Is there a role in COVID-19? *Metabol Open*. 2021;11:100115.
10. Du X, et al. Add-on effect of Chinese herbal medicine in the treatment of mild to moderate COVID-19: A systematic review and meta-analysis. *PLoS One*. 2021;16(8): e0256429.
11. Du XQ, et al. Add-on effect of honeysuckle in the treatment of coronavirus disease 2019: A systematic review and meta-analysis. *Front Pharmacol*. 2021;12: 708636.
12. Fan AY, Gu S, Alemi SF. Chinese herbal medicine for COVID-19: Current evidence with systematic review and meta-analysis. *J Integr Med*. 2020;18(5):385-394.
13. Jiang F, et al. Contribution of traditional Chinese medicine combined with conventional western medicine treatment for the novel coronavirus disease (COVID-19), current evidence with systematic review and meta-analysis. *Phytother Res*. 2021;1-18.
14. Liang SB, et al. Therapeutic effects and safety of oral Chinese patent medicine for COVID-19: A rapid systematic review and meta-analysis of randomized controlled trials. *Complement Ther Med*. 2021;60: 102744.
15. Luo X, et al. The add-on effect of Chinese herbal medicine on COVID-19: A systematic review and meta-analysis. *Phytomedicine*. 2021;85:153282.
16. Oladele JO, et al. A systematic review on COVID-19 pandemic with special emphasis on curative potentials of Nigeria based medicinal plants. *Heliyon*. 2020; 6(9):e04897.
17. Pang W et al. Chinese medical drugs for coronavirus disease 2019: A systematic review and meta-analysis. *Integr Med Res*, 2020;9(3):100477.
18. Shi S, et al. The effect of Chinese herbal medicine on digestive system and liver functions should not be neglected in COVID-19: An updated systematic review and meta-analysis. *IUBMB Life*. 2021; 73(5):739-760.
19. Singh RS, et al. Promising traditional Indian medicinal plants for the management of novel Coronavirus disease: A systematic review. *Phytother Res*. 2021;35(8):4456-4484.
20. Sun CY, Sun YL, Li XM. The role of Chinese medicine in COVID-19 pneumonia: A systematic review and meta-analysis. *Am J Emerg Med*. 2020;38(10): 2153-2159.
21. Umeogaju FU et al., Plant-derived food grade substances (PDFGS) active against respiratory viruses: A systematic review of non-clinical studies. *Front Nutr*. 2021;8: 606782.
22. Wang Y, Greenhalgh T, Wardle J. Chinese herbal medicine ("3 medicines and 3 formulations") for COVID-19: Rapid systematic review and meta-analysis. *J Eval Clin Pract*. 2021;1-20.
23. Wu X, et al. Traditional chinese medicine as an adjunctive therapy for mild and common COVID-19: A systematic review and network meta-analysis. *Medicine (Baltimore)*. 2021;100(40): e27372.
24. Wu XV, et al. Traditional Chinese medicine as a complementary therapy in combat with COVID-19-A review of evidence-based research and clinical practice. *J Adv Nurs*. 2021;77(4):1635-1644.

25. Xiong X et al. Chinese herbal medicine for coronavirus disease 2019: A systematic review and meta-analysis. *Pharmacol Res*, 2020;160:105056.
26. Yan LZ, et al. Clinical effects of the combination of Traditional Chinese and Western Medicines on coronavirus disease 2019: a systematic review and Meta-analysis. *J Tradit Chin Med*. 2021;41(4): 499-506.
27. Yin B, et al. Efficacy of integrated traditional chinese and western medicine for treating COVID-19: A Systematic Review and Meta-Analysis of RCTs. *Front Public Health*. 2021;9:622707.
28. Zeng M, Li L, Wu Z. Traditional Chinese medicine Lianhua Qingwen treating corona virus disease 2019(COVID-19): Meta-analysis of randomized controlled trials. *PLoS One*. 2020;15(9): e0238828.
29. Zhou LP, et al. The effects of traditional chinese medicine as an auxiliary treatment for COVID-19: A systematic review and meta-analysis. *J Altern Complement Med*. 2021;27(3):225-237.
30. Zhuang J, et al. A meta-analysis for Lianhua Qingwen on the treatment of Coronavirus disease 2019 (COVID-19). *Complement Ther Med*. 2021;60:102754.
31. Johnson N, Phillips M. Rayyan for systematic reviews. *Journal of Electronic Resources Librarianship*. 2018;30(1):46-48.
32. Page MJ, et al. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *Bmj*. 2021; 372.
33. Monticolo F, et al. anti-HCoV: A web resource to collect natural compounds against human coronaviruses. *Trends in Food Science & Technology*. 2020;106:1-11.
34. Paraiso IL, Revel JS, Stevens JF. Potential use of polyphenols in the battle against COVID-19. *Curr Opin Food Sci*. 2020;32:149-155.
35. Annunziata G, et al. May polyphenols have a role against coronavirus infection? An Overview of in vitro Evidence. *Front Med (Lausanne)*. 2020;7:240.
36. El-Missiry, M.A., et al., Polyphenols are potential nutritional adjuvants for targeting COVID-19. *Phytother Res*, 2020, p. 2879-2889.
37. Solnier J, Fladerer JP. Flavonoids: A complementary approach to conventional therapy of COVID-19? *Phytochem Rev*. 2020;1-23.
38. Russo M, et al. Roles of flavonoids against coronavirus infection. *Chem Biol Interact*. 2020;328:109211.
39. Keum YS, Jeong YJ. Development of chemical inhibitors of the SARS coronavirus: viral helicase as a potential target. *Biochemical Pharmacology*. 2012; 84(10):1351-1358.
40. Kim DE, et al. Natural bis-benzylisoquinoline alkaloids-tetrandrine, fangchinoline, and cepharanthine, inhibit human coronavirus OC43 infection of MRC-5 human lung cells. *Biomolecules*. 2019;9(11):696.
41. Wink M, Potential of DNA intercalating alkaloids and other plant secondary metabolites against SARS-CoV-2 causing COVID-19. *Diversity*. 2020;12(5):175.
42. Li Sy et al. Identification of natural compounds with antiviral activities against SARS-associated coronavirus. *Antiviral Research*. 2005;67(1):18-23.
43. Jin YH et al. Lycorine, a non-nucleoside RNA dependent RNA polymerase inhibitor, as potential treatment for emerging coronavirus infections. *Phytomedicine*. 2020;153440.
44. Kim DE et al. Natural bis-benzylisoquinoline alkaloids-tetrandrine, fangchinoline, and cepharanthine, inhibit human coronavirus OC43 infection of MRC-5 human lung cells. LID - 10.3390/biom9110696 [doi] LID-696. 2218-273X (Electronic).
45. Wu CY, et al. Small molecules targeting severe acute respiratory syndrome human coronavirus. *Proceedings of the National Academy of Sciences*. 2004;101(27): 10012-10017.
46. Hisham Shady N et al. Sterols and Triterpenes: Antiviral Potential Supported by In-Silico Analysis. *Plants (Basel)*. 2020; 10(1).
47. Wen CC et al. Specific plant terpenoids and lignoids possess potent antiviral activities against severe acute respiratory syndrome coronavirus. *Journal of Medicinal Chemistry*. 2007;50(17):4087-4095.
48. Chang F, et al., Anti-human coronavirus (anti-HCoV) triterpenoids from the leaves

- of *Euphorbia nerifolia*. *Nat Prod Commun*. 2012;7:1415–1417.
49. Lin SC et al. Effective inhibition of MERS-CoV infection by resveratrol. *BMC infectious diseases*. 2017;17(1):1-10.
  50. Hoever G et al. Antiviral activity of glycyrrhizic acid derivatives against SARS– coronavirus. *Journal of Medicinal Chemistry*. 2005;48(4):1256-1259.
  51. Chen CN et al. Inhibition of SARS-CoV 3C-like protease activity by theaflavin-3, 3'-digallate (TF3). *Evidence-Based Complementary and Alternative Medicine*. 2005; 2(2):209-215.
  52. Boukhatem MN, Setzer WN. Aromatic herbs, medicinal plant-derived essential oils, and phytochemical extracts as potential therapies for coronaviruses: Future Perspectives. *Plants (Basel)*. 2020; 9(6):1-23.
  53. Senthil Kumar KJ et al. Geranium and lemon essential oils and their active compounds downregulate angiotensin-converting enzyme 2 (ACE2), a SARS-CoV-2 Spike Receptor-Binding Domain, in Epithelial Cells. *Plants (Basel)*. 2020;9(6): 1-12.
  54. Wilkin PJ, et al. The undiscovered potential of essential oils for treating SARS-CoV-2 (COVID-19). *Curr Pharm Des*. 2020; 26(41):5261-5277.
  55. Boozari M, Hosseinzadeh H. Natural products for COVID-19 prevention and treatment regarding to previous coronavirus infections and novel studies. *Phyther Res*. 2021;35(2):864-876.
  56. Islam MT, et al. Natural products and their derivatives against coronavirus: A review of the non-clinical and pre-clinical data. *Phyther Res*. 2020;34(10):2471-2492.
  57. Khalifa SAM, et al. Screening for natural and derived bio-active compounds in preclinical and clinical studies: One of the frontlines of fighting the coronaviruses pandemic. *Phytomedicine*. 2020;153311.
  58. Zahedipour F et al. Potential effects of curcumin in the treatment of COVID-19 infection. 1099-1573 (Electronic).
  59. Zhang D et al. Valinomycin as a potential antiviral agent against coronaviruses: A review. *Biomed J*. 2020;43(5):414-423.
  60. Park JY et al. Tanshinones as selective and slow-binding inhibitors for SARS-CoV cysteine proteases. *Bioorganic & Medicinal Chemistry*. 2012;20(19):5928-5935.
  61. Giordo R, et al. Therapeutic potential of resveratrol in COVID-19-Associated Hemostatic Disorders. *Molecules*. 2021; 26(4):1-10.
  62. Cinatl J, et al. Glycyrrhizin, an active component of liquorice roots, and replication of SARS-associated corona-virus. *The Lancet*. 2003;361(9374):2045-2046.
  63. Islam MN, et al. Revisiting pharmacological potentials of *Nigella sativa* seed: A promising option for COVID-19 prevention and cure. LID - 10.1002/ptr.6895 [doi].p. 1099-1573 (Electronic).
  64. Orhan IE, Senol Deniz FS. Natural products as potential leads against coronaviruses: Could they be encouraging structural models against SARS-CoV-2? *Natural Products and Bioprospecting*, 2020;10(4):171-186.
  65. Fan HH, et al. Repurposing of clinically approved drugs for treatment of coronavirus disease 2019 in a 2019-novel coronavirus-related coronavirus model. *Chinese Medical Journal*. 2020;133(9): 1051–1056.
  66. Ho TY et al. Emodin blocks the SARS coronavirus spike protein and angiotensin-converting enzyme 2 interaction. *Antiviral Research*. 2007;74(2):92-101.
  67. Lung J, et al. The potential chemical structure of anti-SARS-CoV-2 RNA-dependent RNA polymerase. *Journal of medical virology*. 2020;92(6):693-697.
  68. Zhang ZD et al. Aspirin eugenol ester ameliorates paraquat-induced oxidative damage through ROS/p38-MAPK-mediated mitochondrial apoptosis pathway. *Toxicology*, 2021;453: 152721.
  69. Zhang YN, et al. Gemcitabine, lycorine and oxysphoridine inhibit novel coronavirus (SARS-CoV-2) in cell culture. *Emerging Microbes & Infections*. 2020;9(1):1170-1173.
  70. Pasquereau S, et al. Resveratrol Inhibits HCoV-229E and SARS-CoV-2 Coronavirus Replication In Vitro. *Viruses*. 2021;13(2):1-11.
  71. Barboza JN, et al. An overview on the anti-inflammatory potential and antioxidant profile of eugenol. *Oxid Med Cell Longev*, 2018;2018:3957262.
  72. Huang M, et al. Liposome co-encapsulation as a strategy for the delivery

- of curcumin and resveratrol. *Food Funct*. 2019;10(10):6447-6458.
73. Silva J, et al. Essential oils as antiviral agents. potential of essential oils to treat SARS-CoV-2 infection: An in-silico investigation. *Int J Mol Sci*. 2020;21(10):1-35.
  74. Thuy BTP, et al. Investigation into SARS-CoV-2 Resistance of Compounds in Garlic Essential Oil. 2020;2470-1343. (Electronic data base).
  75. de Ligt M et al. Resveratrol supplementation reduces ACE2 expression in human adipose tissue. *Adipocyte*. 2021;10(1):408-411.
  76. Valizadeh H, et al. Nano-curcumin therapy, a promising method in modulating inflammatory cytokines in COVID-19 patients. *International immunopharmacology*. 2020;89:107088.
  77. López-Alcalde J, et al. Current state of research about Chinese herbal medicines (CHM) for the treatment of coronavirus disease 2019 (COVID-19): A scoping review. *The Journal of Alternative and Complementary Medicine*. 2020;26(7):557-570.
  78. Imran M et al. Thymoquinone: A novel strategy to combat cancer: A review. *Biomedicine & Pharmacotherapy*. 2018; 106:390-402.
  79. Al-Gabri NA, et al. Thymoquinone ameliorates pulmonary vascular damage induced by *Escherichia coli*-derived lipopolysaccharide via cytokine downregulation in rats. *Environ Sci Pollut Res Int*. 2019;26(18):18465-18469.
  80. Çolak M, et al. Efficacy of thymoquinone in the treatment of experimental lipopolysaccharide-induced acute lung injury. *Kardiochir Torakochirurgia Pol*. 2020;17(2): 65-69.
  81. Isik AF, et al. A new agent for treatment of acute respiratory distress syndrome: thymoquinone. An experimental study in a rat model. *Eur J Cardiothorac Surg*. 2005; 28(2):301-5.
  82. Sommer AP, Försterling HD, Naber KG. Thymoquinone: Shield and sword against SARS-CoV-2. *Precis Nanomed*. 2020;3(3): 541-548.
  83. Pathan SA et al. Stability-indicating ultra-performance liquid chromatography method for the estimation of thymoquinone and its application in biopharmaceutical studies. *Biomedical Chromatography*. 2011;25(5):613-620.
  84. Goyal SN et al. Therapeutic potential and pharmaceutical development of thymoquinone: A multitargeted molecule of natural origin. *Front Pharmacol*. 2017;8: 656.
  85. Ansar FHZ et al. Pharmacokinetics and Biodistribution of Thymoquinone-loaded Nanostructured Lipid Carrier After Oral and Intravenous Administration into Rats. *International Journal of Nanomedicine*. 2020;15:7703.
  86. Rathore C et al. Nanocarriers: More than tour de force for thymoquinone. *Expert Opinion on Drug Delivery*. 2020;17(4):479-494.
  87. Cao Z, Yang P, Zhou Q. Multiple biological functions and pharmacological effects of lycorine. *Science China Chemistry*. 2013; 56(10):1382-1391.
  88. Ren L, Zhao H, Chen Z. Study on pharmacokinetic and tissue distribution of lycorine in mice plasma and tissues by liquid chromatography–mass spectrometry. *Talanta*. 2014;119:401-406.
  89. Roy M et al. Lycorine: A prospective natural lead for anticancer drug discovery. *Biomedicine & Pharmacotherapy*. 2018; 107:615-624.
  90. Pastorino G et al. Liquorice (*Glycyrrhiza glabra*): A phytochemical and pharmacological review. *Phytother Res*. 2018. 32(12):2323-2339.
  91. Chen F et al. *In vitro* susceptibility of 10 clinical isolates of SARS coronavirus to selected antiviral compounds. *Journal of Clinical Virology*. 2004;31(1):69-75.
  92. van de Sand L et al. Glycyrrhizin effectively neutralizes SARS-CoV-2 *in vitro* by inhibiting the viral main protease. *BioRxiv*. 2020;1-10.
  93. Yang R et al. The anti-inflammatory activity of licorice, a widely used Chinese herb. *Pharmaceutical Biology*. 2017;55(1):5-18.
  94. Yoshikawa M, et al. Bioactive saponins and glycosides. XII. Horse chestnut.(2): Structures of escins IIIb, IV, V, and VI and isoescins Ia, Ib, and V, acylated polyhydroxyoleanene triterpene oligoglycosides, from the seeds of horse chestnut tree (*Aesculus hippocastanum* L., Hippocastanaceae). *Chemical and Pharmaceutical Bulletin*. 1998;46(11): 1764-1769.

95. Gallelli L et al. Severe acute lung injury related to COVID-19 infection: A review and the possible role for escin. *Journal of Clinical Pharmacology*. 2020;60(7):815-825.
96. Gallelli L. Escin: A review of its anti-edematous, anti-inflammatory, and venotonic properties. *Drug Design, Development and Therapy*. 2019;13:3425.
97. Wu XJ et al. Comparative pharmacokinetics and bioavailability of escin Ia and isoescin Ia after administration of escin and of pure escin Ia and isoescin Ia in rat. *Journal of Ethnopharmacology*. 2012. 139(1):201-206.
98. Marinella MA. Metronomic phytonutrition: How daily, regular intake of plant-based foods may decrease cancer risk. 2017: Beacon Publishing and Design; 2017.
99. Baur JA, Sinclair DA. Therapeutic potential of resveratrol: the in vivo evidence. *Nature Reviews Drug Discovery*, 2006;5(6):493-506.
100. Koushki M, Dashatan NA, Meshkani R. Effect of resveratrol supplementation on inflammatory markers: A systematic review and meta-analysis of randomized controlled trials. (1879-114X (Electronic)).
101. Lin YC et al. Electrospun resveratrol-loaded polyvinylpyrrolidone/cyclodextrin nanofibers and their biomedical applications. *Pharmaceutics*. 2020;12(6):1-16.
102. Movahed A, et al. Efficacy and safety of resveratrol in type 1 diabetes patients: A two-month preliminary exploratory trial. *Nutrients*. 2020;12(1):161.
103. Cottart CH, Nivet-Antoine JL, V Fau -Beaudeux, Beaudeux JL. Review of recent data on the metabolism, biological effects, and toxicity of resveratrol in humans. (1613-4133 (Electronic)).
104. Agrawal DC et al. Medicinal plants and fungi: recent advances in research and development. Springer. (Electronic); 2017.
105. Cui Y et al. The pharmacology, toxicology and therapeutic potential of anthraquinone derivative emodin. (1875-5364 (Electronic)).
106. Zhou Y et al. Network-based drug repurposing for novel coronavirus 2019-nCoV/SARS-CoV-2. *Cell Discovery*. 2020; 6:14-14.
107. Runfeng L et al. Lianhuaqingwen exerts anti-viral and anti-inflammatory activity against novel coronavirus (SARS-CoV-2). *Pharmacological Research*, 2020;156: 104761.
108. Dong Y, et al. Emodin reactivated autophagy and alleviated inflammatory lung injury in mice with lethal endotoxemia. *Exp Anim*. 2019;68(4):559-568.
109. Dong X et al., Emodin: A review of its pharmacology, toxicity and pharmacokinetics. *Phytotherapy Research*. 2016; 30(8):1207-1218.
110. Popovich DG, Yeo CR, Zhang W. Ginsenosides derived from Asian (*Panax ginseng*), American ginseng (*Panax quinquefolius*) and potential cytoactivity. *International Journal of Biomedical and Pharmaceutical Sciences*. 2012;6(1):56-62.
111. Jiang Y et al. Ginsenoside Rb1 treatment attenuates pulmonary inflammatory cytokine release and tissue injury following intestinal ischemia reperfusion injury in mice. *Oxid Med Cell Longev*. 2015;2015: 843721.
112. Garg S et al. Molecular docking analysis of selected phytochemicals against SARS-CoV-2 M(pro) receptor. (0970-4078).
113. Tan ZY et al. [Pharmacokinetics and bioavailability of ginsenoside Rg1 in rats]. *Zhong Yao Cai*. 2013;36(7):1121-3.
114. Zeng M et al. Systematic review of recent advances in pharmacokinetics of four classical Chinese medicines used for the treatment of cerebrovascular disease. *Fitoterapia*. 2013;88:50-75.
115. Tam DNH et al. Ginsenoside Rh1: A Systematic Review of Its Pharmacological Properties. *Planta Med*. 2018;84(3):139-152.
116. Carabin IG, Burdock GA, Chatzidakis C. Safety assessment of *Panax ginseng*. *International Journal of Toxicology*. 2000; 19(4):293-301.
117. Vicidomini C, Roviello V, Roviello GN. Molecular basis of the therapeutical potential of clove (*Syzygium aromaticum* L.) and clues to its anti-COVID-19 utility. *Molecules*. 2021;26(7):1-12.
118. Saxena S et al. 'BhAVI-23'-A spice-herb based dietary infusion possessing in-vitro anti-viral potential. *Journal of Ayurveda and Integrative Medicine*. 2021;312-319.

119. Das Chagas Pereira de Andrade, F. and A.N. Mendes, Computational analysis of eugenol inhibitory activity in lipoxygenase and cyclooxygenase pathways. *Sci Rep.* 2020;10(1):16204.
120. Guénette SA et al. Pharmacokinetics of eugenol and its effects on thermal hypersensitivity in rats. *Eur J Pharmacol.* 2007;562(1-2):60-7.
121. Dhara L, Tripathi A. Sub-acute toxicological and behavioural effects of two candidate therapeutics, cinnamaldehyde and eugenol, for treatment of ESBL producing-quinolone resistant pathogenic Enterobacteriaceae. *Clin Exp Pharmacol Physiol.* 2020;47(6):977-988.
122. Meeting JFWCoFA. W.H. Organization, Evaluation of certain food additives and contaminants: Seventy-seventh report of the joint FAO/WHO expert committee on food additives. World Health Organization. 2013;77.
123. Sarrami N et al. Adverse reactions associated with the use of eugenol in dentistry. *Br Dent J.* 2002;193(5):257-9.
124. Butt MS, Sultan MT. Ginger and its health claims: Molecular aspects. *Critical reviews in Food Science and Nutrition.* 2011;51(5): 383-393.
125. Akinyemi AJ et al. Effect of ginger and turmeric rhizomes on inflammatory cytokines levels and enzyme activities of cholinergic and purinergic systems in hypertensive rats. *Planta Med.* 2016;82(7): 612-20.
126. Villalvilla A et al. 6-Shogaol inhibits chondrocytes' innate immune responses and cathepsin-K activity. *Mol Nutr Food Res.* 2014;58(2):256-66.
127. Montserrat-de la Paz S et al. Ginger rhizome enhances the anti-inflammatory and anti-nociceptive effects of paracetamol in an experimental mouse model of fibromyalgia. *Inflammopharmacology.* 2018;26(4):1093-1101.
128. Kardan M et al. Effect of ginger extract on expression of GATA3, T-bet and ROR- $\gamma$ t in peripheral blood mononuclear cells of patients with Allergic Asthma. *Allergol Immunopathol (Madr).* 2019;47(4):378-385.
129. Yocum GT et al. Ginger and its bioactive component 6-shogaol mitigate lung inflammation in a murine asthma model. *Am J Physiol Lung Cell Mol Physiol.* 2020; 318(2):L296-l303.
130. Li LL et al. Pharmacokinetics and tissue distribution of gingerols and shogaols from ginger (*Zingiber officinale* rosc.) in rats by UPLC–Q-Exactive–HRMS. *Molecules.* 2019;24(3):512.
131. Simon A et al. Blood-brain barrier permeability study of ginger constituents. *Journal of Pharmaceutical and Biomedical Analysis.* 2020;177:112820.
132. Wang S et al. Biological properties of 6-gingerol: A brief review. *Natural product Communications.* 2014;9(7): 1934578X1400900736.
133. Wang Q et al. A novel formulation of [6]-gingerol: Proliposomes with enhanced oral bioavailability and antitumor effect. *International Journal of Pharmaceutics.* 2018;535(1-2):308-315.
134. Wei Q et al. Formulation, characterization, and pharmacokinetic studies of 6-gingerol-loaded nanostructured lipid carriers. *AAPS PharmSciTech.* 2018;19(8):3661-3669.
135. Gilles M et al. Chemical composition and antimicrobial properties of essential oils of three Australian Eucalyptus species. *Food Chemistry.* 2010;119(2):731-737.
136. Gondim FL, Serra DS, Cavalcante F. Effects of eucalyptol in respiratory system mechanics on acute lung injury after exposure to short-term cigarette smoke. *Respir Physiol Neurobiol.* 2019;266:33-38.
137. Reis R et al. Modulation of cigarette smoke extract-induced human bronchial epithelial damage by eucalyptol and curcumin. *Hum Exp Toxicol.* 2021;960327121997986.
138. Wani AR et al. An updated and comprehensive review of the antiviral potential of essential oils and their chemical constituents with special focus on their mechanism of action against various influenza and coronaviruses. *Microb Pathog.* 2021;152:104620.
139. Bastos VP et al. Inhibitory effect of 1, 8-cineole on guinea-pig airway challenged with ovalbumin involves a preferential action on electromechanical coupling. *Clinical and Experimental Pharmacology and Physiology.* 2009;36(11):1120-1126.
140. Juergens LJ, Worth H, Juergens UR. New perspectives for mucolytic, anti-inflammatory and adjunctive therapy with 1, 8-cineole in COPD and asthma: Review on the new therapeutic approach. *Advances in Therapy.* 2020;37(5):1737-1753.

141. Sharma AD, Kaur I. Molecular docking studies on Jensenone from eucalyptus essential oil as a potential inhibitor of COVID 19 corona virus infection. arXiv preprint arXiv:2004.00217, Research & Reviews in Biotechnology & Biosciences. 2020;7(1):59-66.
142. Sadlon AE, Lamson DW. Immune-modifying and antimicrobial effects of Eucalyptus oil and simple inhalation devices. *Alternative Medicine Review*. 2010;15(1):33-43.
143. Kulkarni SA et al. Computational evaluation of major components from plant essential oils as potent inhibitors of SARS-CoV-2 spike protein. *Journal of Molecular Structure*. 2020;1221:128823.
144. Almoiliqy M et al. Protective effects of cinnamaldehyde against mesenteric ischemia-reperfusion-induced lung and liver injuries in rats. *Oxid Med Cell Longev*. 2020;2020:4196548.
145. Huang H, Wang Y. The protective effect of cinnamaldehyde on lipopolysaccharide induced acute lung injury in mice. *Cellular and Molecular Biology*. 2017;63(8):58-63.
146. Zhu R et al. Cinnamaldehyde in diabetes: A review of pharmacology, pharmacokinetics and safety. *Pharmacological Research*. 2017;122:78-89.
147. Isaac-Renton M, Li MK, Parsons LM. Cinnamon spice and everything not nice: many features of intraoral allergy to cinnamic aldehyde. *Dermatitis*. 2015; 26(3):116-121.
148. Panel TRE et al. A toxicologic and dermatologic assessment of related esters and alcohols of cinnamic acid and cinnamyl alcohol when used as fragrance ingredients. *Food and Chemical Toxicology*. 2007;45(1):S1-S23.
149. Zhao H et al. Cinnamaldehyde in a novel intravenous submicrometer emulsion: pharmacokinetics, tissue distribution, antitumor efficacy, and toxicity. *Journal of Agricultural and Food Chemistry*. 2015; 63(28):6386-6392.
150. Alam W et al. Kaempferol as a dietary anti-inflammatory agent: Current therapeutic standing. *Molecules*. 2020;25(18):4073.
151. Khan A et al. *In silico* and *in vitro* evaluation of kaempferol as a potential inhibitor of the SARS-CoV-2 main protease (3CLpro). *Phytotherapy research: PTR*. 2021;10.1002/ptr.6998.
152. Mehmood A et al. *In Silico* analysis of quranic and prophetic medicinal plants for the treatment of infectious viral diseases including corona virus. *Saudi J Biol Sci*. 2021;3137–3151.
153. Dai W et al. Structure-based design, synthesis and biological evaluation of peptidomimetic aldehydes as a novel series of antiviral drug candidates targeting the SARS-CoV-2 main protease. *BioRxiv*. 2020;1331-1335.
154. Jo S et al. Inhibition of SARS-CoV 3CL protease by flavonoids. *Journal of enzyme Inhibition and Medicinal Chemistry*. 2020. 35(1):145-151.
155. Khan A et al. Combined drug repurposing and virtual screening strategies with molecular dynamics simulation identified potent inhibitors for SARS-CoV-2 main protease (3CLpro). *Journal of Biomolecular Structure and Dynamics*. 2020;1-12.
156. Qian J, et al. Kaempferol reduces K63-linked polyubiquitination to inhibit nuclear factor-κB and inflammatory responses in acute lung injury in mice. *Toxicol Lett*. 2019;306:53-60.
157. Yang M et al. Resveratrol inhibits the replication of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in cultured Vero cells. *Phytother Res*. 2021; 35(3):1127-1129.
158. Chen ZP et al. Comparative pharmacokinetics and bioavailability studies of quercetin, kaempferol and isorhamnetin after oral administration of Ginkgo biloba extracts, Ginkgo biloba extract phospholipid complexes and Ginkgo biloba extract solid dispersions in rats. *Fitoterapia*. 2010;81(8):1045-1052.
159. Du Q et al. Comparison of different aliphatic acid grafted N-trimethyl chitosan surface-modified nanostructured lipid carriers for improved oral kaempferol delivery. *International Journal of Pharmaceutics*. 2019;568:118506.
160. Jackson LA et al. An mRNA vaccine against SARS-CoV-2-preliminary report. *New England Journal of Medicine*. 2020;1-8.
161. Jiang P et al. Rutin and flavonoid contents in three buckwheat species *Fagopyrum esculentum*, *F. tataricum*, and *F. homotropicum* and their protective effects against lipid peroxidation. *Food Research International*. 2007;40(3):356-364.

162. Ibrahim MA et al. Rutin and flavone analogs as prospective SARS-CoV-2 main protease inhibitors: In silico drug discovery study. *Journal of Molecular Graphics and Modelling*. 2021;107904.
163. Rizzuti B et al. Rutin is a low micromolar inhibitor of SARS-CoV-2 main protease 3CLpro: Implications for drug design of quercetin analogs. *Biomedicines*. 2021; 9(4):375.
164. da Silva FMA et al. Flavonoid glycosides and their putative human metabolites as potential inhibitors of the SARS-CoV-2 main protease (Mpro) and RNA-dependent RNA polymerase (RdRp). *Memórias do Instituto Oswaldo Cruz*. 2020;115.
165. Huang YC et al. Rutin improves endotoxin-induced acute lung injury via inhibition of iNOS and VCAM-1 expression. *Environ Toxicol*. 2016;31(2):185-91.
166. Dar MA, Tabassum N. Rutin-potent natural thrombolytic agent. *International Current Pharmaceutical Journal*. 2012;1(12):431-435.
167. Barrington R et al. Absorption, conjugation and efflux of the flavonoids, kaempferol and galangin, using the intestinal CaCo-2/TC7 cell model. *Journal of Functional Foods*. 2009;1(1):74-87.
168. Imani A et al. Molecular mechanisms of anticancer effect of rutin. *Phytotherapy Research*. 2021;35(5):2500-2513.
169. Yang QY, Tian XY, Fang WS. Bioactive coumarins from *Boenninghausenia sessilicarpa*. *Journal of Asian Natural Products Research*. 2007;9(1):59-65.
170. Dean FM et al. Leptodactylone, a yellow coumarin from *Leptodactylon* and *Linanthus* species. *Phytochemistry*. 1978; 17(3):505-509.
171. Salminen A et al. Celastrol: Molecular targets of thunder god vine. *Biochemical and biophysical research communications*. 2010;394(3):439-442.
172. Ng SW et al. Molecular modulators of celastrol as the keystones for its diverse pharmacological activities. *Biomedicine & Pharmacotherapy*. 2019;109:1785-1792.
173. Caruso F et al. Interrelated mechanism by which the methide quinone celastrol, obtained from the roots of *tripterygium wilfordii*, inhibits main protease 3CLpro of COVID-19 and Acts as Superoxide Radical Scavenger. *International Journal of Molecular Sciences*. 2020;21(23):9266.
174. Habtemariam S et al. Should we try the antiinflammatory natural product, celastrol, for COVID-19? *Phytotherapy Research*. 2020;1198-1190.
175. Zhang J et al. Oral bioavailability and gender-related pharmacokinetics of celastrol following administration of pure celastrol and its related tablets in rats. *Journal of Ethnopharmacology*. 2012; 144(1):195-200.
176. Qi X et al. Solid self-microemulsifying dispersible tablets of celastrol: Formulation development, characterization and bioavailability evaluation. *International Journal of Pharmaceutics*. 2014;472(1-2): 40-47.
177. Freag MS, Saleh WM, Abdallah OY. Self-assembled phospholipid-based phytosomal nanocarriers as promising platforms for improving oral bioavailability of the anticancer celastrol. *International Journal of Pharmaceutics*. 2018;535(1-2):18 - 26.
178. Wu M et al. Celastrol aggravates LPS-induced inflammation and injuries of liver and kidney in mice. *American Journal of Translational Research*. 2018;10(7): 2078.
179. Xu W et al. Absolute configuration of tetrandrine and isotetrandrine influences their anti-proliferation effects in human T cells via different regulation of NF-κB. *Zeitschrift für Naturforschung C*. 2020; 1(ahead-of-print).
180. Heister PM, Poston RN. Pharmacological hypothesis: TPC2 antagonist tetrandrine as a potential therapeutic agent for COVID-19. *Pharmacology Research & Perspectives*. 2020;8(5):e00653.
181. Su W et al. Inhalation of tetrandrine-hydroxypropyl-β-cyclodextrin inclusion complexes for pulmonary fibrosis Treatment. *Mol Pharm*. 2020;17(5):1596-1607.
182. Kim HJ, Park GM, Kim JK. Anti-inflammatory effect of pristimerin on lipopolysaccharide-induced inflammatory responses in murine macrophages. *Archives of Pharmacal Research*. 2013. 36(4):495-500.
183. Enzolifescience. Pristimerin; 2021. Available:<https://www.enzolifesciences.com/ALX-350-411/pristimerin/>.
184. Gastaldi S et al. New non-covalent inhibitors of NLRP3 inflammasome atpase activity. in XXVI National Meeting in Medicinal Chemistry; 2019.

185. Yang H et al. Pristimerin induces apoptosis by targeting the proteasome in prostate cancer cells. *Journal of Cellular Biochemistry*. 2008;103(1):234-244.
186. Dong C et al. Absorption and metabolism characteristics of pristimerin as determined by a sensitive and reliable LC–MS/MS method. *Fitoterapia*. 2015;106:62-67.
187. Saeedi-Boroujeni A et al. COVID-19: A case for inhibiting NLRP3 inflammasome, suppression of inflammation with curcumin? *Basic Clin Pharmacol Toxicol*, 2021;128(1):37-45.
188. Thimmulappa RK et al. Antiviral and immunomodulatory activity of curcumin: A case for prophylactic therapy for COVID-19. *Heliyon*. 2021;7(2):e06350.
189. Utomo RY, Meiyanto E. Revealing the potency of citrus and galangal constituents to halt SARS-CoV-2 infection; 2020.
190. Khaerunnisa S et al. Potential inhibitor of COVID-19 main protease (Mpro) from several medicinal plant compounds by molecular docking study. 2020;1-18.
191. Ting D et al. Multisite inhibitors for enteric coronavirus: Antiviral cationic carbon dots based on curcumin. *ACS Applied Nano Materials*. 2018;1(10):5451-5459.
192. Dai J et al. Inhibition of curcumin on influenza A virus infection and influenzal pneumonia via oxidative stress, TLR2/4, p38/JNK MAPK and NF-κB pathways. *International Immunopharmacology*. 2018; 54:177-187.
193. Avasarala S et al. Curcumin modulates the inflammatory response and inhibits subsequent fibrosis in a mouse model of viral-induced acute respiratory distress syndrome. *PLoS One*. 2013;8(2):e57285.
194. Xu M, et al. Effects of curcumin in treatment of experimental pulmonary fibrosis: A comparison with hydrocortisone. *Journal of Ethnopharmacology*. 2007; 112(2):292-299.
195. Titto M et al. Curcumin prophylaxis refurbishes alveolar epithelial barrier integrity and alveolar fluid clearance under hypoxia. *Respiratory Physiology & Neurobiology*. 2020;274:103336.
196. Nelson KM et al. The essential medicinal chemistry of curcumin: Miniperspective. *Journal of medicinal chemistry*. 2017; 60(5):1620-1637.
197. Ryan JL et al. Curcumin for radiation dermatitis: A randomized, double-blind, placebo-controlled clinical trial of thirty breast cancer patients. *Radiation Research*. 2013;180(1):34-43.
198. Luber RP et al. Turmeric induced liver injury: A report of two cases. *Case reports in Hepatology*. 2019;2019:1-4.
199. Balaji S, Chempakam B. Toxicity prediction of compounds from turmeric (*Curcuma longa* L). *Food and Chemical Toxicology*. 2010;48(10):2951-2959.
200. Zhang X, Hu J, Chen Y. BTA and the pharmacological effects of tumor suppression. *Molecular medicine reports*. 2016;14(5):4489-4495.
201. Hordyjewska A et al. Betulin and betulinic acid: triterpenoids derivatives with a powerful biological potential. *Phytochemistry Reviews*. 2019;18(3):929-951.
202. Wen CC et al. Specific plant terpenoids and lignoids possess potent antiviral activities against severe acute respiratory syndrome coronavirus. *Journal of Medicinal Chemistry*. 2007;50(17):4087-4095.
203. Pillaiyar T et al. An overview of severe acute respiratory syndrome–coronavirus (SARS-CoV) 3CL protease inhibitors: Peptidomimetics and small molecule chemotherapy. *Journal of Medicinal Chemistry*. 2016;59(14):6595-6628.
204. Nader MA, Baraka HN. Effect of BTA on neutrophil recruitment and inflammatory mediator expression in lipopolysaccharide-induced lung inflammation in rats. *Eur J Pharm Sci*. 2012;46(1-2):106-13.
205. Lingaraju MC et al. BTA attenuates lung injury by modulation of inflammatory cytokine response in experimentally-induced polymicrobial sepsis in mice. *Cytokine*. 2015;71(1):101-8.
206. Gauthier C et al. Glycosidation of lupane-type triterpenoids as potent in vitro cytotoxic agents. *Bioorganic & Medicinal Chemistry*. 2006;14(19):6713-6725.
207. Dutta D et al. Nanoencapsulated BTA analogue distinctively improves colorectal carcinoma in vitro and in vivo. *Sci Rep*. 2019;9(1):11506.
208. Zeng AQ et al. BTA impairs metastasis and reduces immunosuppressive cells in breast cancer models. *Oncotarget*. 2018;9(3): 3794.
209. Sasaki K et al. Improvement in the bioavailability of poorly absorbed glycyrrhizin via various non-vascular administration routes in rats. *International*

- Journal of Pharmaceutics. 2003;265(1-2):95-102.
210. Santos A.C, F. Veiga, A.J. Ribeiro. New delivery systems to improve the bioavailability of resveratrol. Expert opinion on drug delivery. 2011;8(8):973-990.
211. Gambini J et al. Properties of resveratrol: in vitro and in vivo studies about metabolism, bioavailability, and biological effects in animal models and humans. Oxidative Medicine and Cellular Longevity;2015.
212. Liu W. et al. Sensitive and robust UPLC–MS/MS method to determine the gender-dependent pharmacokinetics in rats of emodin and its glucuronide. Journal of Pharmaceutical and Biomedical Analysis. 2011;54(5):1157-1162.
213. Sirtori CR. Aescin: pharmacology, pharmacokinetics and therapeutic profile. Pharmacological Research, 2001;44(3): 183-193.

© 2021 Ali et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/76654>