



Effects of Seed Source, Moisture Content and Duration of Storage on the Viability of *Vitellaria paradoxa* C.F. Gaertn

A. F. Aderounmu¹ and I. O. Asinwa^{2*}

¹*Federal College of Forestry, Jericho Hill Ibadan, Nigeria.*

²*Forestry Research Institute of Nigeria, Jericho Hill Ibadan, Nigeria.*

Authors' contributions

This work was carried out in collaboration between both authors. Author AFA designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Author IOA managed the analyses of the study and the literature searches. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JABB/2019V22i130107

Editor(s):

(1) Dr. Andrzej Kowalski, Department of Biochemistry and Genetics, Institute of Biology, Jan Kochanowski University, Kielce, Poland.

Reviewers:

(1) Afolabi, Qasim Olaitan, Federal College of Animal Health and Production Technology, Nigeria.

(2) Ana Maria Arambarri, La Plata National University, Argentina.

(3) Benjawan Chutichudet, Mahasarakham University, Thailand

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50655>

Original Research Article

Received 19 June 2019
Accepted 08 August 2019
Published 16 August 2019

ABSTRACT

Viability of tropical seeds is crucial in plantation establishment. Temperature – moisture duration relationship of the seed is of great significance in seed viability. 3000 ripe fruits of *V. paradoxa* C. F. Gaertn were collected from each of three sources where it is endemic: Eruwa, Sakiand New-Bussa. The fruits were depulped; a sample of 100 seeds was drawn from each of the three sources and sown in washed and sterilized river sand to monitor the germination percentage of the seeds. The seeds were dried under dehumidifier to attain two moisture contents (43% and 32%). Thereafter, each seed lot from the three sources was divided into two batches (A and B) of 1200 seeds each, divided into four and stored at four different temperature regimes: -20°C, 5°C, 22°C and 28± 2°C. Samples were taken monthly for germination test for six months. Data were subjected to percentages and analysis of variance (ANOVA). Results showed that freshly collected seeds of *V. paradoxa* had 98% germination when sown within 48 hours of fallen and collection which eventually dropped to 4.4% after the six months storage. Storage temperature had significant effect ($P > 0.05$)

*Corresponding author: E-mail: israelasinwa@gmail.com;

on germination of the seeds. Seeds stored at $28\pm 2^{\circ}\text{C}$ had the highest mean germination (72.2%), those stored at 22°C had 70.2% while seeds stored at -20°C and 5°C failed to germinate. The seeds of *V. paradoxa* readily lost viability with time, hence fallen fruits should be collected during fruiting season processed and sown immediately or within 7 days of collection.

Keywords: *Vitellaria paradoxa*; germination; moisture content; storage temperature; duration of storage; seed source.

1. INTRODUCTION

Quality tree plantings with outputs marching the expectations of the tree planters require good quality seed with good germination potential and good genetic properties [1]. In order to ensure successful stand establishment, the need for effective and regular supply of seeds of high genetic and physiological qualities should not be overlooked [2]. This is because seed constitutes unique genetic composition from combined parental genetic materials. Seeds must be viable and available at the appropriate sowing season for production of seedlings for plantation establishment. Most indigenous tree species like many other trees flower and fruit once a year while some do every other year. Therefore, availability of viable seeds is very vital. Seed storage is an important factor in the preservation of viable seeds from the time of collection until they are required for sowing [3,1,2].

Seed production, quality and quantity in most species are not stable from year to year and are influenced by several factors including climate, pest and diseases, animal, human interference and forest fires. In view of this, consideration must be given to appropriate storage to maximize the seeds collected [4]. [5] stated that the two most crucial necessities for storing seeds are constant temperatures and low humidity; although, seeds from different sources respond differently to storage temperatures and durations.

Vitellaria paradoxa C. F. Gaertn seed is recalcitrant hence require adequate silvicultural attention for viability. It is a dicotyledonous plant in the family Sapotaceae. It is commonly known as butter tree, shea tree, shea butter tree or shea nut tree [6]. The seed has high oil content, oil rancidity during storage, oil freezing in cold condition and inaccessibility of the embryo for the supply of the stored nutrient [7]. Recalcitrant seeds have low tolerance to drying and cooling. Many recalcitrant species contain about 40% moisture at maturity and do not tolerate drying to less than 20-30%. Similarly, *V. paradoxa* had various nut fresh weights ranging from 8-10 g

containing 43-68% moisture by weight [8]. In contrast, most orthodox seeds can be dried below 5% moisture content without losing viability. Therefore, this study was carried out to highlight the germination response of *V. paradoxa* seed from different sources to different moisture contents, storage temperature and duration of storage with a view to recommending best storage option(s) for the species with which optimum germination could be achieved .

2. MATERIALS AND METHODS

The study was carried out in three locations namely Eruwa, Saki and New-Bussa in Nigeria. Ripe fruits of *V. paradoxa* were collected from three sites along its natural range: Eruwa (latitudes $7^{\circ}47' \text{N}$ and $7^{\circ}42' \text{N}$; longitudes $3^{\circ}51' \text{E}$ and $3^{\circ}47' \text{E}$); Saki (latitudes $8^{\circ}40' \text{N}$ and $8^{\circ}36' \text{N}$; longitudes $3^{\circ}19' \text{E}$ and $3^{\circ}16' \text{E}$) and New-Bussa (latitudes $9^{\circ}35' \text{N}$ and $9^{\circ}33' \text{N}$; longitudes $3^{\circ}33' \text{E}$ and $3^{\circ}29' \text{E}$) in the derived, southern guinea and northern guinea savanna ecological zones respectively. The forest floor around the selected trees were cleared of fruits, leaves, twigs and fallen fruits before the fallen ripe fruits were collected daily for a period of 7 days. A total of 1500 fruits were collected from each source and packed in jute bags to the laboratory where depulping was carried out to extract the seeds and thereafter separated from dirts. A Sample of 100 clean seeds was drawn from each of the three sources and immediately sown in washed and sterilized river sand under a non-mist propagator to monitor the initial germination percentage of the seed. Each seed lot (2400) from the three sources was divided into two batches of 1200 seeds each (A and B) and dried to two moisture contents (43% and 32%) under a dehumidifier at the Tree Improvement Nursery, Forestry Research Institute of Nigeria (FRIN), Ibadan Nigeria. The moisture content was determined by weighing seeds on a sensitive weighing balance and there after oven drying at 60°C for 17 hours until they attained constant dry weight.

Seedlots from the three sources grouped into A and B were each divided into four (300

seeds/pack) and packed into transparent re-sealable plastic containers before storage at four different temperatures regimes: (-20°C), (5°C), (22°C) and (28± 2°C). The seeds were stored in the Gene bank at National Centre for Genetic Resources and Biotechnology (NACGRAB), Ibadan. Samples of *V. paradoxa* seeds stored under the four temperatures were taken from each batch fortnightly and sown in washed and sterilized river sand under a non-mist propagator. Germination percentage was monitored fortnightly for 6 months. Twelve sampling periods were observed and germination was monitored for two weeks and percentage germination within two weeks of sowing determined. The experimental design used was 3 x 6 x 4 x 2 factorial experiments in Completely Randomized Design where the first factor was the three seed sources, the second factor was six storage duration, the third factor was four storage temperature and the fourth factor was two moisture contents, the treatment combinations were replicated four times. The data collected were subjected to Analysis of variance (ANOVA) and where significant differences were obtained using IBM SPSS software, least significant difference (LSD) test was used to separate the means.

3. RESULTS AND DISCUSSION

3.1 Results

Effect of seed source and moisture content on percentage germination of the different seed lots: The germination of the seeds of *V. paradoxa* varied from 94 to 98%. Seeds from Saki had the highest initial germination of 98%, followed by the seeds from New-Bussa with 97% while the lowest initial percentage germination of 94% was observed among the seeds collected from Eruwa (Table 1). However, sources of seeds and moisture contents did not show any significant effect on the percentage germination of the seeds (Table 2).

Effect of temperature on percentage germination of stored seeds: Storage temperature and duration of storage had

significant effect ($p \leq 0.05$) on germination of the seeds from the sources. The mean percentage germination of seeds stored at 28±2°C and 22°C were not significantly different from each other but differed significantly from the seeds stored at -20°C and 5°C. However, the seeds stored at 28±2°C had higher mean germination (72.2%) than those stored at 22°C with 70.2% while seeds stored at -20°C and 5°C had zero germination each (Table 3).

Effect of duration of storage on percentage germination of stored seeds: Duration of storage substantially influenced the percentage germination of the seeds from the three sources under the four temperature regimes throughout the period of assessment. The variation pattern showed a systematic decline in percentage germination in all the three seedlots as well as in seeds stored at 28±2°C and 22°C over the storage periods. The seedlots from all the sources stored at 28±2°C and 22°C temperature regimes showed consistent decrease in percentage germination ranging from 88.4 to 5.6% while the seeds stored at -20°C and 5°C exhibited sudden germination failure (Table 3).

Effect of temperature and duration of storage on germination of stored seeds: Interaction of storage period and temperature had significant effect on the percentage germination of different seedlots. The percentage germination of seedlots stored at -20°C and 5°C were not significantly different from one another; but were significantly different from those kept in -20°C and 5°C only in the 1st month (Table 2).

4. DISCUSSION

Germination of viable seeds is dependent on varying temperature and moisture contents and other environmental conditions which seeds must be adapted to [9]. Each plant species has a specific optimum and a range within which germination will occur [10,11]. The closer the temperature is to optimum temperature required by seed, the quicker the germination will occur [12,13,14].

Table 1. Germination percentage of *V. paradoxa* seeds from three sources

Seed source	GPS coordinates		Germination %
	Latitude	Longitude	
Saki	8°40'N and 8°36'N	3°19'E and 3°16'E	98
New Bussa	9°35'N and 9°33'N	3°33'E and 3°29'E	97
Eruwa	7°47' N and 7°42'N	3°51'E and 3°47'E	94

Table 2. Analysis of variance for the percentage germination of *V. paradoxa* seeds subjected to viability variables

Parameters	df	ms	f-value	P-level
Source (S)	2	0.04	0.10ns	0.902
Storage duration (SD)	5	42.04	98.69*	0.000*
Storage temperature (ST)	3	114.41	268.61*	0.000*
Moisture content (MC)	1	0.52	1.22ns	0.270
S X SD	10	0.16	0.37ns	0.959
S X ST	6	0.36	0.82ns	0.998
SD X ST	15	14.24	33.44*	0.000*
S X MC	2	0.13	0.31ns	0.734
SD X MC	5	0.51	1.20ns	0.311
ST X MC	3	0.25	0.59ns	0.625
S X SD X ST	30	0.12	0.27ns	0.999
S X SD X MC	10	0.16	0.38ns	0.954
S X ST X MC	6	0.55	0.38ns	0.954
SD X ST X MC	15	0.25	0.58ns	0.892
S X SD X ST X MC	30	0.17	0.40ns	0.998
Error	288	0.43		

Note: *Significant at 5% Probability level, ns = not significant, df = Degree of Freedom, ms = Multiple of Square

Table 3. Mean values of percentage germination of *V. paradoxa* seeds subjected to different temperature and duration of storage

Parameters	Mean germination %
Storage temperature	
-20°C	0.00 ^a
5°C	0.00 ^a
22±1°C	70.4 ^b
28±2°C	72.4 ^b
Duration of storage (Months)	
6	5.6 ^a
5	11.2 ^a
4	22.4 ^b
3	36.8 ^c
2	50.0 ^d
1	88.4 ^e

Note: Means with the same superscripts are not significantly different from each other at $p \leq 0.05$

Seeds of *V. paradoxa* irrespective of source responded to varied storage temperatures as well as storage durations. Germination which was highest when freshly collected seeds were sown and declined in a geometrical progression when stored at 22±1°C and 28±2°C affirm the recalcitrant nature of the species. This implies that its recalcitrant nature of the seeds prevail the effects of the sources. According to [6], *V. paradoxa* is a recalcitrant species which are remarkably short-lived and cannot be dried to moisture content below 20-30% without injury and are unable to tolerate freezing. They are also termed as desiccation sensitive seeds, difficult to be successfully stored and their ex-situ conservation is problematic [2]. This implies that the species should be sown immediately after collection as a substantial level of germination

would be achieved. This finding conforms with [9] who reported that *V. paradoxa* is recalcitrant in nature and thus has short physiological storability and viability status. [15] reported that seed quality maintenance in storage from the time of production until the seed is planted is imperative to assure its planting value [15]. This implies that decrease in seed viability with storage duration in *V. paradoxa* seeds could be ascribed to influence of storage conditions in terms of temperature. The findings of this study where 32 and 43% moisture contents had no significant effect on the viability of the seeds corroborates the result of [16], on storage trials of *V. paradoxa* in Burkina Faso and Senegal with regulated storage temperature and moisture contents over a period of 6 months where the drying to moisture content above 37% did not affect germination percentage

of the seeds of *V. paradoxa*. This is also in line with findings of [17] that mortality of the seeds of *V. paradoxa* increased at moisture contents that ranged from 20 – 26%.

The significance of the effects of storage duration and temperature on germination percentage of *V. paradoxa* seeds recorded in his study is in line with [18] who recorded highest germination in the 1st month at 20-25°C and progressive decline in germination as the duration of storage increased confirming the seed to be recalcitrant and losing viability very rapidly. On the other hand, lower germinations were recorded in low temperature storage. This is probably because of the recalcitrant nature of the seed which requires high moisture for optimum germination [18].

Seed moisture content is a function of environmental conditions which is chiefly influenced by humidity and temperature of the environment. Invariably, the main aim of storing seeds is to maintain the critical moisture content required for the optimum germination or recondition it to attain the optimum moisture level either through drying or soaking [19,20]. The decrease in germination potentials with increase in storage period and decreased temperature confirms the recalcitrant nature of the seed which result in impairment of certain physiological activities at temperatures below the optimum required. This implies that moisture and temperature are basic requirements for germination, they are essential for enzyme activation, breakdown, translocation, and use of reserve storage material. In their resting state, seeds are characteristically low in moisture and relatively inactive metabolically. That is, they are in a state of quiescence [19]. Thus, quiescent seeds are able to maintain a minimum level of metabolic activity that ensures their long-term survival in the soil and during storage. Moisture availability is described in various ways. Field capacity moisture is about optimum for germination in soil; however, germination varies among species and may occur at soil moistures near the permanent wilting point [13]. Most seeds have critical moisture content for germination to occur. For example, Orthodox seeds which are long-lived seeds can be successfully dried to moisture contents as low as 5% without injury while species such as *Chrysophyllum albidum* and *Caesalpinia bonduc* which are recalcitrant seeds cannot germinate readily at moisture content below 20% [21]. Once that critical seed moisture content is attained in the seed, sufficient water is present to initiate germination [22]. On the other hand, if internal moisture

content decreases below the critical moisture content, seeds will essentially decay in the soil [19]. Seed germination is a complex process involving many individual reactions and phases, each of which is affected by temperature [20]. The characteristics exhibited by *V. paradoxa* may be an indication of ecological adaptation of the species since its natural range is within the savanna eco region where temperature is relatively high all the year round. This implies that low temperatures do not favour the viability of the seeds.

5. CONCLUSION

It could be inferred from this finding that seeds of *V. paradoxa* could be collected from any of the sources within the distribution range of the species most especially for germination purpose. There is obvious evidence that the seeds should be sown fresh since seeds from all the sources recorded impressive initial germination percentage. Nursery managers and silviculturists who want to raise the seedlings don't have to bother about provision of cold storage. This information is crucial in ensuring successful germplasm conservation programme so that seedlots that lose their viability quickly may be kept in storage for a short period and sown out before they lose their viability. However, further studies are needed on improving the longevity in storage of *V. paradoxa* seed.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Davidson RH, Edwards DG, Sziklai WO, El-Kassaby YA. Variation in germination parameters among Pacific silver fir populations. *Silvae Genet.* 1996;45:165-171.
2. Toon PG, Haines RJ, Dieters MJ. Relationship between seed weight, germination and seedling-height growth in *Pinus caribae*. Morele. var. hondurensis barre and Golfri. *Seed Science Technology.* 1990;19:389-402.
3. Bewley JD, Bradford K, Hilhorst H. *Seeds: Physiology of development, germination and dormancy.* Springer Science and Business Media. 2012;392.
4. Vertucii CW, Crane J, Vane NC. Physiological aspects of *Taxus brevifolia*. *Physiologia planetarium.* 1996;98:1-12.

5. Lady BJ. Guidelines for seed collection. 18. Available:<http://www.wildflower.org/2002>
6. Hall JB, Aebischer DP, Tomlinson HF, Osei-Amaning A, Hindle JR. *Vitellaria paradoxa*, A Monograph. School of Agricultural and Forest Sciences, University of Bangor. UK. 1996;105.
7. Lamien N, Tigabu M, Guinko S, Oden PC. Variations in dendrometric and fruiting characters of *Vitellaria paradoxa* populations and multivariate models for estimation of fruit yield. *Agroforestry Systems*. 2007;69: 1–11.
8. Bonkougou EG. The shea tree (*Vitellaria paradoxa*) and African Shea parklands, Proceedings of the International Workshop on Processing and Marketing of Shea Products in Africa, Sénégal, 4-6 March 2002. CFC Technical paper No 21, Food and Agriculture Organization. Rome; 2005.
9. Gamaéné SH, Pritchard C, Harris W. *Vitellaria paradoxa*. The project on handling and storage of recalcitrant and intermediate tropical forest tree seeds, Newsletter. Danida Forest Seed Centre, Humlebæk, Denmark. 1999;5:15-16.
10. Cibele CM, Bovi ML, Nakagawa J. Desiccation effects on germination and vigor of King palm seeds. *Hort Brasileira*. 2003;21:88-92.
11. Shaban M. Application of seed equilibrium moisture curves in agro physics. *International Journal of Advanced Biological and Biomedical Research*. 2013;1(9):885-898.
12. Baskin JM, Baskin CC. A classification system for seed dormancy. *Seed Sci. Res*. 2004;14:1–16.
13. Beyranvand H, Farnia A, Nakhjavan SH, Shaban M. Response of yield and yield components of maize (*Zea mayz* L.) to different bio fertilizers. *International Journal of Advanced Biological and Biomedical Research*. 2013;1(9):1068-1077.
14. Bradbeer JW. Seed dormancy and germination. Springer Science and Business Media, 2013;146.
15. Roberts EH. Temperature and seed germination. in Long SP, Woodward FI. eds. *Plants and Temperature*. Cambridge, U.K: Society for Experimental. 1988;109–132.
16. Pritchard HW, Yameogo CS, Daws ML. Desiccation, storage and germination of *Butyrospermum paradoxum* (*Vitellaria paradoxa*) seed from Burkina Faso. *International Plant Genetic Resources Institute Annual Report*. 1996;51.
17. Daws ML, Gamene CS, Glidewell SM, Pritchard HW. Seed mass variation potentially masks single critical water content in recalcitrant seeds. *Seed Science Research*. 2004;14(02):185–195. DOI: 10.1079/SSR2004168
18. Lars S. Guide to handling of tropical and subtropical forest seed (eds) Kersten, O. Danida Forest Seed Centre. 2000;511.
19. Bradford KJ. Water relations in seed germination. In J. Kigel and G. Galili, eds. *Seed Development and Germination*. New York, Marcel Dekker, Inc. 1995;351–396.
20. Covell S, Ellis RH, Roberts EH, Summerfield RJ. The influence of temperature on seed germination rate in grain legumes. I. A comparison of chickpea, lentil, soybean, and cowpea at constant temperatures. *Journal of Botanical Science*. 1986;37:705–715.
21. Covell S, Ellis RH, Roberts EH, Summerfield RJ. The influence of temperature on seed germination rate in grain legumes. I. A comparison of chickpea, lentil, soybean, and cowpea at constant temperatures. *Journal of Botanical Science*. 1986;37:705–715.
22. Ellis RH, Covell S, Roberts EH, Summerfield RJ. The influence of temperature on seed germination rate in grain legumes. II. Intraspecific variation in chickpea (*Cicer arietinum* L.) at constant temperatures. *Journal. Exp. Botany*. 1986; 37:1503-1515.

© 2019 Aderounmu and Asinwa; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/50655>