



Water Use Efficiency of Maize Genotypes of Different Maturity Groups at Seedling and Grain-filling Growth Stages in a Rainforest Location

O. O. Bankole¹, A. Oluwaranti^{1*} and F. E. Awosanmi¹

¹Department of Crop Production and Protection, Obafemi Awolowo University, Ile-Ife, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author OOB managed the study, performed the statistical analysis, managed the literature searches and wrote the first draft of the manuscript. Author AO designed the study, supervised the study and wrote the final draft and author FEA assisted in the supervision of the study and contributed to the final draft. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2019/v29i530154

Editor(s):

(1) Dr. Abhishek Naik, Acen Hyveg Pvt. Ltd., Vegetable Seed Division, West Bengal, India.

Reviewers:

(1) Cristiane Ramos Vieira, University of Cuiabá, Brazil.

(2) R. Mahalakshmi, India.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50836>

Original Research Article

Received 10 June 2019
Accepted 15 August 2019
Published 17 September 2019

ABSTRACT

Aims: The objectives of this study were to evaluate maize genotypes of different maturity groups for seedling and grain filling water use efficiency and determine relationship that exist between the water use efficiency traits and yield of different maize maturity groups.

Study Design: Sixteen maize genotypes were planted in Randomized Complete Block Design in three replicates for emergence, vegetative, water use efficiency traits at the seedling and grain-filling growth stages and yield.

Place and Duration of Study: The sixteen maize genotypes of different maturity groups were evaluated during the early and late cropping seasons of 2016 at the Obafemi Awolowo University Teaching and Research Farm, Ile-Ife, Nigeria

Methodology: Data collected were subjected to Analysis of Variance (ANOVA), correlation analysis among water use efficiency traits and yield for each of the maturity groups.

Results: There was no significant difference among the genotypes within each maturity groups for water use efficiency at seedling and grain filling growth stages.

*Corresponding author: E-mail: boluwaranti@yahoo.com;

The late maturity group of maize used more water at the seedling growth stage than the other maturity groups in the early season of this study while in the late season, the early and extra-early maturity groups used more water than the other maturity groups. Increase in emergence percentage, reduction in speed of germination, and minimal days to complete germination increased water use efficiency at the seedling stage only during the early cropping season. Efficiency of water usage at the seedling growth stage was more among the late and intermediate maturing groups than the extra-early and early maturing groups in the early season while in the late season, the extra-early and early maturing groups used water more efficiently than the late and Intermediate maturing groups

Conclusion: Maturity group played a significant role in the expression and manifestation of water use efficiency traits under different environmental conditions.

Keywords: Maize; maturity groups; water-use efficiency; seedling growth stage; grain filling growth stage.

1. INTRODUCTION

Maize is a cereal crop that is more extensively distributed globally than any other cereal crops because of its wide adaptability to range of climates. It was also reported that maize is the most widely-grown staple food crop in sub-Saharan Africa (SSA) occupying more than 33 million ha each year [1]. Multi-purpose uses of maize have made it a popular and most widely cultivated crop after wheat and rice in the whole world. This is so because maize contributes about 34 - 36% of the average daily calorie intake [2-4]. Maize is used for producing alcohol and nonalcoholic drinks. The stem serves as an important source of bio-fuel [5-7]. It is estimated that maize demand in Sub-Saharan Africa would exceed 52 million tons in 2020 [8]. Despite an increased area of land which has been dedicated to cultivate maize since the mid-2000s, production per hectare in the developing countries is still low (1.3 t/ha) compared to the 8.6 t/ha in developed countries [9].

In most crops, the variation in biomass accumulation and yield production is influenced by the crop's tolerance to water stress, drought, and the efficiency with which the maize crop uses available soil water for growth. This led to the concept of Water Use Efficiency which was broadly defined as the measure of the crop production per unit of water used, irrespective of water source, expressed in units per weight of water depth per unit area, and it can also be explained as the ratio of crop yield over applied water. According to Jensen [10], efficiency has been defined as the ability to produce desired effect with minimum effort, expenses, and waste. Several reports had shown the relationship between water use efficiency and maize production [11-14]. A linear relationship was

reported by [15-18,13] between grain yield and water use efficiency in maize, that is, increase in water use efficiency will lead to an increase in grain yield and vice versa, while a curvi-linear relationship was established between water use efficiency and grain yield of maize as reported by Yazar et al. [16].

Maize has different responses to water deficit according to its developmental stages [19,20]. Drought stress is particularly damaging to grain yield if it occurs early in the growing season (plant establishment), at flowering and during mid to late grain filling [21]. The most critical period for water stress in maize is between 10-14 days before and after flowering, with reduction of 2-3 times more when water deficit coincides with flowering compared with other growing stages [22].

Different maturity groups of maize can be used to rescue shortage of maize supply, to ensure adequate all year round cropping in both seasons, and to close down the gap of high demand for maize. Oluwaranti et al. [23] reported that, different maturity groups of maize have different quality that makes them acceptable as a variety of maize. As reported by Shaibu et al. [24] each maturity group of maize also has its unique advantages and disadvantages with respect to climatic conditions. Drought tolerant maize of different maturity groups has been developed by maize breeders but there is little or no information on how water usage at the seedling and grain filling growth stages of the maize plant are being influenced by their different maturity groups. Therefore, the aim of this study was to evaluate the variations that exist among the genotypes within different maturity group for water use efficiency traits at the seedling and grain filling growth stages and determine the

relationship between the maturity groups and efficiency of water usage at the seedling and grain filling growth stages in a rainforest location.

2. MATERIALS AND METHODS

2.1 Data Collection

Sixteen drought-tolerant maize varieties, consisting of four varieties each of Extra early, Early, Intermediate and Late maturity groups obtained from the Maize Breeding Programme of International Institute of Tropical Agriculture (IITA), Ibadan, Nigeria were used for the study.

The field experiments were carried out at Obafemi Awolowo University Teaching and Research Farm (T&RF), Ile-Ife (latitude 7°25' N, and longitude 4°39'E), Nigeria, during the early and late cropping seasons of 2016. The experiment was laid out in a randomized complete block design with three (3) replicates. Plots were six rows; 5 m long each, with intra and inter-row spacing of 0.5m and 0.75m, respectively.

Emergence counts were taken on 5, 7, and 9 days after planting (DAP) to obtain emergence percentage. Data on fresh weight (FWT) of the maize plants commenced nine DAP and continued at five days interval to 39 DAP. Fresh weight of five ears per plot was taken 60 DAP at five days interval till 90 DAP for extra-early and early maturity groups and started 65DAP till 95 DAP for intermediate and late maturity groups. Dry weight (DWT) of the collected maize plants was determined in the laboratory by oven drying at 80°C to constant weight. Water stored (WSTD) was obtained from the difference between Fresh and Dry weights of the samples. Water used for evapotranspiration was obtained from the product of Potential Evapotranspiration (PET) of the research field and Single Crop Coefficient, (Kc), where (Kc) for maize = 0.35 (FAO 1998). Weather data on the potential evapotranspiration (PET) of the field was obtained from the Automatic Weather Station located at the Teaching and Research Farm, OAU, Ile-Ife. Cumulative Water Used was obtained from the addition of the water stored with the water used for Evapotranspiration by the crop at seedling and grain filling stages. Water Use Efficiency (WUE) for each maize variety was estimated from dry weight of the sample and cumulative water used (Water Use Efficiency (WUE) = DWT/ CWU). Seedling Growth Rate (SDGR) and the Grain Filling Growth Rate

(GFGR); which measures the rate of dry matter production per Unit of time measured as g/day was obtained by regression method with this linear regression model:

$$W = a + bt$$

where,

W = dry weight per plant
t = time in DAP
a = intercept of the regression model and
b = regression coefficient which measures the growth rate (GR)

2.3 Statistical Analyses

Data collected from the field experiments were subjected to statistical analysis of variance (ANOVA) using SAS package version 9.0 of statistical analysis (SAS, 2002). The differences among treatment means were separated using Least Significant Difference (LSD) at 0.05 level of probability. Pearson correlation analysis between water use efficiency traits and yield of the different maturity groups of maize were also carried out.

3. RESULTS

Mean square values from combined analysis of variance due to season was highly significant (P = 0.01) for emergence percentage, seedling and grain-filling water used, seedling and grain-filling dry weights, water use efficiency at seedling and grain-filling growth stages, growth rates at the seedling and grain-filling stages and grain yield (Table 1). Highly significant (P = 0.01) maturity effects were obtained on emergence percent, seedling and grain-filling water used, seedling dry weight, seedling growth rate and grain yield (Table 1). Highly significant (P = 0.01) interaction of the season by maturity group was observed on seedling water used, seedling dry weight, water use efficiency and growth rate at the seedling stage. Significant (P = 0.05) interaction of the season by maturity group were also observed on grain-filling dry weight, water use efficiency and growth rate at the grain-filling growth stage and grain yield (Table 1). The coefficients of variability (CVs) were generally high for emergence, seedling water used, dry weights of the seedling and the filled grains, water use efficiency at the grain filling growth stage, growth rates at the seedling and grain-filling growth stages and grain yield whereas the CV was rather low for water use efficiency at the seedling

Table 1. Mean squares from combined analysis of variance for water use efficiency traits of maize of different maturity groups at seedling and grain filling growth stages and grain yield during the early and late cropping seasons of 2016 at the Teaching and Research Farm, Obafemi Awolowo University, Ile-Ife, Nigeria

Traits	REP (d.f.=2)	SEASON (d.f.=1)	MATGRP (d.f.=3)	V(MATGRP) (d.f.=12)	S*V(MATGRP) (d.f.=12)	SEA.*MTGRP (d.f.=62)	Error	CV	R ²
E%	194.43	14412.44**	5222.25**	505.44	126.77	110.08	721.90	43.78	0.46
SDWUSD	1033.41	94077.46**	20840.20**	1205.50	2379.89	56807.01**	2706.52	35.83	0.68
SDDWT(g)	31.16	3768.16**	368.27**	16.72	34.65	687.71**	28.95	43.67	0.81
SDWUE	0.0001	0.0735**	0.0006*	0.0002	0.0003	0.0023**	0.0002	19.51	0.87
GFWUSD	1050.99	8180.32**	3181.41**	416.83	383.29	133.13	558.93	16.07	0.46
GFDWT(g)	831.88	82354.74**	364.89	348.46	618.99	1526.27*	535.20	43.27	0.75
GFWUE	0.05	3.44**	0.02	0.02	0.03	0.08*	0.02	40.87	0.77
SDGR	0.64	74.20**	6.97**	0.30	0.59	13.91**	0.58	42.19	0.81
GFGR	11.62	2365.72**	9.77	19.82	23.23	41.10*	14.95	48.39	0.77
GYLD	2.96	27.78**	5.77**	0.82	1.16	3.88*	1.22	67.22	0.53

*,** Significant at 0.05 and 0.01 levels of probability respectively

stage and water used at the grain-filling growth stage. The coefficients of determination (R^2) obtained from the model for the water use efficiency traits at the seedling and grain-filling growth stages were generally high which ranged from 68% to 87%, which indicated that the model was highly reliable (Table 1). It was observed from the results of the evaluation of water use efficiency traits and yield, that there were significantly higher means recorded for most traits in the early cropping season compared to the late cropping season of this study except for seedling water used, water use efficiency at the seedling growth stage, growth rate at the grain-filling growth stage and grain yield in which higher values were recorded in the late season (Table 2).

Differences among the means of different maturity groups of maize evaluated for water use efficiency traits, dry matter accumulated at seedling and grain filling stages and grain yield were obtained during the early and late cropping seasons of 2016 (Table 3). The late maturity group of maize used more water at the seedling growth stage than the other maturity groups in the early season of this study while in the late season, the early and extra-early maturity groups used more water than the other maturity groups. In the early cropping season, the late and intermediate maturity groups had the largest seedling dry weight and the least dry weight obtained in the early and extra-early maturity groups while in the late season, the early and extra-early maturity groups had the largest dry weights with the late and intermediate having the least dry weights (Table 3).

Efficiency of water usage at the seedling growth stage was more among the late and intermediate maturing groups than the extra-early and early maturing groups in the early season while in the late season, the extra-early and early maturing groups used water more efficiently than the late and intermediate maturing groups (Table 3). The intermediate and late maturing groups had more dry weight of filled grains than the early and extra-early maturity groups in the early season while there were no significant differences among the four maturity groups for the dry weight of the filled grain in the late season of this study (Table 3). Likewise, this trend of no significant difference among the four maturity groups in the late season for grain-filling dry weight was also obtained for the water use efficiency at the grain filling growth stage while the Late and intermediate maturity groups used water more

efficiently than the extra-early and early maturity groups in the early season (Table 3). The seedling growth rate of the late maturity group was observed to be higher than the other maturity groups in the early cropping season while in the late season, highest growth rate was observed among the early maturity group followed by the extra-early maturity group with the late maturity group having the lowest seedling growth rate in the late season of this study (Table 3). There were no significant differences among the four maturity groups for grain-filling growth rate in the late season while in the early season, the intermediate and late maturity groups had the highest growth rate at the grain-filling growth stage than the early and extra-early maturity groups (Table 3). There were no significant differences in the grain yield among the four maturity groups during the early season of this study while the early maturity group had the highest yield with the intermediate maturity group having the least in the late season of this study (Table 3).

Correlation coefficients among grain yield and water use efficiency traits at the seedling and grain-filling growth stages among the extra-early, early, intermediate and late maturing groups during the early and late cropping seasons of 2016 are presented in Tables 4 and 5.

In the early season of 2016, dry weight at the seedling stage was positively correlated with yield ($P = 0.05$) among the early and late maturity groups of maize in this study (Table 4). Highly significant ($P = 0.01$) positive correlation was obtained between water used for grain filling and yield of the early maturity group of maize used in this study (Table 4). All the maturity groups of maize in this study except the extra-early maturity group had highly significant ($P = 0.01$) positive correlation between the dry weight and water use efficiency at the grain filling stage of development and yield (Table 4). Growth rate at the grain-filling stage of development and yield had significant positive correlation among the early ($P = 0.01$) and late ($P = 0.05$) maturity groups during the early cropping season of this study (Table 4).

During the late season of 2016 of this study, highly significant positive correlation ($P = 0.01$) was obtained between the emergence percent and yield of the extra-early maturity group (Table 5). Only the late maturity group had highly significant ($P = 0.01$) positive correlation between water used for grain filling and yield

Table 2. Emergence, water use efficiency traits and grain yield of the maize genotypes of different maturity groups evaluated during the early and late cropping seasons of 2016 at the Teaching and Research Farm, Obafemi Awolowo University, Ile-Ife, Nigeria

Seasons	E%	SDWUSD	SDDWT	SDWUE	GFWUSED	GFDWT	GFWUE	SDGR	GFGR	GYLD
Early	73.62	113.88	18.59	0.05	156.33	82.76	0.55	0.92	3.03	1.11
Late	49.12	176.49	6.06	0.10	137.87	24.18	0.17	2.8	12.95	2.18
LSD _{0.05}	10.96	21.28	2.20	0.01	9.65	9.45	0.06	0.31	1.58	0.45

E%: Emergence percentage; SDWUS: Seedling water used; SDDWT: Seedling dry weight; WUE: Water use efficiency; GFWUSED: Grain filling water used; GFDWT: Grain filling dry weight; SDGR: Seedling growth rate; GFGR: Grain filling growth rate. LSD: Least significant difference

Table 3. Water use efficiency traits and grain yield of the maturity groups of maize for early and late cropping season of 2016 evaluated at the Teaching and Research Farm, Obafemi Awolowo University, Ile – Ife, Nigeria

MAT.GRPS	SDWUSD		SDDWT		SDWUE		GFDWT Seasons		GFWUE		SDGR		GFGR		GYLD	
	ES	LS	ES	LS	ES	LS	ES	LS	ES	LS	ES	LS	ES	LS	ES	LS
EE	81.94	190.55	3.15	20.94	0.032	0.109	14.37	87.76	0.113	0.609	0.50	3.10	1.67	14.23	1.05	2.12
E	97.45	265.58	5.30	30.11	0.047	0.114	18.24	90.29	0.137	0.603	0.82	4.24	2.29	14.01	0.78	2.91
I	96.27	125.95	5.06	11.67	0.048	0.091	27.55	72.90	0.178	0.451	0.78	1.72	3.58	10.73	0.90	1.09
L	179.85	123.85	10.71	11.62	0.058	0.093	36.56	80.76	0.239	0.521	1.58	1.66	4.56	12.84	1.69	2.61
LSD _{0.05}	24.40	47.93	2.00	5.87	0.012	0.007	11.33	23.78	0.067	0.177	0.30	0.74	1.39	4.75	0.79	0.74

EE: Extra early; E: Early; I: Intermediate; L: Late; ES: Early season; LS: Season 2; SDWUS: Seedling water used; SDDWT: Seedling dry weight; SDWUE: Seedling water use efficiency; GFDWT: Grain filling dry weight; GFWUE: Grain filling water use efficiency SDGR: Seedling growth rate; GFGR: Grain filling growth rate; LSD: Least significant difference

Table 4. Pearson correlation among Water use efficiency traits and grain yield of the maize genotypes of different maturity groups evaluated at the Obafemi Awolowo University Teaching and Research Farm, Ile-Ife, Nigeria during the early cropping season of 2016

		E%	SDWUSD	SDDWT	SDWUE	GFWUSD	GFDWT	GFWUE	SDGR	GFGR
YLD	EE	0.17	-0.06	0.09	0.23	-0.47	-0.16	-0.15	0.37	-0.33
	E	0.30	0.58	0.63*	0.47	0.76**	0.88**	0.85**	-0.17	0.80**
	I	-0.25	-0.21	-0.13	0.06	0.50	0.76**	0.75**	-0.17	0.49
	L	0.31	0.53	0.58*	0.43	0.50	0.75**	0.72**	0.05	0.62*

*,** Correlation is significant at the 0.05 and 0.01 levels of probability respectively

Table 5. Pearson correlation among water use efficiency traits and grain yield of the maize genotypes of different maturity groups evaluated at the Obafemi Awolowo University Teaching ANF Research Farm, Ile-Ife, Nigeria during the late cropping season of 2016

		E%	SDWUSD	SDDWT	SDWUE	GFWUSD	GFDWT	GFWUE	SDGR	GFGR
YLD	EE	0.73**	-0.07	-0.05	0.09	0.12	0.81**	-0.15	-0.01	-0.29
	E	0.47	-0.09	-0.22	-0.53	-0.13	0.24	0.47	-0.14	0.14
	I	0.55	-0.19	-0.14	0.15	-0.01	-0.01	0.22	-0.26	0.07
	L	0.17	0.44	0.36	-0.03	0.78**	-0.33	-0.40	0.36	-0.21

*,** Correlation is significant at the 0.05 and 0.01 levels of probability respectively

whereas highly significant ($P = 0.01$) positive correlation was obtained between dry weight at the grain filling stage and yield among the early maturity group in the late season of 2016 of this study (Table 5).

4. DISCUSSION

The significant difference observed among the early and late cropping seasons of this study for emergence, seedling and grain filling dry weights, water use efficiency at the seedling and grain-filling growth stages, was expected since the two cropping seasons were characterized by different amount of rainfall, temperature, sunshine hour, potential evapotranspiration (PET) among other climatic attributes. This result was also corroborated by the findings of Vina et al. [25] and Sivritepe et al. [26] which showed that abundant supplies of water is required for sustainable organ development. This indicated that with this current global climatic change, favorable seasons will determine the performance of the maturity groups of maize irrespective of their improvement.

The low coefficient of variation recorded for water used at grain filling stage showed that there were no much differences in utilization of water for different maturity groups. While the high value recorded for grain filling growth rate showed much variability in the performance of the different maturity groups which indicated that rate at which different maturity groups of maize acquire dry matter on daily basis under each season varied widely.

Low emergence observed during the late season compared to the early cropping season can be attributed to genetic limitations, environmental factor which resulted in the loss of viability and vigour, during the late season. This was also observed by Ajayi and Fakorede [27] which reported that there was significant reduction in maize seed quality after three months in storage under ambient conditions. This validated the report by Bewley and Black [28] that seed quality has significant direct influence on crop productivity levels.

The significant difference between the maturity groups for the expression of the water use efficiency traits and yield difference could have been due to the difference in the duration of time to maturity of the different maturity groups. This was also observed by Oluwaranti et al. [23] that the time for expression of traits by genotypes

varies with maturity groups. Due to the different durations of the growth stages and the amount of rainfall required by the different maize maturity groups evaluated in this study, the intermediate and late maturity groups with more than 100 – 120 days to physiological maturity used more water for both seedling and grain-filling growths during the early season because of the longer duration of the season. The efficiency of water usage and growth rates at the seedling and grain-filling growth stages was also more among the intermediate and late maturity groups than the early and extra-early maturity groups in the early season since the duration and the amount of rainfall of the early season for their growth and development are more than that of the late season. This result was also corroborated by the findings of Oluwaranti et al. [23] and Ajani et al. [29] which reported that maturity groups highly affected the number of seedlings that emerged and the speed with which they emerged. This is in support of findings by Misra, [30] that the availability of water during the different stages of crop growth also influences crop's survival. These same findings were supported by Vina et al. [25] which reported that if under favorable environmental conditions like temperature, soil moisture content and solar radiation, maize will produce maximum yield at different stages if supplied with adequate nutrients. The seasonal significance for emergence and water use efficiency traits evaluated indicated that the season is taking its toll on seed viability likewise determining the rate of water uptake and utilization for seedling establishment and dry matter accumulation. Meanwhile seasons also determined anthesis-silking interval of the maize plant depending on the availability of water during the flowering period which could later contribute to yield. Interaction of the season by maturity group indicated that the season determined the different performance of maturity groups of maize. Likewise differential efficiency of water usage of the maize maturity groups for the establishment of the plants at the seedling and grain filling growth stages can also be attributed to the differential characteristics of the two growing seasons. This water use efficiency of the plants entails the consumption of little amount of soil and atmospheric water in moving the assimilates from the source to the sink, dry matter accumulation for different maturity groups on daily basis and final yield during the different seasons. Significant maturity group difference for the water use efficiency traits indicated that the level of water absorption and utilization differed at seedling stage and at grain filling stage for the

different maturity groups. Likewise, significance due to maturity group, on yield, indicated the extra-early and late maturity groups produced the highest yield in the early season while in the late season the early and late maturity groups had the highest yields. This supported the findings of Oluwaranti et al. [23] that different maturity groups of maize have different quality that makes them acceptable as a variety of maize. It was also observed that the maize genotypes use water efficiently during the early cropping season at the seedling stage but not at the grain filling stage which also affected the yield. This was because there was poor distribution of rainfall in the month of July 2016 thereby increases the atmospheric temperature for weeks which in turn drastically reduced the soil moisture content during the grain filling period. This was also reported to have been the hottest month since weather record begins and this goes in support of report by NeSmith and Ritchie [31] who observed that maize yield can be reduced by as much as 90% if drought stress occurs between a few days before tassel emergence and the beginning of grain filling. Meanwhile for grain filling water use efficiency and for dry matter accumulation on daily basis on seedling and grain filling growth rate, the late cropping season performed better. The failure of dry matter accumulation during the early cropping season can be attributed to infestation of fall army worms which started right from five days after planting till maturity through perforation of the leaves which is supposed to be used in manufacturing food to fill the grains. The voracious consumption on leaves brings changes in the vegetative, yield and yield components while the efficient use of water during the early cropping season at seedling stage can be attributed to improved soil moisture and reduced temperature. It was also observed that the extra early and early maturity groups of maize use water most efficiently during the early cropping season at the seedling stage indicating the extra early and early maturity group which are bred for drought escape are capable of utilizing water obtained from the soil for stand establishment and dry matter accumulation which can in turn contribute to yield. This corroborates the findings of Ajani et al. [29] that water use efficiency of rain-fed maize is important for identifying maize cultivars that are efficient in the use of limited soil water for biomass and grain yield production. During the late cropping season, the late and intermediate maturity groups of maize used water more efficiently compared to extra early and early maturity groups in the early cropping season

indicating the differences in performance of the maturity groups in their ability to adapt well and utilize the little amount of water in the soil for dry matter accumulation. The performance of the maturity groups also differs in terms of flowering, yield, and yield components as also reported by [23,32,33].

Positive relationship of dry weight and water use efficiency at the grain filling stage with yield for all the maturity groups except the extra-early maturity group during the early cropping season of this study can be attributed to the longer duration of the early cropping season which allowed availability of more efficient growing conditions for their development as reported by Sivritepe et al. [26]. This was contrary to what was obtained during the late season which was characterized by short duration of the growing season which did not allowed the maturity groups to perform maximally.

5. CONCLUSION

Maturity group of maize played a significant role in the expression and manifestation of water use efficiency traits under the different cropping seasons of the rainforest agro-ecology at seedling and grain filling growth stages. The different maturity groups except the extra-early maturity group used water more efficiently during the early season than the late season of this study. High dry weight and water use efficiency can be used as indirect selection for high yield for all the maturity groups except the extra-early maturity group during the early season.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. FAO STAT. Country profile. United Republic of Tanzania; 2015. [Accessed on March 3] Available: <http://faostat3.fao.org/home/E>
2. Amani, HKR Agricultural Development and Food Security in Sub-Saharan Africa Tanzania Country Report. Economic and Social Research Foundation (ESRF) Dares Salaam, Tanzania.
3. Zorya S, Morgan N, Rios LD. Missing food: The case of postharvest grain losses in Sub-Saharan Africa. The International Bank for Reconstruction and Development

- / The World Bank. Report No. 60371-AFR: The World Bank, Washington, DC; 2011.
4. BEFS, (Bioenergy and Food Security Projects). Tanzania. BEFS Country Brief. [Accessed on November 6 2014] Available:www.fao.org/bioenergy/foodsecurity/befs
 5. Bekric V, Radosavljevic M. Savremeni-pristupiupotrebekukuruza. PTEP. 2008;12: 93-96.
 6. Alahdadi I, Oraki, H, Parhizkarkhajani F. Effect of water stress on yield and yield components of sunflower hybrids. Afr J. Biotechnol. 2011;10(34):6504-6509.
 7. Khrodarahmpour Z. Effect of drought stress induced by polyethylene glycol (PEG) on germination indices in corn (*Zea mays* L.) hybrids. African Journal of Biotechnology. 2011;10:18222-18227.
 8. Pingali PL, Pandey S. Meeting world maize needs: Technological opportunities and priorities for public sector. In: Pingali PL. (ed.). CIMMYT 1999-2000. World Maize Facts and Trends. CIMMYT, Mexico. 2001; 1-20.
 9. International Institute for Tropical Agriculture, (IITA). Annual Report for 2005, Ibadan, Nigeria: International Institute for Tropical Agriculture; 2009.
 10. Jensen JR. Remote sensing of the environment: An earth resource perspective. 2nd Edition, Pearson Prentice Hall, Upper Saddle River; 2007.
 11. Bacon MA. Water use efficiency in plant biology; 2004 (CRC Press).
 12. Tijani FO, Oyedele DJ, Aina PO. Soil moisture storage and water-use efficiency of maize planted in succession to different fallow treatments. International Agro-physics. 2008;22:81-87.
 13. Adamtey N, Cofie O, Ofosu-Budu KG, Ofosu-Anim J, Laryea KB, Forester D. Effect of N enriched co-compost on transpiration efficiency and water-use efficiency of maize (*Zea mays* L.) under controlled irrigation. Agricultural Water Management 2010;97:995-1005. DOI: 10.1016/j.agwat.2010.02.004
 14. Frimpong JO, Amoatey HM, Ayeh EO, Asare DK. Productivity and soil water use byrainfed maize genotypes in a coastal savannah environment. International Agrophysics. 2011;25:123-129.
 15. Irmak S, Haman DZ, Basting R. Determination of crop water stress index for irrigation timing and yield estimation of corn. Agronomy Journal. 2000;92:1221-1227. DOI: 10.2134/agronj2000.9261221x
 16. Yazar A, Sezen SM, Gencel B. Drip irrigation of corn in the southeast Anatolia Project(GAP) area in Turkey. Irrigation Drainage Journal. 2002;51:293-300. DOI: 10.1002/ird.63
 17. Istanbuloglu A, Kocaman I, Konuku F. Water production relationship of maize under Tekirday conditions in Turkey. Pakistan Journal of Biological Science. 2002;5:287-291. DOI: 10.3923/pjbs.2002.287.291.
 18. Oktem A, Simsek M, Okem AG. Deficit irrigation effects on sweet corn with drip irrigationsystem in a semi-arid region. I. Water-yield relationship. Agricultural Water Management. 2003;61:63-74. DOI: 10.1016/S0378-3774(02)00161-0
 19. Çakir R. Effect of water stress at different development stages on vegetative and reproductive growth of corn. Field Crops Research. 2004;89:1-16.
 20. Huang R, Birch CJ, Gorge DC. Water use efficiency in maize production – The challenge and improvement strategies, Maize Association of Australia, Australia; 2006.
 21. Heisey PW, Edmeades GO. Maize production in drought-stressed environments: Technical options and research resource allocation. Part 1 of CIMMYT 1997/1998 World Facts and Trends; 1999.
 22. Grant FR, Jackson BS, Kiniry JR, Arkin GF. Water deficit timing effects on yield components in maize. Agronomy Journal. 1989;81:61-65.
 23. Oluwaranti A, Fakorede MAB, Badu-Apraku B. Grain yield of maize varieties of different maturity groups under marginal rainfall conditions. Journal of Agricultural Sciences. 2008;53(3).
 24. Shaibu AS, Rabi IU, Adnan AA. Variability of root and physiological traits of different maturity groups of maize (*Zea mays* L.). J. Plant Breed. Crop Sci. 2015;7(7):233-239.
 25. Vina A, Giitelson AA, Rundquist DC, Keydan G, Leavitt B, Schepers J. Monitoring maize (*Zea mays* L.) phenology with remote sensing. Agron. J. 2004;96: 1139-1147.
 26. Sivritepe HO, Sivritepe N, Senturk B. Correlation between viability and different vigour tests of maize seeds. International

- Journal of Agriculture and Environmental Research. 2016;02(06):1891–1898.
27. Ajayi SA, Fakorede MAB. Effect of storage environments and duration of equilibration on maize seed testing and seedling evaluation. *Maydica*. 2001;46:267-275.
 28. Bewley JD, Black M. *Seeds: Physiology, development and germination*. 2nd Edn. Plenum Press, New York; 1998.
 29. Ajani OT, Oluwaranti, A. Awoniyi AI. Assessment of water-use efficiency of drought tolerant maize varieties in a rain forest location. *Journal of Agriculture and Ecology Research International*. 2016;8(3): 1 -10.
 30. Misra AN. Water use efficiency of pearl millet (*Pennisetum americanum* L. Leeke) genotypes under mid-season moisture stress. *Acta Agronomica Hungarica*. 1991; 40:417-422.
 31. NeSmith DS, Ritchie JT. Effects of soil water-deficits during tassel emergence on development and yield components of maize (*Zea mays* L). *Field Crops Res*. 1992;28:251-256.
 32. Oluwaranti A, Fakorede MAB, Abebe M, Badu-Apraku B. Climatic conditions requirements of maize germplasm for flowering in the rainforest Agro-ecology of Nigeria. *J. Plant Breed. Crop Sci*. 2015; 7(6):170-176.
 33. Akinwale RO, Awosanmi FE, Olufunmike OO, Fadoju AO. Determinants of drought tolerance at seedling stage in early and extra – Early maize hybrids. *Maydica*. 2017;62.

© 2019 Bankole et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/50836>