



Influence of Various Organic Amendments on Growth and Yield Attributes of Mung Bean (*Vigna radiata* L.)

Smita Kumari ^a, Rajneesh Kumar ^{b*}, Shambhu Chouhan ^a
and Pravin L. Chaudhary ^b

^a Department of Agronomy, School of Agriculture, Lovely Professional University, Phagwara, Punjab-144402, India.

^b Department of Genetics and Plant Breeding, School of Agriculture, Lovely Professional University, Phagwara, Punjab-144402, India.

Authors' contributions

This work was carried out in collaboration among all authors. Author SK designed the study, wrote the protocol and first draft of the manuscript. Author RK performed the statistical analysis and managed the analyses of the study. Author SC managed the literature searches and author PLC cited the references and proofread the article. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i122975

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/99539>

Original Research Article

Received: 15/02/2023

Accepted: 01/05/2023

Published: 13/05/2023

ABSTRACT

A field experiment was conducted during summer season of 2022 to study the influence of various organic amendments on various attributes of growth and yield of mung bean. The experiment was laid out in four replications with 11 treatments in Randomized Block design (RBD). The results revealed that the highest plant height (52.8 cm), number of leaves plant⁻¹ (35.3), number of branches plant⁻¹ (11.1), chlorophyll content using SPAD (58.4), leaf area index (1.85) was recorded from the combined application of T11= Recommended dose of fertilizers(RDF) + Vermicompost +

*Corresponding author: E-mail: r4rajneesh1997@gmail.com;

Jeevamrutam + Rhizobium. Similarly for yield attributes the highest number of pods plant⁻¹ (31.35), pod's length (11.05 cm), number of grains pods⁻¹ (12.2), test weight (45.7 g), grain yield (1242.5 kg ha⁻¹), straw yield (2181.1 kg ha⁻¹), biological yield (3405.3 kg ha⁻¹) and harvest index (36%) was recorded in combined application of same treatment T11. The result showed that combined application of organic manures had positive effects on growth and yield parameters of mung bean.

Keywords: RDF; Vermicompost; Jeevamrutam; Rhizobium and RBD.

1. INTRODUCTION

Mung bean, a member of the Fabaceae family, is an ancient pulse crop that originated from Southeast Asia and has been widely cultivated for human consumption [1]. It can be consumed in various forms including boiled dry beans, stew, flour, sprouts and immature pods as a vegetable. Additionally, the roasted or boiled dry beans can be used for animal food, especially poultry, while its biomass can serve as fodder [2]. Overall, mung bean is a valuable source of cheap protein for human and animal consumption [3].

India is the leading producer of mung bean globally and ranks third among important pulse crops grown in India. It covers around 16% of the total pulse area of the country. The crop is cultivated in about 4.5 million hectares with a total production of 2.5 million tonnes, which contributes around 10% to the total pulse production. The third advance estimates by the Government of India in 2020-21 suggest a mung bean production of 2.64 million tonnes (India stat mung bean outlook report, 2021). However, despite the significance of pulses for protein requirements, the availability of pulses in India is only around 40 grams per capita per day, which falls far below under daily intake recommendation of 85 grams of pulses per capita (FAO/WHO recommendations).

Organic amendments refer to natural products that are derived from organic sources such as animal waste, vegetable compost, agricultural residues, and human excreta. Rhizobium, vermicompost, and liquid formulations of jeevamrutam are biological products mix with the soil, providing plant nutrients and increasing soil fertility, organic matter, microbial activity, and aggregate stability. Rhizobium, in particular known for its ability to enhance biological fixation of atmospheric nitrogen into ammonia and improve phosphorus availability to crops [4-5].

Vermicompost is an organic amendment that is rich in nutrients, containing high levels of humus, 3% nitrogen, 1% P₂O₅, 1.5% K₂O, micronutrients, and beneficial soil microbes like nitrogen-fixing bacteria and mycorrhizal fungi. Scientific research has proven that vermicompost is an effective enhancer of plant growth [6]. Vermicomposting involves the biological degradation of organic waste by earthworms to form vermicast by Edwards and Burrows [7]. Studies by Ansari and Ismail [8] have shown that vermicast produced by worms contains 7.37% nitrogen and 19.58% P₂O₅.

Jeevamrutam is a natural fertilizer made from cow dung, cow urine, pulse flour, jaggery, and a small amount of soil to increase the microbial population. It is a highly effective plant growth stimulant that enhances the biological efficiency of crops. It is used to activate soil and protect plants from diseases and increases the nutritional quality of fruits and vegetables by Devakumar et al. [9]. Cow urine is an excellent source of plant nutrients and has anti-fungal properties. Objective of this experiment to check the influence of various of organic amendments on growth and yield of mung bean.

2. MATERIALS AND METHODS

The field experiment was conducted at Agriculture research farm of Lovely Professional University, Phagwara, Punjab during *Spring* season of 2022. The experiment was conducted using a randomized block design (RBD) with four replications and each plot measurement covered an area of 3m x 5m. The variety SML-668 was used for sowing green gram. Treatment considering three level of organic manures *viz.* Vermicompost, Rhizobium and Jeevamrutam. in the experiment, which consisted of 11 treatments labelled as T1- control, T2- Rhizobium, T3- Jeevamrutam, T4- Vermicompost, T5- Vermicompost + Rhizobium + Jeevamrutam, T6- Recommended Dose of Fertilizer or RDF, T7- RDF + Rhizobium, T8- RDF + Rhizobium + Jeevamrutam, T9- RDF + Vermicompost, T10-

RDF + Vermicompost + Rhizobium and T11-RDF + Vermicompost + Rhizobium + Jeevamrutam. Please specify the soil type and its properties where field experiment conducted. Variety is SML-668, seed rate 38 kg/ha, 5 Irrigation given for proper plant growth. Plant population is 6349.21 plants ha⁻¹. Soil type sandy loam, pH – 8.60, EC- 0.19, Organic Carbon 0.32% Urea – 27.5 kg ha⁻¹, SSP- 250 Kg and potassium is available in sufficient amount. Jeevamrutam – 500 l ha⁻¹ in 3 doses. Vermicompost - @ 2 tons ha⁻¹ seed was treated with rhizobium culture (*Rhizobium leguminosarum*). First selected healthy leaves, calibrate the SPAD meter, place the leaf on the meter and take a reading, Recorded the SPAD reading, repeat on several leaves and average the readings, higher readings mean more chlorophyll.

The growth attributes viz. plant height (cm), number of leaves plant⁻¹ and number of branches recorded at harvest whereas Leaf area index and Chlorophyll index at 55 DAS. Yield parameters viz. number of pods plant⁻¹, Pod length (cm), Number of grains pod⁻¹ recorded at harvest whereas test weight (g), grain yield, Straw yield, Biological yield (kg ha⁻¹) and Harvest index (%) recorded after threshing.

How you estimated chlorophyll content

2.1 Formula Used

Where,

Leaf area index was recorded in cm by formula:

$$\frac{\text{Leaf area per plant}}{\text{Ground area}} \times 100$$

Harvest index was recorded by in % by formula:

$$\frac{\text{Economic yield}}{\text{Biological yield}} \times 100$$

2.2 Statistical Analysis

LSD test was performed to determines the impact of different treatments s used in the experiment, statistical analysis of data was performed using the OPSTAT software with a 95% confidence level. This means that the obtained results are considered statistically significant, with a probability of less than 5% that they are due to chance.

3. RESULTS AND DISCUSSION

3.1 Growth Attributes

The data presented in Table 1 and Fig. 1 revealed crop growth and development measured in the terms of plant height (cm), number of leaves plant⁻¹, number of branches plant⁻¹, chlorophyll content using SPAD and Leaf area index (%).

Table 1. Influence of organic amendments on growth attributes of mung bean

Treatment		Growth attributes				
		Plant height (cm)	No of leaves plant ⁻¹	No of Branches Plant ⁻¹	Chlorophyll content	Leaf Area Index
T _c	T1	37.5	21.5	6.5	44.1	1.22
T _R	T2	40.0	26.6	8.2	50.1	1.48
T _J	T3	38.7	22.4	6.8	46.9	1.33
T _V	T4	39.8	25.7	7.9	49.7	1.37
T _{V+R+J}	T5	42.7	29.6	9.2	50.5	1.50
T _{RDF}	T6	44.5	30.8	9.6	54.8	1.54
T _{RDF+R}	T7	47.2	31.7	9.9	55.4	1.63
T _{RDF+R+J}	T8	49.5	32.3	10.1	56.0	1.69
T _{RDF+V}	T9	46.4	30.5	9.5	51.9	1.58
T _{RDF+V+R}	T10	49.9	34.1	10.7	57.1	1.76
T _{RDF+V+R+J}	T11	52.8	35.3	11.1	58.4	1.85
LSD (0.05)		3.52	2.15	0.90	3.19	0.13

T_c = Control, T_R = Rhizobium, T_J = Jeevamrutam, T_V = Vermicompost, T_{V+R+J} = Vermicompost + Rhizobium + Jeevamrutam, T_{RDF} = Recommended dose of fertilizers, T_{RDF+R} = Recommended dose of fertilizers + Rhizobium, T_{RDF+R+J} = Recommended dose of fertilizers + Rhizobium + Jeevamrutam, T_{RDF+V} = Recommended dose of fertilizers + Vermicompost, T_{RDF+V+R} = Recommended dose of fertilizers + Vermicompost + Rhizobium, T_{RDF+V+R+J} = Recommended dose of fertilizers + Vermicompost + Rhizobium + Jeevamrutam




Fig. 1. Influence of organic amendments on growth attributes of mung bean

3.1.1 Plant height, number of leaves plant⁻¹, number of branches plant⁻¹, chlorophyll content and leaf area index

The data recorded at harvest revealed that the plant height in the treatment T11 (52.8) that was significantly higher than T5, T6, T7 and T9 whereas T8 and T10 were statistically similar with T11. The findings reported are in agreement with the results obtained by [10-13]. For number of leaves plant⁻¹ in the treatment T2, T4, T5, T6, T7, T8, T9, T10 and T11 were significantly higher than control. The highest number of leaves plant⁻¹ was observed in the treatment T11 (35.3) that was significantly higher than T5, T6, T7, T8 and T9 whereas T10 was statistically similar with T11. Similar findings reported were in consistent with the results obtained by [11-13]. Similarly, number of branches plant⁻¹ in the treatment T2, T4, T5, T6, T7, T8, T9, T10 and T11 were significantly higher than control. The highest number of branches plant⁻¹ was observed in the treatment T11 (11.1) that was significantly higher than T5, T6, T7 and T9 whereas T10 and T8 was statistically similar with T11. [11-13 and 14] also reported similar results. However, at 55 DAS the chlorophyll content in the treatment T2, T4, T5, T6, T7, T8, T9, T10 and T11 were significantly higher than control. The highest chlorophyll content using SPAD was observed in T11 (58.4) that was significantly higher than T5 and T9 whereas T6, T7, T8 and T10 were statistically similar with T11. [12] also reported similar results. For leaf area index at 55 DAS revealed that treatment T2, T5, T6, T7, T8, T9, T10 and T11 were significantly higher than control. The highest Leaf Area Index was observed in T11 (1.85) that was significantly higher than T5, T6, T7, T8 and T9 whereas T10 were statistically

similar with T11. [12-13] also reported similar results.

3.2 Yield attributes

The data presented in Table 2 and Fig. 2. yield and yield attributes on mung bean was measured in the terms of number of pods plant⁻¹, pods length (cm), number of grains pod⁻¹, test weight (g), grain yield (kg ha⁻¹), straw yield (kg ha⁻¹), biological yield (kg ha⁻¹) and harvest index (%).

3.2.1 Number of pods plant⁻¹, pods length, number of grains pod⁻¹, test weight, grain yield, straw yield, biological yield and harvest index

The data recorded at harvest revealed that the highest number of pods plant⁻¹ of mung bean was recorded by treatment T11 (31.35) and the lowest in T1 (19.10). Treatment T2, T3, T4, T5, T6, T7, T8, T9 and T10 were significantly higher than T1. T2 was significantly higher than T3 and T4 whereas T5 was statistically similar with T6. T11 was significantly higher than T7, T8 and T9 while T11 was statistically similar with T10. Similar findings reported were consistent with the results obtained by [14-15]. Highest pods length of mung bean was recorded by treatment T11 (11.05) whereas T4, T5, T6, T7, T8, T9 and T10 were significantly higher than control. The pods length in T5 was significantly higher than T3 and T4 but statistically similar with T2. T11 was significantly higher than T6, T7, T8, and T9 while T11 was statistically similar with T10. The outcomes reported are consistent with the results obtained by [16-17]. For number of grains pod⁻¹ of mung bean was recorded by treatment T11 (12.2) whereas T2, T4, T5, T6, T7, T8, T9 and

T10. T5 was significantly higher than T3 and T4 but statistically similar with T2. [14-15 and 17] also reported similar results. The data recorded after threshing revealed that the highest test weight of mung bean was recorded by treatment T11 (45.7). T2, T5, T6, T7, T8, T9 and T10 were significantly higher than T1 however, test weight in T5 was significantly higher than T2, T3 and T4. The test weight in T11 was significantly higher than T6, T7 and T9 while T11 was statistically similar with T8 and T10. [14 and 16] also reported similar results. The highest grain yield

(kg ha⁻¹) of mung bean was recorded by treatment T11 (1242.5). T2, T4, T5, T6, T7, T8, T9 and T10 were significantly higher than T1. However, grain yield in T5 was significantly higher than T2, T3 and T4. The grain yield in T11 was significantly higher than T6, T7, T8, T9 and T10. The outcomes reported are consistent with the results obtained by [14-15,17-18]. Similarly, the highest straw yield (kg ha⁻¹) of mung bean was recorded by treatment T11 (2181.1). T2, T5, T6, T7, T8, T9 and T10 were significantly higher than T1.

Table 2. Influence of organic amendments on yield attributes of mung bean

Treatment	Yield attributes								
	No of pods plant ⁻¹	Pods length (cm)	No of grains Pod ⁻¹	Test weight (g)	Grain yield (kg ha ⁻¹)	Straw yield (kg ha ⁻¹)	Biological yield (kg ha ⁻¹)	Harvest index (%)	
T _c	T1	19.10	4.73	7.5	32.4	748.5	1612.5	2397.0	32.7
T _R	T2	27.00	7.33	10.3	36.3	855.5	1750.9	2606.8	32.8
T _J	T3	24.00	5.03	7.8	34.6	778.0	1654.0	2432.3	31.9
T _V	T4	25.20	6.40	8.4	35.8	816.5	1688.0	2504.8	32.6
T _{V+R+J}	T5	27.85	8.70	10.5	40.8	926.3	1815.5	2741.3	33.7
T _{RDF}	T6	26.80	8.38	9.5	39.1	911.5	1785.8	2696.8	33.8
T _{RDF+R}	T7	29.20	9.50	11.2	42.2	952.3	1871.5	2823.5	34.0
T _{RDF+R+J}	T8	29.85	9.80	11.6	42.8	995.3	1908.0	2903.0	34.3
T _{RDF+V}	T9	28.05	9.23	10.7	41.5	883.6	1838.8	2767.3	32.0
T _{RDF+V+R}	T10	30.40	10.28	11.8	43.5	1127.3	2086.3	3212.5	35.0
T _{RDF+V+R+J}	T11	31.35	11.05	12.2	45.7	1242.5	2181.1	3405.3	36.0
LSD (0.05)		1.67	1.19	0.67	3.00	38.52	128.07	31.64	0.90

T_c = Control, T_R = Rhizobium, T_J = Jeevamrutam, T_V = Vermicompost, T_{V+R+J} = Vermicompost + Rhizobium + Jeevamrutam, T_{RDF} = Recommended dose of fertilizers, T_{RDF+R} = Recommended dose of fertilizers + Rhizobium, T_{RDF+R+J} = Recommended dose of fertilizers + Rhizobium + Jeevamrutam, T_{RDF+V} = Recommended dose of fertilizers + Vermicompost, T_{RDF+V+R} = Recommended dose of fertilizers + Vermicompost + Rhizobium, T_{RDF+V+R+J} = Recommended dose of fertilizers + Vermicompost + Rhizobium + Jeevamrutam




Fig. 2. Influence of organic amendments on yield attributes of mung bean

However, straw yield in T5 was significantly higher than T3 and T4 while statistically similar with T2. The straw yield in T11 was significantly higher than T6, T7, T8 and T9 and T11 was statistically similar with T10. These results are in accordance with [15 and 17]. Moreover, biological yield (kg ha^{-1}) of mung bean was recorded by treatment T11(3405.3). T2, T3, T4, T5, T6, T7, T8, T9 and T10 were significantly higher than T1. However, biological yield in T5 was significantly higher than T2, T3 and T4. The biological yield in T11 was significantly higher than T6, T7, T8, T9, and T10. [16] also reported similar results. The harvest index (%) is the proportionality of grain yield to the biological yield multiplied by 100. The data recorded after threshing revealed that the highest harvest index of mung bean was recorded by treatment T11 (36). T2, T4 and T9 were statistically similar with T1 while treatment T11 was significantly higher than T7, T8 and T10. However, harvest index in T5 was statistically similar with T6. The biological yield in T11 was significantly higher than T6, T7, T8, T9, and T10. [16 and 18] also reported similar results.

4. CONCLUSION

From the above results it can be concluded that the combination of organic and inorganic fertilizers is more productive compared to sole application of organic as well as inorganic fertilizers. By combining both organic and inorganic fertilizers we got highest growth and yield parameters. The combination of organic and inorganic fertilizers have positive effects on the growth and yield parameters of mung bean. From my research I concluded that the highest plant height (52.8 cm), number of leaves plant^{-1} (35.3), number of branches plant^{-1} (11.1), chlorophyll content using SPAD (58.4), leaf area index (1.85) was recorded from combined application of RDF + vermicompost + jeevamrutam + Rhizobium similarly for yield parameters the highest number of pods plant^{-1} (31.35), pods length (11.05 cm), number of grains pods^{-1} (12.2), test weight (45.7 g), grain yield ($1242.5 \text{ kg ha}^{-1}$), straw yield ($2181.1 \text{ kg ha}^{-1}$), biological yield ($3405.3 \text{ kg ha}^{-1}$) and harvest index (36%) was recorded in combined application of RDF + vermicompost + jeevamrutam + Rhizobium. Therefore, I recommended farmers to use combine application of organic and inorganic fertilizers instead of using their sole application.

ACKNOWLEDGEMENTS

I would like to express my gratitude to Dr. Harmit Singh Thind, Department of Agronomy, School of Agriculture at LPU, for being my advisor and providing continuous support and guidance throughout my research study. I also thank my co-author for their valuable contributions. Special thanks to Rajneesh Kumar for his assistance with data analysis and formatting of the article.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mogotsi K. Effect of jeevamrut on growth, yield and quality of mung bean (*Vigna radiata* L.). Ph.D. Thesis, Maharana Pratap University of Agriculture and Technology., Udaipur, Rajasthan, India; 2006.
2. Winch T. Description and characteristics of the main food crops. In Growing Food a Guide to Food Production. Springer., Dordrecht. 2006;18-22.
3. Oplinger ES, Hardman LL, Kaminski AR. Varietal response to different levels of fertility and bio-fertilizer on yield and quality of summer mung bean. Indian Journal of Agronomy. 1990;47:473-476.
4. Bhat TA, Gupta M, Ganai MA, Ahangar RA, Bhat HA. Yield, soil health and nutrient utilization of field pea (*Pisum sativum* L.) as affected by phosphorus and biofertilizers. International Journal of Modern Plant and Animal science. 2013;1:1-8.
5. Kumar A, Maurya BB, Raghuwanshi R. Effects of rhizobium on the growth, yield and soil nutrient status in mung bean. Journal of Science and Technology. 2014;12:23-28.
6. Chaoui HI, Zibilske LM, Ohno T. Effects of earthworm casts and compost on soil microbial activity and plant nutrient availability. Soil Biology and Biochemistry. 2003;35(2):295-302.
7. Edwards CA, Burrows I. The potential of earthworms composts as plant growth media. Simon Peter Bakker Academic Publication. 1988;34:211–220.
8. Ansari AA, Ismail SA. Role of earthworms in vermitechnology. Journal of Agricultural Technology. 2012;8(2):403-415.

9. Devakumar N, Shubha S, Gowder SB, Rao GG. Microbial analytical studies of traditional organic preparations beejamrutha and jeevamrutha. Building Organic Bridges. 2014;2:639-642.
10. Balachandran D, Nagarajan P. Dual inoculation of Rhizobium and phosphobacteria with phosphorus on black gram Madras Agriculture Journal. 2002;89:691693.
11. M. Srinivas and Shaik M. Effect of Rhizobium inoculation and different levels of nitrogen and phosphorus on yield and economics of greengram (*Vigna radiata*. L. Wil.). Crop Research. 2012;24:463-466.
12. Singh B, Pareek RG. Studies on phosphorus and bio-inoculants on biological nitrogen fixation, concentration, uptake, quality and productivity of mungbean. Annals of Agricultural Research. 2003;24(3):537-41.
13. Rajesh M, Jayakumar K, Kannan TM. Effect of biofertilizers application on growth and yield parameters of green gram (*Vigna radiata* L.). International journal of environment and bioenergy. 2013;7(1): 43-53.
14. Verma R, Singh M, Dawson J, Muddassir P, Khan I. Effect of bio-fertilizers and organic manures on growth and yield of greengram (*Vigna radiata* L.). The Pharma Innovation Journal. 2022;11(4): 1599-602.
15. Marko GS, Kushwaha HS, Singh S, Namdeo KN, Sharma RD. Effect of sulphur and biofertilizers on growth, yield and quality of blackgram (*Phaseolus mungo* L.). Crop Research. 2013;45:175-178.
16. Dongare DM, Pawar GR, Murumkar SB, Chavan DA. To study the effect of different fertilizer and biofertilizer levels on growth and yield of summer greengram. International Journal of Agricultural Sciences. 2016;12(2):151-157.
17. Mohammad I, Yadav BL, Ahamad A. Effect of phosphorus and bio-organics on yield and soil fertility status of mungbean (*Vigna radiata* L.) Wilczek] under semi-arid condition of Rajasthan, India. International Journal of Current Microbiology and Applied Sciences. 2017; 6:1545-1553.
18. Dhakal Y, Meena RS, Kumar S. Effect of INM on nodulation, yield, quality and available nutrient status in soil after harvest of greengram. Legume Research- An International Journal. 2016;39(4): 590-594.

© 2023 Kumari et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/99539>