



# Effect of Preparation, Microwave and Pressing on Quality of Carrot Powder

Dev Raj <sup>a\*</sup>, Ashuqullah Atif <sup>a</sup>, D. R. Bhandari <sup>b</sup>,  
A.K. Senapati <sup>a</sup> and A.K. Pandey <sup>c</sup>

<sup>a</sup> Department of Post Harvest Technology, ASPEE College of Horticulture, Navsari Agricultural University, Navsari, Gujarat, India.

<sup>b</sup> Department of Vegetable Science, ASPEE College of Horticulture, Navsari Agricultural University, Navsari, Gujarat, India.

<sup>c</sup> Department of Fruit Science, ASPEE College of Horticulture, Navsari Agricultural University, Navsari, Gujarat, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i91397>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/123377>

**Original Research Article**

**Received: 05/07/2024**

**Accepted: 07/09/2024**

**Published: 12/09/2024**

## ABSTRACT

The experiment was carried out with the aim to study effects of preparation, microwave and pressing on quality of carrot powder. The experiment was conducted at Department of Post Harvest Technology, ASPEE College of Horticulture, Navsari Agricultural University, Navsari, Gujarat during 2021-22. Single strength pressing was done to fasten the drying rate. The experiment was replicated three times using completely randomised design (CRD) with factorial concept and comprised of 18 treatments viz. preparation techniques as Factor-1 ["Shredding" (T<sub>1</sub>), "Crushing" (T<sub>2</sub>) and "Slicing" (T<sub>3</sub>)]; microwave pre-treatments as Factor-2 [Control (M<sub>1</sub>), Microwave pre-treatment 450 W for 30 seconds (M<sub>2</sub>) and microwave pre-treatment 450 W for 60 seconds (M<sub>3</sub>)] and

\*Corresponding author: E-mail: [drpandir\\_phtc@nau.in](mailto:drpandir_phtc@nau.in);

**Cite as:** Raj, Dev, Ashuqullah Atif, D. R. Bhandari, A.K. Senapati, and A.K. Pandey. 2024. "Effect of Preparation, Microwave and Pressing on Quality of Carrot Powder". *Journal of Advances in Biology & Biotechnology* 27 (9):1265-82. <https://doi.org/10.9734/jabb/2024/v27i91397>.

pressing as Factor-3 [without pressing ( $P_1$ ) and pressing ( $P_2$ )]. After single strength pressing, the samples were dehydrated in mechanical dehydrator using three stage dehydration temperatures (75°C for 2 hours, 65°C for 2 hours and 60°C for about 11-12 hours) till final moisture content of  $5.50 \pm 1\%$  following by packing. The results of the investigation revealed that best quality carrot powder can be obtained from carrot which was prepared by dehydrating crushed materials by giving microwave for 60 seconds followed by pressing ( $T_2M_3P_2$ ) and carrot powder obtained from this treatment possess higher rehydration ratio, dehydration ratio and sensory score while lowest drying time, moisture content, water activity, reducing sugars and non-enzymatic browning (NEB). While, the carrot powder which was prepared by dehydrating crushed materials by giving microwave pre-treatment for 60 seconds followed by pressing ( $T_2M_3P_2$ ) exhibited minimum changes in nutritional as well as sensory attributes during six months' storage when packed in polypropylene bags of 400-gauge thickness. It can be concluded that carrot powder can be prepared by dehydrating crushed materials by giving microwave pre-treatment for 60 seconds followed by pressing ( $T_2M_3P_2$ ) and the powder can remain microbiologically safe for storage period of six months.

**Keywords:** Carrot; powder; dehydration; quality; pre-treatments.

## 1. INTRODUCTION

“Among major vegetables, carrot (*Daucus carota* L.) belonging to Apiaceae family is the most important root crop grown throughout the world. It is a root vegetable, usually orange, purple, red, white or yellow, with a crisp texture when fresh. It is a rich source of  $\beta$ -carotene and contains other vitamins, like thiamine, riboflavin, vitamin B-complex and minerals” [1]. “The carrot is consumed in different forms as raw, juice, salads, cooked vegetable sweet dishes etc. Carrots provide great health benefits to the human body as they are good sources of carotenoids, bioactive compounds, vitamins and minerals” [2]. “Bioactive compounds present in the carrots are  $\alpha$ - and  $\beta$ -carotene, phenolic acids and lycopene which are responsible for their antioxidant activity. Carrot contains approximately 86.00 per cent moisture, 10.60 g carbohydrates, 0.20 g fat, 1.20 g fiber, 1890  $\mu$ g carotene, 0.04 g thiamine, 0.02 g riboflavin, 0.60 g niacin, 530.00 mg phosphorus, 1.10 g minerals, 3.00 mg vit C and 26.60  $\mu$ g total phenolic content per 100 g” [3]. “Due to its good flavor and nutrition, carrot is regarded as a very important vegetable owing to its various properties like anticancer, anti-anemic, antioxidant, sedative and healing which are directly related to human health” [4]. “Due to the perishable and seasonal nature of carrots, it is not possible to make fresh carrots available throughout the year. Therefore, consumer demand has increased toward processed foods that have more natural flavour and colour with high nutritional quality and sufficient storage safe for distribution and consumption. This can be achieved by adopting suitable processing methods which inactivate micro-organisms as

well as enzymes with little loss of pigments, flavor compounds, and vitamins” [5].

“Dehydrated carrots in the form of grating can be used in the preparation of *halwa*, discs made in to chips” [6]. “Pre-treatment improves the nutritional, sensorial and functional properties of the dehydrated food without changing its integrity. It also improves the texture as well as stability of the pigment during dehydration and the storage of dehydrated product” [7]. Carrots are dehydrated in the form of slices, shreds, crushed, cubes and strips. High solids and freedom from woody fiber are desired qualities of carrots to be dehydrated. The low rehydration ratio, discoloration, and development of off-flavour are the commonly reported problems with dehydrated carrot products. Removal of biologically active water is the main task while preserving food [8] by reducing the moisture contents to a level, that allows safe storage over an extended period of time with microbial safety. “During microwave treatment, the generation of heat within tissue could create pressure, which ruptures the cell walls and increases the mass transfer of carotene into extraction solvent. As a result, the time required for microwave-assisted extraction of  $\beta$ -carotene from carrot can be reduced” [9]. Microwave treatment also known to enhance juice recovery and decrease dehydration time. Beside, use of different preparation techniques also possess effect on yield and nutritional quality. Very limited literature is available on the effect of preparation techniques, microwave pre-treatments and preservatives on quality of carrot powder during storage. Thus, the present investigation was carried out to study the effect of preparation,

microwave and pressing on the quality of carrot powder.

## 2. MATERIALS AND METHODS

**Raw materials:** Fresh matured but tender, disease, and injury-free, orange-colored carrots (*var.* Chantney) selected and purchased from Morarji yard, APMC, Navsari. The carrots were cleaned and used for further processing into dehydrated powder.


**Location:** The experiment was conducted in the Department of Post Harvest Technology, ASPEE College of Horticulture and Forestry, Navsari Agricultural University, Navsari-396450, Gujarat (India).

**Experiment detail:** A total of 18 treatments were used for dehydration of carrot into powder using combinations of different treatments viz. preparation techniques as Factor-1 ["Shredding" (T<sub>1</sub>), "Crushing" (T<sub>2</sub>) and "Slicing" (T<sub>3</sub>)]; microwave pre-treatments as Factor-2 [Control (M<sub>1</sub>), Microwave pre-treatment 450 w for 30 seconds (M<sub>2</sub>) and microwave pre-treatment 450 w for 60 seconds (M<sub>3</sub>)] and pressing as Factor-3 [ without pressing- Control (P<sub>1</sub>) and pressing for juice extraction (P<sub>2</sub>)].

**Sample Preparation and treatments:** Good quality carrots of uniform size were selected during experimentation. Then carrots were thoroughly washed to remove any adhering dust and dirt particles. After washing, the carrots were cut from the end and top to remove undesirable portions with a stainless steel (SS) knife. Then cut carrots were shredded / crushed / slices (2-3mm) with the help of a shredder/ crusher / slicer machines (Factor 1) as per experiment. The prepared shreds, crushed materials, and slices of carrots (Factor-1) were subjected to microwave

pre-treatments (Factor-2). After microwave pre-treatments, the shreds, crushed materials, and slices were subjected to pressing (Factor-3) for single-strength pressing by hydraulic press followed by loading into a mechanical dehydrator using three stage dehydration temperatures (75°C for 2 hours, 65°C for 2 hours and 60°C for about 11-12 hours) till final moisture content of 5.50 ± 1% as reported by Raj et al. [10] for dehydration of onions. After dehydration, dried shreds, crushed materials and slices were ground into powder using a lab scale mill-grinder. The prepared powder was packed in polypropylene bags of 400-gauge thickness and stored for 6 months at ambient temperature to evaluate the nutritional as well as sensory quality of developed products at every two month intervals. The principal steps used for the dehydration of carrots into powder are illustrated in Fig. 1.

**Physico-chemical parameters:** Physico-chemical parameters of fresh carrots used for dehydration (n=10) were recorded. The powder recovery was arrived at by recording the final weight of powder obtained after drying and the total weight of raw materials and expressed as recovery per cent. The moisture was estimated by drying the weighted samples in a hot air oven at 70±2°C to a constant weight [11]. For estimation of β-carotene five-gram sample was crushed into 10-15 ml of acetone followed by addition of anhydrous sodium sulphate crystals. Supernatant was collected into beaker and procedure was repeated twice. Then sample were transferred into separating funnel. After this 10-15 ml petroleum ether was added, mixed thoroughly and make up the volume of 100 ml. Optical density was recorded at 452nm using petroleum ether as blank and contents were calculated by using the using the following formula [12].



**Fig. 1. Principal steps used for dehydration of carrot into powder**

$$\beta\text{-carotene (mg /100 g)} = \frac{\text{OD} \times 13.9 \times 10000 \times 10}{\text{Weight of sample (g)} \times 560 \times 1000} \times 100$$

**Sensory evaluation:** Sensory evaluation of carrot powder was conducted during storage to assess the consumer's acceptance for the products. The prepared samples of powder were evaluated for sensory qualities on based on overall acceptability on 9 Point Hedonic scale (9 mean Extremely liked and 1 extremely disliked) according to the method of Amerine et al. [13]. Sensory panelists (7-9 members) comprised of faculty members and PG students of Department of Post Harvest Technology, NAU, Navsari (Gujarat) were used for sensory analysis throughout the period of storage. Coded samples of products were served. Plain tap water was provided to the panelists for mouth rinsing in between the sensory evaluation.

**Statistical methodology:** Experiment was carried out in Completely Randomized Design with factorial concept (FCRD). The data were collected on physico-chemical parameters of fresh carrot and expressed as mean value. The data on physico-chemical parameters of powder were statistical analyzed following a completely randomized design as described by Panse and Shukhatme [14]. The treatment differences were tested by 'F' test of significance on the basis of null hypothesis. The critical differences (C.D.) at 5 per cent level of probability were worked out to compare treatments mean for predicting significance among treatments.

### 3. RESULTS AND DISCUSSION

#### Physico-chemical attributes of fresh carrots:

The physico-chemical characteristics of fresh carrots which were used for the preparation of juice and powder are given in Table 1. The data on physical parameters of fresh carrots revealed the mean weight, peeling, and trimming losses of the carrots in the range of 75.45 g and 7.40 per cent having the orange-color of skin/ peel and flesh. The data on chemical parameters of carrots revealed mean moisture, TSS, acidity, reducing sugars, non-reducing sugars, total sugars, beta-carotene, proteins, crude fibre, ascorbic acid in the range of 9.53 °Brix, 0.12%, 2.74%, 4.15%, 6.89%, 38.65 mg/100g, 0.95%, 1.80% and 3.91mg/100g, respectively. Our results in the present investigation for physico-chemical parameters of carrots are similar to earlier studies as reported by Gopalan et al. [3]. The chemical constituents of carrot are reported to contain 86% moisture with 9.55 °Brix TSS,

2.77% reducing sugars, 6.92% total sugars, 0.12% acidity, 10.6 g carbohydrates, 0.2 g fat, 1.2 g fiber, 3965 micro g beta-carotene and 3 mg per 100 g vitamin C [3]. Sharma et al. [15] also reported similar variations for moisture (86 to 88%), carbohydrate (6 to 10.6%), crude fiber (1.2 to 2.4%), total sugars (5.6%), beta-carotene (39.6 mg/ 100 g).

**Table 1. Physico-chemical characteristics of fresh carrot used for dehydration**

Parameters*	Mean
Weight of carrot (g)	75.45
Peeling and trimming loss (%)	7.40
Peel color, visual	Orange
Flesh color, visual	Orange
Moisture (%)	86.12
TSS(°Brix)	9.53
Acidity (%)	0.12
Total Sugars (%)	6.89
Reducing sugars (%)	2.74
Non-reducing sugars (%)	4.15
Beta-carotene (mg/100g)	38.65
Proteins (%)	0.95
Crude fibre (%)	1.80

**Quality of powder:** As per the dehydration experiment treatment details, one factor was taken to study the effect of pressing along with control (without pressing) on the quality of the dehydrated carrot powder. The factor aimed to evaluate the feasibility of pressing for juice extraction, to utilize left over carrot pomace for dehydration and to evaluate dehydrated carrot powder based on physico-chemical, sensory and microbial characteristics. Thus, the prepared dehydrated carrot powder was evaluated for physico-chemical, and sensory to study the effect of different treatments during six-month storage.

**Powder yield:** The perusal of data on the effect of different treatments on powder yield of carrot powder have been presented in Table 2.

**Effect of preparation techniques:** Among different preparation techniques (T), the powder yield of carrot powder varied significantly between 9.09 and 12.38%, with minimum powder yield (9.09%) in carrot powder which was prepared by giving crushing treatment (T<sub>2</sub>) and maximum powder yield (12.38%) in slicing (T<sub>3</sub>). The minimum powder yield of carrot which was prepared by giving crushing treatment might be attributed to higher juice recovery in carrot juice. Crushing of carrots resulted in immediate release of the moisture and its soluble compounds upon pressing due to more disintegration of the cell structure. Wilczynski et al. [16] also reported a

higher yield (71.6%) of apple juice of different varieties in screw type of juice extractor due to the crushing action of fruit tissues as compared to basket press (61.9%). Raj et al. [17] reported yield of potato flour among different potato cultivars varied from 13.23 to 20.73 per cent, with the maximum yield in Kufri Chipsona-2 followed by Kufri Chipsona-1 (20.63%) and Kufri Chandramukhi (18.53%) and minimum in Kufri Jyoti followed by Kufri Badshah (14.70%). The increase in the dehydration ratio of the flour in blanching treatment might be attributed to leaching losses of the nutrients and solids.

**Effect of microwave pre-treatment:** Data shows that among different microwave pre-treatments grand mean powder yield (M) of microwave pre-treated carrot powder varied significantly between 10.16 and 10.96%, with minimum powder yield (10.16%) in carrot powder pre-treated with microwave for 60 seconds (M<sub>3</sub>) and maximum powder yield (10.96%) in carrot powder prepared without microwave pre-treatment (M<sub>1</sub>). The differences in powder yield in carrot under different microwave pre-treatments might be due to the extraction efficacy of soluble compounds owing to variations in pre-treatment time. The exposure to microwave causes disintegration of the cell structure due to vibration of the water molecules and results in higher juice recovery upon pressing and lower powder yield because the juice itself contains soluble solids along with other micro-nutrients. Gerard and Roberts [18] reported that Fuji and McIntosh apple mashes heated to bulk temperatures of 60 °C by a 2450 MHz microwave oven at 1500 W increased juice yield when the mash was treated before pressing. Rayman and Baysal [19] also reported an increase in carrot juice yield of 9.7% (50.90 to 55.84%) juice yield due to applications of the electrical methods (electroplasmolysis + microwave). The juice yield was found to vary with the variation of voltage and time of EP application. These treatments affect the cell permeability and make the juice extraction process easier. Leeratanarak et al. [20] reported that blanching helps to increase the dehydration ratio of the product which represents the decrease in the yield of the dehydrated product. The decrease in yield of the flour in blanching treatment might be attributed to leaching losses of the nutrients and solids

**Effect of pressing:** Further, it was observed that the mean powder yield (P) of the carrot when dehydrated followed by pressing treatment varied significantly from 9.19 to 11.87%, with minimum

powder yield (9.19%) in carrot powder when dehydrated followed by pressing treatment (P<sub>2</sub>) and maximum powder yield (11.87%) in carrot powder when dehydrated without pressing treatment (P<sub>1</sub>). The minimum powder yield of the carrot which was prepared for dehydration by giving pressing treatment might be attributed to the transfer of soluble solids to carrot juice during the extraction process and thus decreasing the weight of soluble solids for dehydration. Wilczynski et al. [16] also reported a higher yield (71.6%) of apple juice of different varieties due to higher pressure in screw-type juice extractor as compared to basket press (61.9%) leaving less pomace for dehydration. Leeratanarak et al. [20] reported that blanching helps to increase the dehydration ratio of the product which represents the decrease in the yield of the dehydrated product. The decrease in yield of the flour in blanching treatment might be attributed to leaching losses of the nutrients and solids.

**Effect of treatment interactions:** The interaction of microwave and pressing treatments (MP) depicted variation in carrot powder yield from 8.64 to 12.16%, with a minimum powder yield of 8.64% in powder which was dehydrated by pressing the raw material followed by pre-treatment with microwave for 60 seconds (M<sub>3</sub>P<sub>2</sub>) and maximum powder yield (12.16%) in powder which was dehydrated without pressing and microwave pre-treatment (M<sub>1</sub>P<sub>1</sub>). Interaction of preparation techniques and pressing technique (TP) depicted variation in carrot powder yield from 6.72 to 12.53%, with a minimum powder yield of 6.72% in powder which was prepared by crushing technique (T<sub>2</sub>P<sub>2</sub>) and maximum powder yield 12.53% in powder which was dehydrated by without pressing and microwave pre-treatment (T<sub>3</sub>P<sub>1</sub>). Interaction of preparation techniques, microwave and pressing techniques (TMP) depicted variation in carrot powder yield from 6.20 to 12.92%, with a minimum powder yield of 6.72% powder which was prepared by dehydrating crushed materials by giving microwave for 60 seconds followed by pressing treatment (T<sub>2</sub>M<sub>3</sub>P<sub>2</sub>) and maximum powder yield of 12.92% in powder which was prepared by dehydrating carrot slices without microwave and pressing pre-treatment (T<sub>3</sub>M<sub>1</sub>P<sub>1</sub>). Interactions of TxM, TxP, MxP, and TxMxP were found to have significant effects.

**Dehydration rate:** The perusal of data of the effect of different treatments on the dehydration rate (g/min) of carrot powder during dehydration has been presented in Table 3.

**Effect of preparation techniques:** Among different preparation techniques (T), the dehydration rate of carrot powder varied significantly between 6.89 and 8.28 (g/min), with minimum dehydration rate (6.89 g/min) in carrot powder which was prepared by giving crushing treatment (T<sub>2</sub>) and maximum dehydration rate (8.28 g/min) in slicing (T<sub>3</sub>). The minimum dehydration rate of the carrot which was prepared by giving crushing treatment might be attributed to less moisture available in the carrot pomace for drying after pressing because the pressing treatment causes higher yield of the juice and pomace remains with less moisture. Crushing of carrots followed by pressing resulted immediate release of the moisture due to more disintegration of the cell structure and thus less available moisture remained in the carrot pomace for drying. Leeratanarak et al. [6] reported that blanching time had effect on the drying rates at all drying temperature in hot air drying. The samples prepared by blanching dried faster than the unblanched samples. The increase in the drying rate of the flour in blanching treatment might be attributed to faster diffusion of the water out of the tissues. Hatamipour et al. [21] reported that an increase in blanching time decreased the drying rate. Dinrifo [22] recorded variations in the drying rate of sweet potato slices when pre-treated with blanching at 100°C and sulphitation with sodium metabisulphite. The pre-treatment reduces the resistance to moisture transport thereby increasing the drying rates. The drying of sweet potato slices exhibited higher drying rates for pre-treated samples than for untreated samples, especially at higher temperatures.

**Effect of microwave pre-treatment:** Data shows that among different microwave pre-treatments grand mean dehydration rate (M) of microwave pre-treated carrot powder varied significantly between 7.56 and 7.83 (g/min), with maximum dehydration rate (7.83 g/min) in carrot powder prepared pre-treated with microwave for 60 seconds (M<sub>3</sub>) and minimum dehydration rate (7.56 g/min) in carrot powder pre-treated with microwave for 30 seconds (M<sub>2</sub>). The differences in drying rate during carrot dehydration under different microwave pre-treatments might be due to the extraction efficacy of soluble compounds owing to variation in pre-treatment time. The exposure to microwave causes disintegration of the cell structure due to vibration of the water molecules and results from easy water removal during dehydration. Sabry et al. [23] reported that microwave treatment of carrot slices (2-3 MM)

with 360 watts for 6 min can reduce weight loss from 81.75% moisture content to 52.32% on dry weight basis with a slight reduction of drying period as compared to other treatments. Kaur and Singh [24] reported that drying of beet slices at high air temperature (75°C) followed by high microwave power (1080 W) witnessed increased drying rates and substantial shortening of the drying time. Andres et al. [25] reported that when drying of apple with a combined air microwave system, the drying rate curve could be divided into four periods limited by four critical points resulting faster drying.

**Effect of pressing:** Further, it was observed that mean Dehydration rate (P) of the carrot powder when dehydrated followed by pressing treatment varied significantly from 6.82 to 8.50 (g/min), with minimum dehydration rate (6.82 g/min) in carrot powder when dehydrated followed by pressing treatment (P<sub>2</sub>) and maximum dehydration rate (8.50 g/min) in carrot powder when dehydrated without pressing treatment (P<sub>1</sub>). The minimum dehydration rate of the carrot which was prepared by giving pressing treatment might be attributed to less moisture available in the carrot pomace for drying after pressing because pressing treatment causes higher juice extraction and pomace remained with less moisture. Pressing of prepared carrots resulted in immediate release of the moisture due to more disintegration of the cell structure.

**Effect of treatment interactions:** Interaction of preparation techniques, microwave and pressing techniques (TMP) depicted variation in carrot powder dehydration rate from 5.00 to 9.08 g/min, with minimum a dehydration rate of 5.00 g/min in powder which was prepared by dehydrating crushed materials by giving microwave for 30 seconds followed by pressing treatment (T<sub>2</sub>M<sub>2</sub>P<sub>2</sub>) and maximum dehydration rate of 9.08 g/min in powder which was prepared by dehydrating shredded materials by giving microwave for 60 seconds without pressing pre-treatment (T<sub>1</sub>M<sub>3</sub>P<sub>1</sub>). Interactions of TxM, TxP, MxP, and TxMxP were found to have significant effect. However, interactions of MxP were found to have a non-significant effects. Abano et al. [26] reported improvement in drying rates of the garlic slices followed by a drying temperature of 45 °C compared to control dried samples that were harder than all the pre-treated dried samples. Wu et al. (2013) reported higher drying rates and greater effective diffusivities under catalytic infrared (CIR) than hot air (HA) heating before the moisture contents reached the range of 16 to 30% on a wet basis (w.b.).

**Table 2. Effect of preparation, microwave, and pressing on the dehydrated powder yield of carrot (10 kg tray load) during preparation of powder**

Storage Intervals(S)	Preparation techniques (T)	Pressing (P)								Grand Mean (T)
		P <sub>1</sub> : Without pressing - (Control)				P <sub>2</sub> : Pressing				
		Microwave pre-treatment (M)				Microwave pre-treatment (M)				
		M <sub>1</sub> : control	M <sub>2</sub> : 30 S	M <sub>3</sub> : 60 S	Mean	M <sub>1</sub> : control	M <sub>2</sub> : 30 S	M <sub>3</sub> : 60 S	Mean	
<b>Powder yield (%)</b>	T <sub>1</sub> : Shredding	11.97	11.43	11.51	<b>11.64</b>	9.47	8.60	7.80	<b>8.62</b>	10.13
	T <sub>2</sub> : Crushing	11.58	11.41	11.38	<b>11.46</b>	7.26	6.70	6.20	<b>6.72</b>	9.09
	T <sub>3</sub> : Slicing	12.92	12.51	12.16	<b>12.53</b>	12.53	12.23	11.93	<b>12.23</b>	12.38
	<b>Mean (M, P of S<sub>1</sub>)</b>	<b>12.16</b>	<b>11.78</b>	<b>11.68</b>	<b>11.87</b>	<b>9.75</b>	<b>9.18</b>	<b>8.64</b>	<b>9.19</b>	
Grand Mean		<b>10.96</b>	<b>10.48</b>	<b>10.16</b>						
			<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>		<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>			
	<b>T</b>		0.042	0.121	<b>TxP</b>	0.060	0.171			
	<b>M</b>		0.042	0.121	<b>MxP</b>	0.060	0.171			
	<b>P</b>		0.034	0.099	<b>TxMxP</b>	0.103	0.297			
	<b>TxM</b>		0.073	0.210						
	<b>CV%</b>		1.70							

**Table 3. Effect of preparation, microwave, and pressing on the dehydration rate of carrot (10 kg tray load) during preparation of powder**

Storage Intervals(S)	Preparation techniques (T)	Pressing (P)								Grand Mean (T)
		P <sub>1</sub> : Without pressing - (Control)				P <sub>2</sub> : Pressing				
		Microwave pre-treatment (M)				Microwave pre-treatment (M)				
		M <sub>1</sub> : control	M <sub>2</sub> : 30 S	M <sub>3</sub> : 60 S	Mean	M <sub>1</sub> : control	M <sub>2</sub> : 30 S	M <sub>3</sub> : 60 S	Mean	
<b>Dehydration rate (g/min)</b>	T <sub>1</sub> : Shredding	8.88	8.97	9.08	<b>8.98</b>	7.04	6.76	6.15	<b>6.65</b>	7.81
	T <sub>2</sub> : Crushing	8.38	8.55	8.62	<b>8.52</b>	5.24	5.00	5.54	<b>5.26</b>	6.89
	T <sub>3</sub> : Slicing	7.97	8.02	8.07	<b>8.02</b>	8.04	8.08	9.49	<b>8.54</b>	8.28
	<b>Mean (M, P of S<sub>1</sub>)</b>	<b>8.41</b>	<b>8.51</b>	<b>8.59</b>	<b>8.50</b>	<b>6.77</b>	<b>6.61</b>	<b>7.06</b>	<b>6.82</b>	
Grand Mean		<b>7.59</b>	<b>7.56</b>	<b>7.83</b>						
			<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>		<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>			
	<b>T</b>		0.060	0.172	<b>TxP</b>	0.085	0.243			
	<b>M</b>		0.060	0.172	<b>MxP</b>	0.085	NS			
	<b>P</b>		0.049	0.140	<b>TxMxP</b>	0.147	0.421			
	<b>Tx M</b>		0.104	0.298						
	<b>CV%</b>		3.32							

Table 4. Effect of preparation, microwave, and pressing on the rehydration ratio of carrot of powder

Storage Intervals (S)	Preparation techniques (T)	Pressing (P)								Grand Mean (T)
		P <sub>1</sub> : Without pressing - (Control)				P <sub>2</sub> : Pressing				
		Microwave pre-treatment (M)				Microwave pre-treatment (M)				
		M <sub>1</sub> : control	M <sub>2</sub> : 30 S	M <sub>3</sub> : 60 S	Mean	M <sub>1</sub> : control	M <sub>2</sub> : 30 S	M <sub>3</sub> : 60 S	Mean	
Rehydration ratio	T <sub>1</sub> : Shredding	5.31	5.42	5.74	<b>5.49</b>	6.27	6.58	6.98	<b>6.61</b>	6.05
	T <sub>2</sub> : Crushing	6.12	6.33	6.7	<b>6.38</b>	6.48	6.58	7.11	<b>6.72</b>	6.55
	T <sub>3</sub> : Slicing	4.39	4.55	4.84	<b>4.59</b>	4.65	4.81	5.12	<b>4.86</b>	4.73
Grand Mean	Mean (M, P of S <sub>1</sub> )	<b>5.27</b>	<b>5.43</b>	<b>5.76</b>	<b>5.49</b>	<b>5.80</b>	<b>5.99</b>	<b>6.40</b>	<b>6.06</b>	
		<b>5.54</b>	<b>5.71</b>	<b>6.08</b>						
			<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>		<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>			
		T	0.035	0.100	<b>TxP</b>	0.049	0.141			
		M	0.035	0.100	<b>MxP</b>	0.049	NS			
		P	0.029	0.082	<b>TxMxP</b>	0.086	NS			
		TxM	0.060	NS						
	CV%	2.56								



Table 5. Effect of preparation, microwave, and pressing on the moisture content (%) of carrot powder during storage

Storage Intervals(S)	Preparation techniques (T)	Pressing (P)								Grand Mean (S, TS)	Grand Mean (T)
		P <sub>1</sub> : Without pressing - (Control)				P <sub>2</sub> : Pressing					
		Microwave pre-treatment (M)				Microwave pre-treatment (M)					
		M <sub>1</sub> :control	M <sub>2</sub> : 30 S	M <sub>3</sub> : 60 S	Mean (P <sub>1</sub> , S)	M <sub>1</sub> :control	M <sub>2</sub> : 30 S	M <sub>3</sub> : 60 S	Mean (P <sub>2</sub> , S)		
<b>S<sub>1</sub>: Initial</b>	T <sub>1</sub> : Shredding	5.98	5.82	5.46	5.75	4.97	4.65	4.44	4.69	5.22	<b>6.69</b>
	T <sub>2</sub> : Crushing	5.63	5.52	5.45	5.53	4.53	4.42	4.15	4.37	4.95	<b>5.91</b>
	T <sub>3</sub> : Slicing	7.30	6.24	5.71	6.42	5.82	5.35	4.85	5.34	5.88	<b>7.53</b>
	<b>Mean (MP of S<sub>1</sub>)</b>	<b>6.30</b>	<b>5.86</b>	<b>5.54</b>	<b>5.90</b>	<b>5.11</b>	<b>4.81</b>	<b>4.48</b>	<b>4.80</b>	<b>5.35</b>	
<b>S<sub>2</sub>:2 Months</b>	T <sub>1</sub> : Shredding	7.62	7.46	7.10	7.39	6.61	6.04	5.94	6.20	6.80	
	T <sub>2</sub> : Crushing	6.27	6.16	6.09	6.17	5.67	5.36	5.19	5.41	5.79	
	T <sub>3</sub> : Slicing	8.94	7.88	7.35	8.06	7.46	6.99	6.49	6.98	7.52	
	<b>Mean (MP of S<sub>2</sub>)</b>	<b>7.61</b>	<b>7.17</b>	<b>6.85</b>	<b>7.21</b>	<b>6.58</b>	<b>6.13</b>	<b>5.87</b>	<b>6.19</b>	<b>6.70</b>	
<b>S<sub>3</sub>:4 months</b>	T <sub>1</sub> : Shredding	7.86	7.70	7.34	7.63	6.85	6.28	6.18	6.44	7.04	
	T <sub>2</sub> : Crushing	6.51	6.40	6.33	6.41	5.71	5.57	5.33	5.54	5.98	
	T <sub>3</sub> : Slicing	9.18	8.12	7.59	8.30	8.70	7.23	6.73	7.55	7.93	
	<b>Mean (MP of S<sub>3</sub>)</b>	<b>7.85</b>	<b>7.41</b>	<b>7.09</b>	<b>7.45</b>	<b>7.09</b>	<b>6.36</b>	<b>6.08</b>	<b>6.51</b>	<b>6.98</b>	
<b>S<sub>4</sub>:6 months</b>	T <sub>1</sub> : Shredding	8.80	8.23	7.93	8.32	7.22	7.13	6.88	7.08	7.70	
	T <sub>2</sub> : Crushing	7.85	7.78	7.63	7.75	6.27	6.12	5.86	6.08	6.92	
	T <sub>3</sub> : Slicing	10.41	9.65	8.27	9.44	8.88	8.18	7.34	8.13	8.79	
	<b>Mean (MP of S<sub>4</sub>)</b>	<b>9.02</b>	<b>8.55</b>	<b>7.94</b>	<b>8.51</b>	<b>7.46</b>	<b>7.14</b>	<b>6.69</b>	<b>7.10</b>	<b>7.80</b>	
Grand Mean (P)		<b>7.70</b>	<b>7.25</b>	<b>6.85</b>	<b>7.27</b>	<b>6.56</b>	<b>6.11</b>	<b>5.78</b>	<b>6.15</b>		
Grand Mean (M)		<b>7.13</b>	<b>6.68</b>	<b>6.32</b>							
	<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>		<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>		<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>		<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>
<b>T</b>	0.021	0.061	<b>TxP</b>	0.030	0.086	<b>S</b>	0.025	0.071	<b>TxMxS</b>	0.076	0.213
<b>M</b>	0.021	0.061	<b>MxP</b>	0.030	0.086	<b>TxS</b>	0.044	0.123	<b>TxPxS</b>	0.062	0.174
<b>P</b>	0.017	0.049	<b>TxMxP</b>	0.052	0.148	<b>MxS</b>	0.044	0.123	<b>MxPxS</b>	0.062	0.174
<b>TxM</b>	0.037	0.105				<b>PxS</b>	0.036	0.100	<b>TxMxPxS</b>	0.107	0.301
<b>CV%</b>	2.67					<b>CV%</b>	2.77				

**Table 6. Effect of preparation, microwave, and pressing on the  $\beta$ - carotene (mg/100g) of carrot powder during storage**

Storage Intervals(S)	Preparation techniques (T)	Pressing (P)								Grand Mean (S, TS)	Grand Mean (T)
		P <sub>1</sub> : Without pressing - (Control)				P <sub>2</sub> : Pressing					
		Microwave pre-treatment (M)				Microwave pre-treatment (M)					
		M <sub>1</sub> :control	M <sub>2</sub> : 30 S	M <sub>3</sub> : 60 S	Mean (P <sub>1</sub> , S)	M <sub>1</sub> :control	M <sub>2</sub> : 30 S	M <sub>3</sub> : 60 S	Mean (P <sub>2</sub> , S)		
<b>S<sub>1</sub>: Initial</b>	T <sub>1</sub> : Shredding	175.95	177.43	178.04	177.14	139.18	133.52	120.67	131.12	154.13	<b>118.63</b>
	T <sub>2</sub> : Crushing	177.28	178.45	180.13	178.62	111.13	104.81	114.90	110.28	144.45	
	T <sub>3</sub> : Slicing	171.87	170.44	178.40	173.57	171.11	171.11	176.82	173.01	173.29	
	<b>Mean (MP of S<sub>1</sub>)</b>	<b>175.03</b>	<b>175.44</b>	<b>178.86</b>	<b>176.44</b>	<b>140.47</b>	<b>136.48</b>	<b>137.46</b>	<b>138.14</b>	<b>157.29</b>	
<b>S<sub>2</sub>:2 Months</b>	T <sub>1</sub> : Shredding	154.33	155.86	156.47	155.55	114.09	117.25	110.03	113.79	134.67	<b>105.91</b>
	T <sub>2</sub> : Crushing	155.70	156.88	158.56	157.05	80.61	92.11	101.13	91.28	124.17	
	T <sub>3</sub> : Slicing	150.30	148.87	156.77	151.98	149.48	148.41	155.24	151.04	151.51	
	<b>Mean (MP of S<sub>2</sub>)</b>	<b>153.44</b>	<b>153.87</b>	<b>157.27</b>	<b>154.86</b>	<b>114.73</b>	<b>119.26</b>	<b>122.13</b>	<b>118.71</b>	<b>136.78</b>	
<b>S<sub>3</sub>:4 months</b>	T <sub>1</sub> : Shredding	116.48	117.96	118.63	117.69	88.16	89.74	95.38	91.09	104.39	<b>124.72</b>
	T <sub>2</sub> : Crushing	117.81	118.98	120.72	119.17	70.85	72.87	77.01	73.58	96.37	
	T <sub>3</sub> : Slicing	112.40	110.98	118.93	114.10	111.64	110.52	117.40	113.19	113.65	
	<b>Mean (MP of S<sub>3</sub>)</b>	<b>115.56</b>	<b>115.97</b>	<b>119.43</b>	<b>116.99</b>	<b>90.22</b>	<b>91.04</b>	<b>96.60</b>	<b>92.62</b>	<b>104.80</b>	
<b>S<sub>4</sub>:6 months</b>	T <sub>1</sub> : Shredding	67.58	69.05	69.72	68.78	42.45	52.97	55.23	50.22	59.50	<b>59.53</b>
	T <sub>2</sub> : Crushing	68.90	70.07	71.81	70.26	44.20	42.16	54.80	47.05	58.66	
	T <sub>3</sub> : Slicing	50.55	62.12	70.02	60.90	53.73	61.61	64.49	59.94	60.42	
	<b>Mean (MP of S<sub>4</sub>)</b>	<b>62.34</b>	<b>67.08</b>	<b>70.52</b>	<b>66.65</b>	<b>46.79</b>	<b>52.25</b>	<b>58.17</b>	<b>52.40</b>	<b>59.53</b>	
<b>Grand Mean (P)</b>		<b>126.60</b>	<b>128.09</b>	<b>131.52</b>	<b>128.73</b>	<b>98.05</b>	<b>99.76</b>	<b>103.59</b>	<b>100.47</b>		
<b>Grand Mean (M)</b>		<b>112.32</b>	<b>113.92</b>	<b>117.55</b>							
	<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>		<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>		<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>		<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>
<b>T</b>	0.405	1.161	<b>T<sub>x</sub>P</b>	0.572	1.641	<b>S</b>	0.512	1.436	<b>T<sub>x</sub>M<sub>x</sub>S</b>	1.537	4.309
<b>M</b>	0.405	1.161	<b>M<sub>x</sub>P</b>	0.572	1.641	<b>T<sub>x</sub>S</b>	0.887	2.488	<b>T<sub>x</sub>P<sub>x</sub>S</b>	1.255	3.518
<b>P</b>	0.330	0.947	<b>T<sub>x</sub>M<sub>x</sub>P</b>	0.991	2.842	<b>M<sub>x</sub>S</b>	0.887	2.488	<b>M<sub>x</sub>P<sub>x</sub>S</b>	1.255	3.518
<b>T<sub>x</sub>M</b>	0.171	0.490				<b>P<sub>x</sub>S</b>	0.725	2.031	<b>T<sub>x</sub>M<sub>x</sub>P<sub>x</sub>S</b>	2.174	6.094
<b>CV%</b>	3.02					<b>CV%</b>	3.31				

**Table 7. Effect of preparation, microwave, and pressing on the sensory overall acceptability (9-point Hedonic scale) of carrot powder during storage**

Storage Intervals(S)	Preparation techniques (T)	Pressing (P)								Grand Mean (S, TS)	Grand Mean (T)
		P <sub>1</sub> : Without pressing - (Control)				P <sub>2</sub> : Pressing					
		Microwave pre-treatment (M)				Microwave pre-treatment (M)					
		M <sub>1</sub> :control	M <sub>2</sub> : 30 S	M <sub>3</sub> : 60 S	Mean (P <sub>1</sub> , S)	M <sub>1</sub> :control	M <sub>2</sub> : 30 S	M <sub>3</sub> : 60 S	Mean (P <sub>2</sub> , S)		
<b>S<sub>1</sub>: Initial</b>	T <sub>1</sub> : Shredding	6.20	6.75	6.85	6.60	6.88	6.75	7.00	6.88	6.74	<b>6.25</b>
	T <sub>2</sub> : Crushing	7.30	7.95	7.95	7.73	8.15	8.35	8.65	8.38	8.06	<b>7.56</b>
	T <sub>3</sub> : Slicing	6.95	7.00	7.18	7.04	7.10	7.10	7.40	7.20	7.12	<b>6.59</b>
	<b>Mean (MP of S<sub>1</sub>)</b>	<b>6.82</b>	<b>7.23</b>	<b>7.33</b>	<b>7.13</b>	<b>7.38</b>	<b>7.40</b>	<b>7.68</b>	<b>7.49</b>	<b>7.31</b>	
<b>S<sub>2</sub>:2 Months</b>	T <sub>1</sub> : Shredding	5.85	6.40	6.75	6.33	6.70	6.40	6.65	6.58	6.46	
	T <sub>2</sub> : Crushing	6.95	7.60	7.70	7.42	7.80	8.00	8.25	8.02	7.72	
	T <sub>3</sub> : Slicing	6.70	6.85	6.90	6.82	6.60	6.75	7.20	6.85	6.83	
	<b>Mean (MP of S<sub>2</sub>)</b>	<b>6.50</b>	<b>6.95</b>	<b>7.12</b>	<b>6.86</b>	<b>7.03</b>	<b>7.05</b>	<b>7.37</b>	<b>7.15</b>	<b>7.00</b>	
<b>S<sub>3</sub>:4 months</b>	T <sub>1</sub> : Shredding	5.58	6.13	6.48	6.06	6.93	6.13	6.38	6.48	6.27	
	T <sub>2</sub> : Crushing	6.68	7.33	7.58	7.19	7.53	7.73	8.03	7.76	7.48	
	T <sub>3</sub> : Slicing	6.43	6.58	6.63	6.54	6.83	6.48	6.93	6.74	6.64	
	<b>Mean (MP of S<sub>3</sub>)</b>	<b>6.23</b>	<b>6.68</b>	<b>6.89</b>	<b>6.60</b>	<b>7.09</b>	<b>6.78</b>	<b>7.11</b>	<b>6.99</b>	<b>6.79</b>	
<b>S<sub>4</sub>:6 months</b>	T <sub>1</sub> : Shredding	4.81	5.40	5.58	5.26	5.63	5.70	6.03	5.78	5.52	
	T <sub>2</sub> : Crushing	6.01	6.72	6.87	6.53	7.07	7.39	7.78	7.41	6.97	
	T <sub>3</sub> : Slicing	5.15	5.63	5.92	5.57	5.70	5.87	6.33	5.96	5.76	
	<b>Mean (MP of S<sub>4</sub>)</b>	<b>5.32</b>	<b>5.92</b>	<b>6.12</b>	<b>5.79</b>	<b>6.13</b>	<b>6.32</b>	<b>6.71</b>	<b>6.39</b>	<b>6.09</b>	
<b>Grand Mean (P)</b>	<b>6.22</b>	<b>6.69</b>	<b>6.86</b>	<b>6.59</b>	<b>6.91</b>	<b>6.89</b>	<b>7.22</b>	<b>7.00</b>			
<b>Grand Mean (M)</b>	<b>6.56</b>	<b>6.79</b>	<b>7.04</b>								
	<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>		<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>		<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>		<b>S.Em. ±</b>	<b>CD<sub>0.05</sub></b>
<b>T</b>	0.021	0.059	<b>T<sub>x</sub>P</b>	0.029	0.084	<b>S</b>	0.026	0.073	<b>T<sub>x</sub>M<sub>x</sub>S</b>	0.079	0.220
<b>M</b>	0.021	0.059	<b>M<sub>x</sub>P</b>	0.029	0.084	<b>T<sub>x</sub>S</b>	0.045	0.127	<b>T<sub>x</sub>P<sub>x</sub>S</b>	0.064	0.180
<b>P</b>	0.017	0.049	<b>T<sub>x</sub>M<sub>x</sub>P</b>	0.051	0.145	<b>M<sub>x</sub>S</b>	0.045	0.127	<b>M<sub>x</sub>P<sub>x</sub>S</b>	0.064	0.180
<b>T<sub>x</sub>M</b>	0.036	0.103				<b>P<sub>x</sub>S</b>	0.037	0.104	<b>T<sub>x</sub>M<sub>x</sub>P<sub>x</sub>S</b>	0.111	0.312
<b>CV%</b>	2.59					<b>CV%</b>	2.84				

**Rehydration ratio:** The perusal of data on the effect of different treatments on the rehydration ratio of carrot powder has been presented in Table 4.

**Effect of preparation techniques:** Among different preparation techniques (T), the rehydration ratio of carrot powder varied significantly between 4.73 and 6.55, with a maximum rehydration ratio of 6.55 in carrot powder which was prepared by giving crushing treatment (T<sub>2</sub>) and a minimum rehydration ratio of 4.73 in slicing (T<sub>3</sub>). Loesecke and Willard [27] reported that the rehydration characteristics of the dehydrated food are influenced by processing conditions as well as the sample composition used in dehydration.

**Effect of microwave pre-treatment:** Data shows that among different microwave pre-treatments grand mean rehydration ratio (M) of microwave-pretreated carrot powder varied significantly between 5.54 and 6.08, with maximum rehydration ratio (6.08) in carrot powder pre-treated with microwave for 60 seconds (M<sub>3</sub>) and minimum rehydration ratio (5.54) in carrot powder prepared without microwave pre-treatment for (M<sub>1</sub>). Pandit (2015) reported similar variations in the rehydration ratio (2.20 to 2.94) of dehydrated elephant foot yam due to variation in microwave exposure time and power. This can be explained by the high internal pressure produced by microwave heating which can cause the structure of carrot slices to expand and puff during dehydration and thus cause high rehydration ratio. Similar results were obtained by several researchers in other foods viz. kiwi fruits [28] and onions [9]. Kaur and Singh [24] reported that the rehydration characteristics of beetroot improved for samples dehydrated at 55°C air temperature followed by high microwave power (1080 W) which can be help in the reduction of shrinkage. Zielinska et al. [29] also reported an increase in the rehydration ratio due to microwave treatment in dried cranberries.

**Effect of pressing:** Further, it was observed that mean rehydration ratio (P) of the carrot powder when dehydrated followed by pressing treatment varied significantly from 5.49 to 6.06, with maximum rehydration ratio (6.06) in carrot powder when dehydrated by pressing treatment (P<sub>2</sub>) and minimum rehydration ratio (5.49) in carrot powder when dehydrated without pressing treatment (P<sub>1</sub>). Loesecke and Willard [27] reported that the rehydration characteristics of the dehydrated food are influenced by

processing conditions as well as the sample composition used in dehydration.

**Effect of treatment interactions:** The interaction of preparation techniques and pressing technique (TP) depicted variation in carrot powder rehydration ratio from 4.59 to 6.72, with maximum rehydration ratio of 6.72 in powder which was prepared by crushing technique followed by pressing treatment (T<sub>2</sub>P<sub>2</sub>) and minimum rehydration ratio of 4.59 in powder which was dehydrated by slicing technique without pressing treatment (T<sub>3</sub>P<sub>1</sub>).

**Moisture:** The perusal of data on the effect of different treatments on the moisture content of carrot powder during six months storage period is presented in Table 5.

**Effect of preparation techniques:** Among different preparation techniques (T), the moisture of carrot powder varied significantly between 5.91 and 7.53%, with minimum moisture in carrot powder which was prepared by giving crushing treatment (5.97%) and maximum moisture in slicing (7.53%). The lower moisture in carrot powder prepared using crushing treatment followed by pressing might be attributed to the release of most of the water from the tissue due to pressing and higher the dehydration rate for left over carrot pomace which requires less dehydration time for drying. Earlier studies also observed similar results for moisture in treatments that requires less dehydration time for okra, cauliflower, and onions [30]. Lower moisture and water activity were also observed in food products which requires less dehydration time as reported by Vaghashiya et al. [31] in dehydrated Aloe vera gel and Zinzala [32] for bael candy. Alam et al. [33] variations in water activity in dried carrot pomace powder prepared using different pre-treatments i.e., water blanching (WB), steam blanching (SB), citric acid blanching (CB), potassium metabi sulphate (KMS) dipping after blanching (WBS) and reported highest water activity (0.558) in untreated samples and lowest (0.379) value in WBS pre-treated samples.

**Effect of microwave pre-treatment:** Data shows that among different microwave pre-treatments grand mean moisture (M) of microwave- pretreated carrot powder varied significantly between 6.32 and 7.13%, with minimum moisture (6.32%) in carrot powder pre-treated with microwave for 60 seconds (M<sub>3</sub>) and maximum moisture (7.13%) in carrot powder

prepared without microwave pre-treatment ( $M_1$ ). Zielinska et al. [29] reported lower moisture content in microwave pre-treated samples compared to the control. Higher levels of moisture could lead to microbial spoilage and subsequent deterioration in quality. Sabry et al. [23] reported variations in the moisture content of the dried carrot powder due to microwave treatments. Water loss during microwave pre-treatments increased with an increase in microwave power and time.

**Effect of pressing:** Further, it was observed that the mean moisture (P) of the carrot powder when dehydrated followed by pressing treatment varied significantly from 6.15 to 7.27%, with minimum moisture (6.15%) in carrot powder when dehydrated followed by pressing treatment ( $P_2$ ) and maximum moisture (7.27%) in carrot powder when dehydrated without pressing treatment ( $P_1$ ). The lower moisture in carrot powder prepared pressing treatment might be attributed to the release of most of the water from the tissue due to pressing and higher the dehydration rate for left over carrot pomace which requires dehydration time for drying. Earlier studies also observed similar results for moisture in treatments that requires less dehydration time for okra, cauliflower and onions (Anon., 2017). Lower moisture and water activity were also observed in food products which require less dehydration time as reported by Vaghashiya [31] in dehydrated *Aloe vera* gel and Zinzala [32] for bael candy.

**Effect of storage:** Storage of the carrot powder for six months resulted significant increase in moisture (S) from an initial value of 5.35 to 7.80%. Chukwu and Lawal [34] reported that the moisture content of sweet potato flour increased from 4.32 to 5.36% within the first week of storage but decreased to 5.27% at the end of two weeks storage period and attributed this to loss of moisture during storage. Alinnor and Akalezi [35] reported that dehydrated flour can be stored successfully during storage if the moisture content of flour is below 8.3%. Maneepun et al. [36] reported increased the moisture content of sweet potato flour upto 8% in four and a half months. Raj et al. [17] registered a moisture content increase in potato flour of different cultivars from 3.07 to 4.78 per cent during storage period of six months. Sra et al. [8] reported that the moisture content of dried carrot slices increased from 6.3 to 8.4%. Abano et al. [26] reported that KMS pre-treated samples performed better in terms of moisture diffusivity

at all drying temperatures. Tiwari and Sarkar [37] reported 87.02% moisture content in fresh carrot and 4.2% in dried carrot powder. Ramachandra and Rao (2011) also reported a gradual increase in the moisture content of dehumidified air-dried *Aloe vera* gel powder during accelerated storage. Sharada [23] also reported a gradual increase in the moisture content of foam-mat-dried guava and banana powder during storage. The increase in moisture during the storage period of six months might be attributed to an increase in relative humidity in the atmosphere due to the continuations of rainy monsoons and due to the permeability of packaging materials to water vapours. A similar reason for other processed products had been reported by Raj et al. [38] for dehydrated onion rings.

**Effect of treatment interactions:** Interaction of preparation techniques, microwave pre-treatment, pressing and storage (TMPS) depicted variation in the moisture of carrot powder from 4.15 to 10.41%, with minimum increase of moisture from 4.15 to 5.85% ( $T_2M_3P_2S_1$  to  $T_2M_3P_2S_4$ ) in powder which was prepared by dehydrating crushed materials by giving microwave for 60 seconds followed by pressing and maximum increase of moisture from 7.30 to 10.41% ( $T_3M_1P_1S_1$  to  $T_3M_1P_1S_4$ ) in carrot powder which was prepared by dehydrating carrot slices without microwave and pressing pre-treatment. Interactions of  $T_xM$ ,  $T_xP$ ,  $M_xP$ ,  $T_xM_xP$ ,  $T_xS$ ,  $M_xS$ ,  $P_xS$ ,  $T_xM_xS$ ,  $T_xP_xS$ ,  $M_xP_xS$ , and  $T_xM_xP_xS$  were found to have significant effect.

**$\beta$ -carotene:** The perusal of data on the effect of different treatments on beta carotene of carrot powder during six months storage period has been presented in Table 6.

**Effect of preparation techniques:** Among different preparation techniques (T), the beta carotene of carrot powder varied significantly between 105.91 and 124.72 mg/100g, with minimum beta carotene (105.91 mg/100g) in carrot powder which was prepared by giving crushing treatment ( $T_2$ ) and maximum beta carotene (124.72 mg/100g) in slicing ( $T_3$ ). The food commodity having a smaller size possesses a large surface area and can release most of the soluble solids along with water. The crushing treatment observed minimum beta carotene in carrot powder which might be attributed to the release of large amounts of soluble solids along with water in the form of juice and thus less beta carotene remained in the dried carrot powder

due to pressing treatment. Tadesse et al. [39] reported the highest  $\beta$ -carotene (74.97 ppm) for the combination of the samples blanched at 55°C and soaked in 5% salt solutions. Alam et al. [33] reported the maximum  $\beta$ -carotene (633.57  $\mu\text{g } 100 \text{ g}^{-1}$ ) value for citric acid blanching (CB) pre-treated samples of carrot pomace followed by solar dried and the minimum value (186.01  $\mu\text{g } 100 \text{ g}^{-1}$ ) for CB pre-treated samples followed by sun-dried.

**Effect of microwave pre-treatment:** Data shows that among different microwave pre-treatments grand mean beta carotene (M) of microwave-pretreated carrot powder varied significantly between 112.32 and 117.55 mg/100g, with maximum beta carotene (117.55 mg/100g) in carrot powder pre-treated with microwave for 60 seconds ( $M_3$ ) and minimum beta carotene (112.32 mg/100g) in carrot powder prepared without microwave pre-treatment ( $M_1$ ). Sabry et al. [23] reported similar variations in beta carotene content of the dried carrot powder due to microwave treatments and higher beta carotene content was observed in the samples treated with microwave as compared to without microwave. Cui et al. (2004) also reported higher beta carotene content in dried carrot slices and Chinese chive leaves when given microwave treatment. Arjmandi et al. [40] reported an increase in phenolics and lycopene contents in smoothie due to microwave treatment.

**Effect of pressing:** Further, it was observed that the mean beta carotene (P) of the carrot powder when dehydrated followed by pressing treatment varied significantly from 100.47 to 128.73 mg/100g, with minimum beta carotene 100.47 mg/100 g in carrot powder when dehydrated followed by pressing treatment ( $P_2$ ) and maximum beta carotene 128.73 mg/100g in carrot powder when dehydrated without pressing treatment ( $P_1$ ). The food commodity having smaller size possess a large surface area can release most of the soluble solids along with water due to pressing. The pressing treatment observed minimum beta carotene in carrot powder which might be attributed to the release of large amounts of soluble solids along with water in the form of juice and thus less beta carotene remained in the dried carrot powder due to the pressing treatment.

**Effect of storage:** Storage of the carrot powder for six months resulted significant decrease in beta carotene (S) from an initial value of 157.29 to 59.53 mg/100g. Hal [41] reported that during

storage, auto-oxidation of carotenoids may take place, leading to a loss of color and an undesirable decline in beta-carotene value. The stability of beta-carotene proved to be strongly and adversely affected by storage temperature and light. Chaturvedi et al. [42] reported that carotenoid content decreased during storage from 59.8 per cent to 52 per cent of  $\beta$ -carotene after six months of storage. Sra et al. [8] reported a decrease in the  $\beta$ -carotene content from 69.4 to 49.2 mg/100 g during storage. Carrot slices pre-treated with 6% KMS and packed in AFL pouches were found to retain the highest carotene content. Singh et al. [43] reported that the pressure-cooking treatment utilized for the preparation of carrot powder and grits revealed that carrot powder had higher  $\beta$ -carotene levels (36.94 mg/100 g) than carrot grits (33.48 mg/100 g) which further showed a gradual loss of  $\beta$ -carotene during storage in all the packaging materials but higher  $\beta$ -carotene content reported to be retained in the aluminium laminated pack and can be stored safely up to 6 months at room temperature. A similar decrease in  $\beta$ -carotene content (S) from an initial value of 17.53 mg/100g to 9.84 mg/100g was observed during six months of storage of the sweet potato flour by Ashuqullah [44].

**Effect of treatment interactions:** Interaction of preparation techniques, microwave pre-treatment, pressing and storage (TMPS) depicted variation in beta carotene of carrot powder from 42.16 to 177.28 (mg/100g), with a minimum decrease of beta carotene from 144.90 to 54.80 mg/100g ( $T_2M_3P_2S_1$  to  $T_2M_3P_2S_4$ ) in powder which was prepared by dehydrating crushed materials by giving microwave for 60 seconds followed by pressing and maximum decrease of beta-carotene from 171.87 to 50.55 mg/100g ( $T_3M_1P_1S_1$  to  $T_3M_1P_1S_4$ ) in carrot powder which was prepared by dehydrating carrot slices without microwave and pressing pre-treatment. Interactions of TxM, TxP, MxP, TxMxP, TxS, MxS, PxS, TxMxS, TxPxS, MxPxS, and TxMxPxS were found to have significant effect.

**Overall acceptability:** The perusal of data pertaining on the different treatments on the overall acceptability of carrot powder during six months storage period has been presented in Table 7.

**Effect of preparation techniques:** Among different preparation techniques (T), the overall acceptability (9-point Hedonic scale) of

carrot powder varied significantly between 6.25 and 7.56, with maximum overall acceptability (7.56) in carrot powder which was prepared by giving crushing treatment ( $T_2$ ) and minimum overall acceptability (6.25) in shredding ( $T_1$ ). Nath et al. (2012) reported variations in taste in ginger candy when prepared using different slice thicknesses (5-25 mm) and blanched for different times (10-30 min). The higher overall acceptability in carrot powder prepared using crushing treatment followed by pressing might be attributed to a higher dehydration rate and requirement of less dehydration time for drying. Earlier studies also observed similar results for overall acceptability in treatments that requires less dehydration time for okra, cauliflower, and onions [30]. Higher overall acceptability was also observed in food products that requires less dehydration time as reported by Zinzala [32] for bael candy. The higher overall acceptability score in carrot powder which was prepared by giving crushing treatment attributed to the retention of more colour, taste, flavour, and texture in dehydrated powder. Chaturvedi et al. [42] reported variations in overall acceptability in dehydrated carrots using different pre-treatments and drying techniques i.e., tray dried (TD), tray dried and radiated (TDR), Infra-red dried (IR), Infra-red dried and radiated (IRR). They reported a maximum overall acceptability value (4.93) in IRR and the lowest value (4.17) in TD. Kukanoor et al. [45] reported the highest value (4.46) of texture and overall acceptability (4.55) in the dehydrated carrot slices blanched for 2 minutes and steeped in 300 brix syrup followed by 6% salt for 2 hours. Whereas, Raj et al. [17] reported significant variations in the overall acceptability of the potato flours of the varieties due to variations in the NEB, sensory color, texture score, and these scores of the product directly correlated with the overall acceptability of the product.

**Effect of microwave pre-treatment:** Data shows that among different microwave pre-treatments grand mean overall acceptability (M) of microwave-pretreated carrot powder varied significantly between 6.56 and 7.04, with maximum overall acceptability (7.04) in carrot powder pre-treated with microwave for 60 seconds ( $M_3$ ) and minimum overall acceptability (6.56) in carrot powder prepared without microwave pre-treatment ( $M_1$ ). Pandit [46] reported similar variations in the overall acceptability of dehydrated elephant foot yam due to variations in microwave exposure time and power. This can be explained by the high internal pressure produced by microwave heating

which can cause the structure of carrot slices to expand and puff during dehydration which provides a more porous structure in the food product. Zielinska et al. [29] also an reported improvement in color due to microwave treatment in dried cranberries.

**Effect of pressing:** Further, it was observed that the mean overall acceptability (P) of the carrot powder when dehydrated followed by pressing treatment varied significantly from 6.59 to 7.00, with maximum overall acceptability (7.00) in carrot powder when dehydrated followed by pressing treatment ( $P_2$ ) and minimum overall acceptability (6.59) in carrot powder when dehydrated without pressing treatment ( $P_1$ ). The higher overall acceptability in carrot powder prepared using pressing treatment might be attributed to a higher dehydration rate and requirement of less dehydration time for drying.

**Effect of storage:** Storage of the carrot powder for six months resulted significant decrease in overall acceptability (S) from an initial value of 7.31 to 6.09. The decrease in overall acceptability score might be attributed to an increase in moisture content, water activity and NEB in dehydrated carrot powder during storage. Sra et al. [8] reported a decrease in the overall acceptability of dried carrot slices from 8.2 to 7.1 during six months storage period. Chaturvedi et al. [42] reported a decrease in the overall acceptability of dehydrated carrots during six months of storage. Pandit [46] reported similar variations in the overall acceptability of dehydrated elephant foot yam due to variations in microwave exposure time and power during six months of storage. This can be explained by the high internal pressure produced by microwave heating which can cause the structure of carrot slices to expand and puff during dehydration which provides a more porous structure in the food product.

**Effect of treatment interactions:** Interaction of preparation techniques, microwave pre-treatment, pressing and storage (TMPS) depicted variation in overall acceptability of carrot powder from 4.81 to 8.65, with minimum a decrease of overall acceptability from 8.65 to 7.78 ( $T_2M_3P_2S_1$  to  $T_2M_3P_2S_4$ ) in powder which was prepared by dehydrating crushed materials by giving microwave for 60 seconds followed by pressing and maximum decrease of overall acceptability from 6.95 to 5.15 ( $T_3M_1P_1S_1$  to  $T_3M_1P_1S_4$ ) in carrot powder which was prepared by dehydrating carrot slices without microwave

and pressing pre-treatment. Interactions of TxM, TxP, MxP, TxMxP, TxS, MxS, PxS, TxMxS, TxPxS, MxPxS, and TxMxPxS were found to have significant effect [47,48].

#### 4. CONCLUSION

It can be concluded that carrot powder prepared by dehydrating crushed materials by giving microwave pre-treatment for 60 seconds followed by pressing (after single-strength juice extraction) can remain safe for six months in PP bags (400 gauge) with minimum changes in chemical and sensory quality. This technique possesses the additional benefit of preparing two products simultaneously i.e. juice and powder instead of the product either alone. Thus, the developed technologies can commercially be explored by the food processing industry. Therefore, profitable utilization of carrots grown in India by processing can ensure better returns to the growers, processors and consumers.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

#### ACKNOWLEDGEMENT

Authors acknowledge Department of Post Harvest Technology for the funding of the research and providing facilities.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Jabbar S, Abid M, Hu B, Wu T, Hashim MM, Lei S, Zeng X. Quality of carrot juice as influenced by blanching and sonication treatments. *LWT-Food Sci. and Tech.* 2013;55(1):16-21.
2. Qin L, Xu S, and Zhang W. Effect of enzymatic hydrolysis on the yield of cloudy carrot juice and the effects of hydrocolloids on color and cloud stability during ambient storage. *J. Sci. Food. Agric.* 2005; 8(5):505–512.

3. Gopalan C, Shastri RB, Balasubramanian SC. Nutritive value of Indian foods. National Institute of Nutrition. Ind. Council of Medical Res., Hyderabad; 2004.
4. Shivhare US, Gupta M, Basu S, Raghavan GSV. Optimization of blanching process for carrots. *J. Food. Process. Eng.* 2009;3(2):587-605.
5. Dede S, Alpas H, Bayındırlı A. High hydrostatic pressure treatment and storage of carrot and tomato juices: Antioxidant activity and microbial safety. *J. Sci. Food and Agri.* 2007;87(5):773-782.
6. Khan MR. Osmotic dehydration technique for fruits preservation-a review. *Pak. J. Food. Sci.* 2012;22(2):71-85.
7. Kumar N, Kumar K. Development of carrot pomace and wheat flour based cookies. *J. of Pure and App. Sci. and Tech.* 2011;1(1):5-11.
8. Sra SK, Sandhu KS, Ahluwalia P. Effect of treatments and packaging on the quality of dried carrot slices during storage. *J. food sci. technol.* 2014;51(4):645-654.
9. Hiranvarachat B, Devahastin S, Chiewchan N, Raghavan GV. Structural modification by different pretreatment methods to enhance microwave-assisted extraction of  $\beta$ -carotene from carrots. *J. Food Eng.* 2013;115(2):190-197.
10. Raj D, Huddar AG, Gupta P. Effect of dehydration temperatures on the quality characteristics of dehydrated onion rings during storage. *Udyanika J. Horti. Sci.* 2004;10(4):47-52.
11. AOAC. Official methods of analysis, Association of Official Analytical Chemists. 14<sup>th</sup> edn, Arlington, Virginia, USA; 1984.
12. Ranganna S. Handbook of analysis and quality control for fruit and vegetable products. Tata McGraw Hill Publishing Co. Ltd., New Delhi (India); 1997.
13. Amerine MA, Pangborn RM, Roessler EB. Principles of sensory evaluation of food. Acad. Press, New York: Academic Press; 1965.
14. Panse VG, Sukhatme PV. Statistical methods for research workers. ICAR publication, New Delhi. 1967;20-40.
15. Sharma KD, Karki S, Thakur NS, Attri S. Chemical composition, functional properties and processing of carrot a review. *J. Food Sci. and Tech.* 2012; 49(1):22-32.
16. Wilczyński K, Kobus Z, Dżiki D. Effect of press construction on yield and quality of



- apple juice. Sustainability. 2019;11(13):3630.
17. Raj D, Lal BB, Joshi VK. Yield, quality and storability of the potato flour of different Indian cultivars. Int. J. Fd. Fermentation. 2011;1(1):111-117.
  18. Gerard KA, Roberts JS. Microwave heating of apple mash to improve juice yield and quality. LWT-Food Sci. and Tech. 2004;37(5):551-557.
  19. Rayman A, Baysal T. Yield and quality effects of electropulsation and microwave applications on carrot juice production and storage. J. Food Sci. 2011;76(4):C598-C605.
  20. Leeratanarak N, Devahastin S, Chiewchan N. Drying kinetics and quality of potato chips undergoing different drying techniques. Journal of Food Engineering. 2006;77(3):635-643.
  21. Hatamipour MS, Kazemi HH, Nooralivand A, Nozarpoor A. Drying characteristics of six varieties of sweet potatoes in different dryers. Journal of Food and Bio Products Processing. 2007;85(3):171-177.
  22. Dinrifo RR. Effects of pre-treatments on drying kinetics of sweet potato slices. Agricultural Engineering International Journal. 2012;14(3):136-145.
  23. Sabry ZA, Bahloul HE, El-Desouky AI and Assous MTM. Effect of microwave pretreatment and drying methods on the carrot powder quality. Middle East J. App. Sciences. 2016;6(2):349-356.
  24. Kaur K, Singh AK. Drying kinetics and quality characteristics of beetroot slices under hot air followed by microwave finish drying. African J. Agril. Res., 2014;9(12):1036-1044.
  25. Andres A, Bilbao C, Fito P. Drying kinetics of apple cylinders under combined hot air-microwave dehydration. Journal of Food Engineering. 2004;63(1):71-78.
  26. Abano EE, Ma H, Qu W. Effects of pretreatments on the drying characteristics and chemical composition of garlic slices in a convective hot air dryer. J Agric. Food Tech. 2011;1(5):50-58.
  27. Loesecke V, Willard H. Drying and dehydration of foods. Reinhold Publishing Corporation; 1955.
  28. Maskan M. Kinetics of colour change of kiwifruit during hot air and microwave drying. J. Food Eng; 2001.
  29. Zielinska M, Zielinska D, Markowski M. The effect of microwave-vacuum pretreatment on the drying kinetics, color and the content of bioactive compounds in osmo-microwave-vacuum dried cranberries (*Vaccinium macrocarpon*). Food and Bioprocess Tech. 2018;11(3):585-602.
  30. Anonymous. AGRESKO report of horticulture sub-committee, navsari agricultural university. In: 13<sup>th</sup> Combined Joint Agresko Meeting, Sardar Krushinagar Dantiwada Agricultural University, Dantiwada; April 5-7, 2017.
  31. Vaghashiya JM, Raj D, Bhandari DR, Desai CS, Patel JM. Health drink preparation using Aloe vera, bitter gourd, aonla and guava. Green Farming. 2016;7(6):1486-1490.
  32. Zinzala PB. Processing and value addition of Bael (*Aegle marmelos* L.). Thesis Ph.D., Navsari Agricultural University, Navsari, India; 2019.
  33. Alam MS, Gupta K, Khaira H, Javed M. Quality of dried carrot pomace powder as affected by pretreatments and methods of drying. Agril. Eng. International: CIGR Journal. 2013;15(4):236-243.
  34. Chukwu O, Lawal AO. Comparative study of storage stability of sweet potato and yam flours. International Journal of Emerging Technologies in Engineering Research, 2015;3(3):44-49.
  35. Alinnor IJ, Akalezi CO. Proximate and mineral compositions of *Dioscorea rotundata*. American Journal Research Communication. 2014;2(3):119 – 126.
  36. Maneepun SS, Reungmanee-paitoon M. Yunchalad. Product development for root and tuber crops, Vol I –Asia (G. J. Scott, S. Wiersema, and P. I. Ferguson, eds.), CIP, Lima: 1992;229.
  37. Tiwari S, Sarkar N. Development and evaluation of carotene rich carrot powder. Int. J. Res. Biosci. Agric. Tech. 2018;6(1):123-131.
  38. Raj D, Huddar AG, Subanna VC, Gowda IND. Effect of packaging on the quality characteristics of dehydrated onion rings during storage. Beverages Fd. World. 2006;36(5):33-35.
  39. Tadesse TF, Abera S, Worku S. Nutritional and sensory properties of solar-dried carrot slices as affected by blanching and osmotic pre-treatments. Int. J. Food Sci. and Nutrition Eng. 2015;5(1):24-32.
  40. Arjmandi M, Oton M, Artés-Hernandez F, Gomez PA, Artes F, Aguayo E. Effect of microwave treatments on the quality of a smoothie. In VIII International Postharvest

- Symposium: Enhancing Supply Chain and Consumer Benefits-Ethical and Technological Issues. 2016;1194:1481-1486.
41. Hal VM. Quality of sweet potato flour during processing and storage. *Food Reviews International*. 2007;16:1-37.
  42. Chaturvedi A, Sujatha V, Ramesh C, Babu JD. Development of shelf stable intermediate moisture carrot (*Daucus carota*) shreds using radiation as hurdle technology. *Int. Food Res. J.* 2013;20(2):775.
  43. Singh P, Kulshrestha K, Kumar S. Effect of storage on  $\beta$ -carotene content and microbial quality of dehydrated carrot products. *Food Biosci.* 2013;2:39-45.
  44. Ashuqullah. Evaluation of sweet potato {*Ipomoea batatas* (L.) Lam} varieties and pre-treatments for dehydration into flour. M.Sc. Thesis, Navsari Agricultural University, Navsari, Gujarat, India; 2017.
  45. Kukanoor L, Karadiguddi M, Rayar S, Jaishankar HP. Effect of pre-treatments on physical and sensory qualities of dehydrated carrot slices. *J. Hort.* 2014;1(3):1-3.
  46. Pandit PD. Microwave assisted convective drying characteristics of elephant foot yam [*Amorphallus paeoniifolius* (Densst.) NICOLSON] Thesis. M.Sc. (Agri.), Navsari Agricultural University, Navsari, India. 2015;87-91.
  47. Kalse SB, Patil MM, Jain SK. Microwave drying of onion slices. *Res. J. Chemical Sci.* 2012;2(4):57-60.
  48. Sharada S. Studies on effect of various operating parameters and foaming agents- Drying of fruits and vegetables. *Inter. J. Modern Engin. Res.* 2013;3(3):1512-1519.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/123377>