



Water Quality Distribution in Situ Gunung Putri Bogor Regency, Indonesia

**Zahidah Hasan^{1*}, Thoriq Ilham¹, Yuli Andriani¹, Heti Herawati¹
and Fachmijany Sulawesty²**

¹*Faculty of Fisheries and Marine Sciences, Padjadjaran University, Indonesia.*

²*Limnology Research Center, Indonesian Institute of Sciences, Indonesia.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJFAR/2019/v4i330056

Editor(s):

(1) Dr. Telat Yanik, Professor, Department of Aquaculture, Faculty of Fisheries, Atatürk University, Turkey.

Reviewers:

(1) Moses Mwajar Ngeiywa, University of Eldoret, Kenya.

(2) K. D. Mini, Mahatma Gandhi University, India.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50649>

Original Research Article

Received 01 July 2019

Accepted 02 September 2019

Published 12 September 2019

ABSTRACT

The purpose of this study was to determine water quality distribution in Situ Gunung Putri. This research was conducted from February to March 2019, carried out in Situ Gunung Putri and at the Limnology Research Center of the Indonesian Institute of Sciences, both located in Bogor, West Java. The method used was the survey method. The results showed that, at the surface level, the temperature ranged between (28.14±0.52 and 30.61±1.76)°C, transparency between (0.38±0.12 and 0.65±0.06) m, dissolved oxygen (DO) was between 1.92±1.10 and 5.7± 2.55 mg / L, chemical oxygen demand (COD) was between 42.22±9.86 and 54.09±15.89, nitrate between (0.047±0.009 and 0.05±0.015) mg / L, phosphate of (0.015±0.002 and 0.022± 0.004) mg / L and pH was between 7.11± 0.27 and 7.33 ±0.34. Meanwhile at the Secchi depth values of the temperature ranged between (2.93±0.71 and 29.96 ±1.71)°C, DO was 1.14 ±0.90 and 3.37±1.63 mg / L, nitrate was 0.045±0.008 and 0.056±0.019 mg / L, phosphate was 0.016±0.003 and 0.035± 0.043 mg / L and pH was between 6.68± 0.33 and 7.11 ±0.28. Based on these values water quality parameters except COD in Situ Gunung Putri is still at the optimal range for aquatic organisms.

Keywords: Water quality; Situ Gunung Putri; Bogor.

**Corresponding author: Email: ibuzah@gmail.com;*

1. INTRODUCTION

Situ is a Sundanese for small lake that is formed naturally or artificially, the source of water comes from springs, rainwater or surface runoff. Situ or lake has two main functions, ecological functions and socio-economic-cultural functions. The ecological function of situ as an area of biodiversity conservation, both flora and fauna and as a water regulatory systems, flood control, habitat for wild life or protected species, while the socio-economic-cultural function is to meet the needs of human life, among others as a usable source of water by the community both directly in agriculture, fisheries, industry and household activities [1].

Situ Gunung Putri is located in Gunung Putri Village, Gunung Putri District, Bogor Regency, West Java. The location is right by the Jagorawi toll road and is located amongst industrial area, housing area and agricultural area. Situ Gunung Putri was once used as a tourism object around 2010 to 2013, but discontinued since 2014. Since then the location has been used as a fishing area. Activities around Situ Gunung Putri are industries, agriculture and housing. these activities will affected on several water quality, including COD, dissolved oxygen nitrate and phosphate. Changes in the condition of Situ Gunung Putri waters over time due to increased waste originating from industrial, agricultural and household activities. Especially after being used as a water tourism destination, the water quality of Situ Gunung Putri tend to decreased.

Water quality characteristics can be distinguished physical and chemical parameters. Physical parameters such as temperature and transparency, chemical parameters such as pH, DO, BOD, and COD. The solubility of gas in water as well as biological activity in waters is strongly influenced by temperature changes. Each increase in temperature of 10°C will cause an increase in metabolic rate of 2-3 times. Increased metabolic rate of organisms will cause oxygen demand to increase, on the other hand increasing temperatures will cause oxygen solubility in water to decrease. This phenomenon will cause aquatic biota to experience difficulty in the process of respiration [2]. Aquatic organisms can live in ideal conditions in the pH of the waters ranging from weak acids to weak bases or in the range of pH values 6-9. Conditions of waters that are strong acids (less than 6) and strong bases (more than 9) will endanger the survival of the biota in the waters. Therefore,

research on the water quality of Situ Gunung Putri is needed in order to obtain better aquatic resources management recommendations.

2. METHODS

This research was carried out from February to March 2019 at Situ Gunung Putri, Bogor and in the laboratory of the Limnologi Research Center, Indonesian Institute of Sciences, Cibinong, Bogor, West Java.

The method used in this study was a survey method. Samples obtained from five stations that were defined by land use as it can be seen in Fig. 1. Sampling was done six times with one week interval. Water quality parameters that were analyzed are temperature, Transparency, DO, COD, Nitrate, Phosphate and pH, both from surface level and secchi depth except tranparency and COD not taken in secchi depth.

Temperature, DO, pH measured *in situ* using Horiba Water Quality Checker and transparency measured using the Secchi Disc. measurement of nitrate, phosphate concentrations and COD value was carried out in the laboratory. Measurement of nitrate concentration was carried out using the brucine method using a UV-Visible spectrophotometer at a wavelength of 420 nm. Phosphate concentration was measured by ascorbic acid method using a UV-Visible spectrophotometer at a wavelength of 880 nm. Meanwhile COD measurement was carried out in the Environmental Productivity laboratory at Bogor Agriculture University, using the closed reflux method with a spectrophotometer at a wavelength of 600 nm [3].

3. RESULTS AND DISCUSSION

3.1 Water Quality Parameters

The result of water quality parameters consisted of physical and chemical during observations presented in Table 1.

3.2 Temperature

The highest temperature was found on the surface of the water at the station 1 which reaches 30.61°C. The lowest temperature on the water surface was 28.14°C at station 5 is also still within the ideal temperature limit for the survival of aquatic organisms. Average tempature during study can be seen in Fig. 2.




Fig. 1. Data collection location at Situ Gunung Putri

Table 1. Physical and chemical parameters during study

Parameters/depth	Station				
	1	2	3	4	5
Surface					
Temperature (°C)	30.61±1.76	29.54±1.12	29.73±1.36	28.58±0.77	28.14±0.52
Transparency (m)	0.38±0.12	0.5±0.17	0.53±0.09	0.65±0.06	0.63±0.12
DO (mg/L)	5.7±2.55	2.81±1.38	3.96±3.73	3.06±2.69	1.92±1.10
COD (mg/L)	46.59±10.37	46.89±13.33	54.09±15.89	42.22±9.86	44.59±7.65
Nitrate (mg/L)	0.049±0.006	0.058±0.015	0.05±0.009	0.047±0.009	0.049±0.007
Phosphate (mg/L)	0.018±0.005	0.017±0.003	0.015±0.002	0.022±0.004	0.015±0.003
pH	7.25±0.23	7.11±0.27	7.27±0.10	7.15±0.17	7.33±0.34
Secchi depth					
Temperature (°C)	29.96±1.71	28.56±0.95	29.03±1.49	28.31±0.64	27.93±0.71
DO (mg/L)	3.37±1.63	1.30±0.94	2.61±5.05	2.07±2.07	1.14±0.90
Nitrate (mg/L)	0.051±0.004	0.046±0.007	0.045±0.008	0.050±0.013	0.056±0.019
Phosphate (mg/L)	0.019±0.001	0.020±0.004	0.035±0.043	0.024±0.004	0.016±0.003
pH	7.11±0.28	6.86±0.33	6.93±0.30	6.90±0.32	7.02±0.27

The temperature in the secchi depth was also measured to determine the effect on the transparency of sunlight. According to [4] sunlight entering the waters will be absorbed and change into heat energy. In the surface layers waters will have a higher temperature and lower density than in the lower layers.

The highest temperature at secchi depth was obtained at station 1 reaching 29.96°C. The lowest temperature measured in secchi depth was at station 5 of 27.93°C. The difference between the temperature on the surface with secchi depth was not too significant at all stations. This is in line with the findings of [5] that

the temperature obtained at each depth is not much different especially between surface layers and secchi depth. Overall, temperature value di Situ Gunung Putri still in tolerable range for aquatic organisms as stated in the Republic of Indonesia Government Regulation No. 82 of 2001 for class III (suitable for aquatic organisms to live in, deviation 3°C from normal temperature).

3.3 Transparency

The value of light penetration or transparency as measured by secchi disc is strongly influenced by the intensity of sunlight at the surface water, turbidity and the density of plankton in a waters

column. Light penetration is a limiting factor for photosynthetic organisms (phytoplankton) and also causes death in certain organisms [6]. Fig. 3 showed transparency during study.

The lowest transparency value at station 1 was 0.38 ± 0.12 m and the highest transparency at station 4 was 0.65 ± 0.06 m. The transparency value according to [7] was still within normal limits because the transparency of light is good for plankton growth optimally which is 30 cm (0.30 m) to 50 cm (0.50 m). Further stated that phytoplankton growth at euphotic zone that can be estimated with multiplying secchi depth visibility by 2.




Fig. 2. Temperature during observation




Fig. 3. Water transparency during study

3.4 Dissolved Oxygen

There was a difference in dissolved oxygen concentration between the surface water and the secchi depth of each station caused by differences in sunlight penetration. The highest DO obtained at station 1 and the lowest one at station 5 both in surface water and secchi depth. In surface water the highest average value was 5.70 ± 2.55 mg/L and at secchi depth the value was 3.37 ± 1.63 g/L. meanwhile lowest value was 1.92 ± 1.1 mg/L and at secchi depth was 1.14 ± 0.90 mg/L.

According to [8] DO concentrations which have decreased as depth increase due to photosynthesis activities is higher in the surface waters level than in deeper waters level.

In addition, at the bottom layer of waters decomposition of organic matter that utilize oxygen occur more [9]. Based on Fig. 4, dissolved oxygen in station 2 and 5 under the quality standard for class III according to [10] both at surface and secchi depth, meanwhile in station 1 the concentration suitable with standard both in two layer. On the other hand, in others two station DO concentration at surface layer suitable with the standard but not in the secchi depth. minimum concentration of 1.0 mg/L is essential to sustain fish for long period, and 5.0 mg/L are adequate for fish growth [11].

3.5 Acidity/Alkalinity (pH)

The highest average pH value in surface obtained at station 5 with a value of 7.33 ± 0.34 and the lowest was found at station 2 with a value of 7.11 ± 0.27 . Whereas the highest value in secchi depth occurred at station 1 with a value of 7.11 ± 0.28 and the lowest ones at station 2 with a value of 6.86 ± 0.33 . The difference in pH between surface and secchi depth is normal condition. according to [12] statement, that the decrease in pH at the deeper waters level occurs due to an increase in microbial activity to decompose organic matter so that O_2 decreases and CO_2 increases. which will cause the waters to become more acidic. Overall, pH value di Situ Gunung Putri still in tolerable range for aquatic organisms as stated in the Republic of Indonesia Government Regulation No. 82 of 2001 for class III with value range between 6-9.

3.6 COD

The highest average COD (Chemical Oxygen Demand) value was found at station 3 with an

average value of 54.09 ± 15.89 mg / L while the lowest COD concentration was at station 4 with an average value of 42.22 ± 9.86 mg / L. Based on Indonesian Government regulation Number 82 of 2001. Those values were still below the threshold limit for class III of 50 mg/L (suitable for aquaculture activities, and aquatic organisms), but its very close to border and need to get serious attention. COD distribution during study are showed in Fig. 6.

3.7 Nitrate

Based on the results as shown in Fig. 7, the highest average value of nitrate concentration in surface layer was found at station 2 at 0.058 ± 0.015 mg / L and the lowest one was obtained at station 4 of 0.047 ± 0.009 mg / L. On the other hand the highest average nitrate concentration in secchi depth found at station 5 of 0.056 ± 0.019 mg / L and the lowest occurred at station 3 with a value of 0.045 ± 0.008 mg / L.

Concentration of nitrate in waters are influenced by several parameters such as dissolved oxygen and organic matter content. According to [13], if dissolved oxygen concentration is low, the activity of microorganisms in decomposing of organic matter and nitrification process will be affected. This result is similar to [14] findings in Jakarta Bay, who found that there is relationship between nitrate concentrations with oxic condition. In Fig. 7 it can be seen that there is no significant difference between surface level and secchi depth in nitrate concentration. Compare [15] that conducted research Haryana Lakes, India found nitrate concentration range between 26.1-47.67 mg/l, nitrate concentration in Situ Gunung Putri were very low.

3.8 Phosphate

Phosphate concentration can be seen in Fig. 8. The highest average phosphate concentration in surface level was at station 4 with a value of 0.022 ± 0.004 mg / L and the lowest occurred at station 3 with a value of 0.015 ± 0.002 mg / L. Meanwhile the highest phosphate concentration in secchi depth was at station 3 with a value of 0.035 ± 0.0043 mg / L and the lowest is at station 3 with a value of 0.016 ± 0.003 mg / L.

Phosphates are needed in small amounts, as a transfer of energy from the outside into cells of the organism [4]. According to Indonesian Government regulation Number 82 of 2001,

phosphate concentration in all of the station was still far below the threshold of phosphate concentration for aquaculture and aquatic organisms (class III), which was less than 1 mg / L. From Fig. 8 it can be seen that there is no

significance difference between surface layer and secchi depth in phosphate concentration. Low concentration of phosphate also found in Naivasha Lake, Kenya as stated by [16] that range between 0.001-0.079 mg/L.




Fig. 4. Dissolved oxygen during study




Fig. 5. pH during observation




Fig. 6. COD value during observation




Fig. 7. Nitrate concentration during study




Fig. 8. Phosphate concentration during observation

4. CONCLUSIONS

Based on this research, it can be concluded that water quality in Situ Gunung Putri is still at the optimal range according to Indonesian Government regulation Number 82 of 2001, except for Chemical Oxygen Demand and Dissolved Oxygen at stasion 2 and 5 which is very close to threshold value.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Sittadewi EH. Fungsi Strategis Danau Tondano, Perubahan Ekosistem dan

2. Masalah yang Terjadi. *Jurnal Teknik Lingkungan*. 2008;99(1):59-66. (In Indonesian).
3. Satino. *Handout Limnologi*. Yogyakarta. 2010: FMIPA UNY, Yogyakarta (in Indonesian).
4. APHA. *Standard methods for the examination of waters and wastewater*. 21st edition. American Public Health Association. American Water Works Association. Water Pollution Control Federation. Washington. D.C. 2015;1467.
5. Effendi H. *Telaah kualitas air: Bagi Pengelolaan Sumber Daya dan Lingkungan Perairan*. Kanisius. Yogyakarta; 2003. (In Indonesian).
6. Adiwilaga EM, Hariyadi S. dan N. T. M. Pratiwi. *Perilaku oksigen terlarut selama 24 Jam Pada Lokasi Karamba Jaring*

- Apung di Waduk Saguling, Jawa Barat. *Limnotek*. 2009;16(2):109-118. (In Indonesian).
6. Barus TA. Pengantar Limnologi. USU Press. Medan; 2004. (In Indonesian).
 7. Boyd EC. Water quality: An introduction. Springer Science & Business Media. 2000;330.
 8. Reeb SG. Oxygen and fish behaviour. 2009;13. Available:www.howfishbehave.ca
 9. Lukman, Ridwansyah I. Kajian kondisi morfometri dan beberapa parameter stratifikasi perairan danau toba. *Jurnal Limnotek*. 2010;17(2):158-170. (In Indonesian).
 10. Republic of Indonesia Government Regulation No. 82 of 2001. Water quality mangement and water pollution control.
 11. Bhatnagar A, Devi P. Water quality guidelines for management of pond fish culture. *International Journal of Environmental Science*. 2013;3(6):1980-2009.
 12. Araoye PA. The seasonal variation of pH and dissolved oxygen (DO) concentration in Asa Lake Ilorin, Nigeria. *International Journal of Physical Science*. 2009;4(5):271-274.
 13. Zahidah. Produktivitas Perairan. Unpad Press. 2017;105. (In Indonesian)
 14. Yuliana, Adiwilaga EM, Harris E, Pratiwi NTM. Hubungan antara kelimpahan fitoplankton dengan parameter fisik-kimiawi perairan di teluk jakarta. *Jurnal Akuatika*. 2012;3(20):169-179. (In Indonesian)
 15. Kumar R, Grover AS, Wats M. Assessment of water quality status of lakes in Haryana, India. *Int J Recent Sci Res*. 2018;9(7):27831-27835.
 16. Ndungu JN. Assessing water quality in Lake Naivasha. Dissertation at Twente. Netherlands; 2014.

© 2019 Hasan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/50649>